

## APPENDIX 7

### References

- Abrams, M.D. (1992). Fire and the development of oak forests in eastern North America. *BioScience*, 42, 346-352.
- Adams, S.M., & Hackney, C.T. (1992). Ecological processes in southeastern United States aquatic ecosystems. In C. T. Hackney, S. M. Adams, & W. H. Martin (Eds.), *Biodiversity of the Southeastern United States: Aquatic Communities*. New York: John Wiley & Sons, Inc.
- Adams, W. J., Toll, J. E., Brix, K. V., Tear, L. M., & DeForest, D. K. (2000). Site-specific approach for setting water quality criteria for Selenium: differences between lotic and lentic systems. *Williams Lake, BC. Proceedings of Mine Reclamation Symposium; 2000 Jun 19-22*, 231-241.
- Ahern, M. M., Mullett, M., MacKay, K., & Hamilton, C. (2010). Residence in Coal-Mining Areas and Low-Birth-Weight Outcomes. *Maternal and Child Health Journal*, DOI 10.1007/s10995-009-0555-1.
- Alexander, R. B., Bohlke, J.K., Boyer, E.W., David, M.B., Harvey, J.W., Mulholland, P.J., Seitzinger, S.P., Tobias, C.R., Tonitto, C., & Wollheim, W.M.. (2009). Dynamic modeling of nitrogen losses in river networks unravels the coupled effects of hydrological and biogeochemical processes. *Biogeochemistry*, 93, 91–116.
- American Bird Conservancy. (2009.) Mountaintop Removal/Valley Fill Coal Mining Impacts on Birds. Retrieved December 29, 2009 from: <http://www.abcbirds.org/conservationissues/threats/energyproduction/mountaintop.html>.
- Amichev, B.Y., Burger, J. A., & Rodrigue, J. A. (2008). Carbon sequestration by forests and soils on mined land in the Midwestern and Appalachian coalfields of the US. *Forest Ecology Management* 256, 1949-1959.
- Andrle, R. F., & Carroll, J. R. (1988.) The atlas of breeding birds in New York State. Ithaca: Cornell University Press.
- Anderson, M. G., Merrill, M. D., & Biasi, F. D. (1998). Connecticut River Watershed Analysis: Ecological communities and neo-tropical migratory birds. *Final report summary to USGS Biological Resources Division*. The Nature Conservancy, Boston MA. 36pp.
- Appalachian Mountains Joint Venture (AMJV). (2007). Implementation plan for the Appalachian Mountains Joint Venture: a foundation for all-bird conservation in the region. 87 pp. <http://www.amjv.org/>
- Armstead, M. Y., Yaeger-Seagle, J., & Emerson, L. (2004). Benthic macroinvertebrate studies conducted in mountaintop mining/valley fill influenced streams in conjunction with the USEPA Environmental Impact Study. *Presented at the 2004 National Meeting of the American Society of Mining and Reclamation and the 25th West Virginia Surface Mine Drainage Task Force*, April 18-24, 2004. Lexington: ASMR.
- Armstead, M. Y. (2006). Status of the Industry Understanding of the Effects of Mountaintop Mining and Valley Fills on Aquatic Resources. *Report prepared by Potesta & Associates, Inc. for Jackson & Kelly, PLLC, Charleston, WV*.
- Arcott, D. B., Jackson, J. K., & Kratzer, E. B. (2006). Role of rarity and taxonomic resolution in a regional and spatial analysis of stream macroinvertebrates. *Journal of the North American Benthological Society*, 25, 977–997.
- Askins, R. A. (1993). Population trends in grassland, shrubland, and forest birds in eastern North America. *Current Ornithology* 11, 1-34.
- Arthur, M.A., G.B. Coltharp, & D.L. Brown (1998). Effects of best management practices on forest streamwater quality in eastern Kentucky. *J. Amer. Wat. Resour. Assoc.* 34:481-495.
- Bailey, B. (1999). Social and economic impacts of wild harvested products. *Ph. D. Dissertation, West Virginia University. Morgantown, WV*.

- Baker, J. W., Grover, J. P., Ramachandranair, R., Black, C., Valenti Jr., T. W., Brooks, B. W., & Roelke, D. L. (2009). Growth at the edge of the niche: an experimental study of the harmful alga *Prymnesium parvum*. *Limnology and Oceanography* 54,1679-1687.
- Baker, J.W., Grover, J. P., Brooks, B. W., Urena-Boeck, F., Roelke, D. L., Errera, R. (2007). Growth and toxicity of *Prymnesium parvum* (haptophyta) as a function of salinity light and temperature. *Journal of Phycology*, 43, 219–227.
- Barber-James, H. M., Gattolliat, J., Sartori, M., & Hubbard, M. D. (2008). Global diversity of mayflies (Ephemeroptera, Insecta) in freshwater. *Hydrobiologia*, 595, 339-350.
- Barbour, M. T., Gerritsen, J., Snyder, B. D., & Stribling, J. B. (1999). Rapid Bioassessment Protocols for use in streams and wadeable rivers: periphyton, benthic macroinvertebrates and fish, second edition. *EPA 841-B-99-002. U.S. Environmental Protection Agency; Office of Water; Washington, D.C. <http://www.epa.gov/owow/monitoring/rbp/>*
- Barbour, R. W., & Davis, W. H. (1974). Mammals of Kentucky. *Lexington: University of Kentucky Press.*
- Barrett, J. W. (1995). Regional silviculture of the United States. *New York: John Wiley & Sons.*
- Bauernfeind, E., & Moog O. (2000). Mayflies (Insecta: Ephemeroptera) and the assessment of ecological integrity: a methodological approach. *Hydrobiologia*, 422/423, 71-83.
- Baxter, C.V., Fausch, K.D., & Saunders, W.C. (2005). Tangled webs: reciprocal flows of invertebrate prey link streams and riparian zones, *Freshwater Biology* 50, 201–220
- Beketov, M. A. (2004). Different sensitivities of mayflies (Insecta, Ephemeroptera) to ammonia, nitrite and nitrate: linkage between experimental and observational data. *Hydrobiologia*, 528, 209-216.
- Bernhardt, E.S., Brian D. Lutz, B.D., Carter, C.E., Fay, J.P., King, R.S., Margaret A. Palmer, M.A., David Campagna, D., & Amos, J. (2010). How many mountains can we mine? Ecological thresholds for freshwater ecosystems of the Central Appalachians. *In Review, Copyrighted.*
- Bernhardt, E. S., & Palmer, M. A. (2007). Restoring streams in an urbanizing world. *Freshwater Biology*, 52, 738–751.
- Bernhardt, E. S., Sudduth, E.B., Palmer, M.A., Allan, J.D., Meyer, J.L., Alexander, G., Follstad-Shah, J., Hassett, B., Jenkinson, R., Lave R., Rumps J., Pagano, L. (2007). Restoring rivers one reach at a time: Results from a survey of U.S. river restoration practitioners. *Restoration Ecology*, 15, 482–493.
- Bevans, H., Parsons, S., Ziemkiewicz, P., and Winters, B. (2005). *Prediction and Treatment of Selenium – A framework for selenium Studies in the Appalachian Plateaus. Topic WQ-4. Accessed at <http://aciddrainage.com/ps/wq-4.cfm>*
- Bevier, L. R. (1994). The atlas of breeding birds of Connecticut. *State Geologic and Natural History Survey of Connecticut. Bulletin 113.*
- Bishop, K., Buffam, I., Erlandsson, M., Fölster, J., Laudon, H., Seibert, J., & Temnerud, J. (2008) *Aqua Incognita: the unknown headwaters. Hydrological Processes*, 22, 1239–1242.
- Bonta, J. V., & Dick, W. A. (2003). Impact of coal surface mining and reclamation on surface water chemical concentrations and load rates in three Ohio watersheds. *Journal of the American Water Resources Association*, 39(4), 793-816.
- Bormann, F. H. & Likens, G. E. (1979) *Pattern and Process in a Forested Ecosystem, New York, Springer-Verlag.*
- Boulton, A. J. (1989). Over-summering refugees of aquatic macroinvertebrates in two intermittent streams in Central Victoria. *Transactions of the Royal Society of South Australia*, 113, 23-34.

- Bray, J. R., & Gorham, E. (1964). Litter production in forests of the world. *Advanced Ecological Restoration*, 2, 101-157.
- Brewer, R., McPeck, G. A., & Adams, Jr. R. J. (1991). The atlas of breeding birds of Michigan. *East Lansing: Michigan State University Press*.
- Brinkman, S. F., & Johnston, W. D. (2008). Acute toxicity of aqueous copper, cadmium, and zinc to the mayfly *Rhithrogena hageni*. *Archives of Environmental Contamination and Toxicology*, 54, 466-472.
- Brittain, J. E., & Saltveit, S. J. (1989). Review of effect of river regulation on mayflies (Ephemeroptera). *Regulated Rivers Research and Management*, 3, 191-204. <http://onlinelibrary.wiley.com/doi/10.1002/rrr.3450030119/abstract>
- Brooks, R. T. (2009). Potential impacts of global climate change on the hydrology and ecology of ephemeral freshwater systems of the forests of the northeastern United States. *Climatic Change*, (95), 469-483.
- Brown, P., Cabarle B., & Livernash R.. (1997). Carbon Counts: Estimating Climate Change Mitigation in Forestry Projects. *Washington, DC: World Resources Institute*.
- Bryant, G., McPhilliamy, S., & Childers H. (2002). A survey of the water quality of streams in the primary region of mountaintop/valley fill coal mining October 1999 to January 2001. *In U.S. Environmental Protection Agency, Mid Atlantic Region, Mountaintop Mining Environmental Impact Statement, Appendix D, Stream chemistry final report, 102 p.* <http://www.epa.gov/region03/mtntop/eis2005.htm>
- Bryce, S. A., Omernik J. M., & Larsen D. P. (1999). Ecoregions - a geographic framework to guide risk characterization and ecosystem management. *Environmental Practice* 1(3),141-155.
- Buckelew Jr., A. R., & Hall, G. A. (1994). *The West Virginia Breeding Bird Atlas*. Pittsburgh: University of Pittsburgh Press.
- Buehler, D. A., Giocomo, J.J., Jones, J., Hamel, P.B., Rogers, C.M., Beachy, T.A., Varble, D.W., Nicholson, C.P., Roth, K.L., Barg, J., Robertson, R.J., Robb, J.R., Islam, K.. (2008). Cerulean warbler reproduction, survival, and models of population decline. *Journal of Wildlife Management*, 72, 646-653.
- Buffington, J. M., Kilgo, J.C., Sargent, R. A., Miller, K. V., & Chapman, B. R. (1997). Comparison of breeding bird communities in bottomland hardwood forests of different successional stages. *Wilson Bulletin*, 109(2), 314-319.
- Bulluck, L. P. (2007). *Golden-winged warbler (Vermivora chrysoptera) demographics and habitat use and the potential effects of land use change on golden-winged and cerulean warblers (Dendroica cerulea) in the Cumberland Mountains of Tennessee*. Ph.D. Dissertation, University of Tennessee, Knoxville. 200 pp.
- Burton, T. M. (1976). An analysis of the feeding ecology of the salamanders (Amphibia, Urodela) of the Hubbard Brook experimental forest, New Hampshire. *Journal of Herpetology*, 10, 187-284.
- Burton, T. M., & Likens, G. E. (1975). Salamander populations and biomass in Hubbard Brook Experimental Forest, New Hampshire. *Copeia*, 1975, 541-546.
- Buss, D. F., & Salles, F. F. (2007). Using Baetidae species as biological indicators of environmental degradation in a Brazilian River basin. *Environmental Monitoring and Assessment*, 130, 365-372.
- Canterbury, R.A. (2002). *Mountaintop removal and valley-fill mining environmental impact study: bird populations along edges*. Draft Report 2001. Final 2002. Concord College, Appendix in Mountaintop mining/valley fills in Appalachia. Final programmatic environmental impact statement. US Environmental Protection Agency, Philadelphia, Pennsylvania. <http://www.epa.gov/region03/mtntop/eis2003appendices.htm#appd>

- Carlisle, D.M. & Clements, W.H.. (1999). Sensitivity and variability of metrics used in biological assessments of running waters. *Environmental Toxicology and Chemistry* 18, 285-291.
- Caruso, B. S. & Haynes, J. (2010). Connectivity and jurisdictional Issues for Rocky Mountains and Great Plains aquatic resources. *Wetlands*. DOI 10.1007/s13157-010-0084-0
- Casas, J. J., Zamora-Munoz, C., Archila, F., & Alba-Tercedor, J. (2000). The effect of a headwater dam on the use of leaf bugs by invertebrate communities. *Regulated Rivers Research and Management*, 16, 577-591.
- Center for Biological Diversity (CBD). (2010). Petition to list the eastern-small footed bat (*Myotis leibii*) and northern long-eared bat (*Myotis septentrionalis*) as threatened or endangered under the Endangered Species Act. Center for Biological Diversity, Richmond, VT. 61 pp.
- Chadwick, M. A., Hunter, H., Feminella, J. W., & Henry, R. P. (2002). Salt and water balance in *Hexagenia limbata* (Ephemeroptera: Ephemeridae) when exposed to brackish water. *Florida Entomologist*, 85(4), 650-651.
- Chambers, D. B., & Messinger, T. (2001). Benthic invertebrate communities and their responses to selected environmental factors in the Kanawha River Basin, West Virginia, Virginia, and North Carolina. *Water-Resources Investigations Report 01-4021*, Department of the Interior, U.S. Geological Survey, Charlestown, WV. <http://pubs.usgs.gov/wri/wri014021/>
- Chapman, P. M., Adams, W.J., Brooks, M.L., Delos, C.G., Luoma, S.N., Maher, W.A., Ohlendorf, H.M., Presser, T.S., & ShawD.P. (2009). Ecological assessment of selenium in the aquatic environment: Summary of a SETAC Pellston Workshop. *Pensacola: Society of Environmental Toxicology and Chemistry (SETAC)*. <http://www.setac.org/sites/default/files/SELSummary.pdf>
- Chapman, P. M., Adams, W.J., Brooks, M.L., Delos, C.G., Luoma, S.N., Maher, W.A., Ohlendorf, H.M., Presser, T.S., & ShawD.P. (Ed.). (2010). Ecological Assessment of Selenium in the Aquatic Environment. *CRC Press, Taylor & Francis Group*
- Clarke, A., Mac Nally, R. Bond, N., & Lake, P. S. (2008). Macroinvertebrate diversity in headwater streams: a review. *Freshwater Biology*, 53, 1707–1721.
- Clarke, R. T., Wright, J. F., & Furse, M. T. (2003). RIVPACS models for predicting the expected macroinvertebrate fauna and assessing the ecological quality of rivers. *Ecological Modeling* 160, 219-233.
- Clawson, R. L. (2002). Trends in population size and current status. In *The Indiana bat: biology and management of an endangered species: 2–8*. Kurta, A. & Kennedy, J. (Eds). Austin, TX: Bat Conservation International.
- Clements, W. H. (1994). Benthic invertebrate community responses to heavy metals in the upper Arkansas River Basin, Colorado. *Journal of the North American Benthological Society*, 19, 30-44.
- Clements, W. H. (2004). Small-scale experiments support causal relationships between metal contamination and macroinvertebrate community response. *Ecological Applications*, 14, 954-967.
- Clements, W. H., Cherry, D. S., & Van Hassel, J. H. (1992). Assessment of the impact of heavy metals on benthic communities at the Clinch River (Virginia): evaluation of an index of community sensitivity. *Canadian Journal of Fisheries and Aquatic Science*, 49, 1686-169.
- Coal River TMDL Main Report. Total Maximum Daily Loads for Selected Streams in the Coal River Watershed, West Virginia. Final Approved Report, September 2006.

- Collins, B.M., Sobczka, W.V., & Colburn, E.A., . (2007). Subsurface flowpaths in a forested headwater stream harbor a diverse macroinvertebrate community. *Wetlands*, 27(2).
- Conant, R., & Collins, J. T. (1991). *A Field Guide to Reptiles and Amphibians: Eastern and Central North America*, 3<sup>rd</sup> Ed. Boston: Houghton Mifflin.
- Conley, J. M., Funk, D. H., & Buchwalter, D. B. (2009). Selenium bioaccumulation and maternal transfer in the mayfly *Centroptilum triangulifer* in a life-cycle, periphyton-biofilm trophic assay. *Environmental Science & Technology*, 43, 7952-7957.
- Courtney, L. A., & Clements, W. H. (2000). Sensitivity to acidic pH in benthic invertebrate assemblages with different histories of metal exposure. *Journal of the North American Benthological Society*, 19, 112-127.
- Courtney, L. A., & Clements, W. H. (2002). Assessing the influence of water and substratum quality on benthic macroinvertebrate communities in a metal-polluted stream: an experimental approach. *Freshwater Biology*, 47, 1766-1778.
- Cummins, K. W. (1973). Trophic relations of aquatic insects. *Annual Review of Entomology* 18, 183-206.
- Cummins, K. W. (1980). The natural stream ecosystem, pp. 7-24. In J.V. Ward and J.A. Stanford (Eds). *The ecology of regulated streams*. New York: Plenum Press.
- Cummins, K. W., & Klug, M. J. (1979). Feeding ecology of stream invertebrates. *Annual Review of Ecology, Evolution, and Systematics*, 10, 147-172.
- Cutter, G.A., & Bruland, K.W. (1984). The marine biogeochemistry of selenium: A re-evaluation, *Limnol. Oceanogr.*, 29(6), 1984, 1179-1192
- Daugherty, C. "Priority Delivery: 2008WVImpairedStreams.zzz". E-mail message to Christine Mazzarella. 30 July, 2009.
- Davic, R.D. and Welsh Jr., H.H. (2004) On the Ecological Roles of Salamanders. *Annual Review of Ecology, Evolution and Systematics*, 35: 405-434.
- DeBruyn, A., Chapman, M. H., & Chapman, P. M. (2007). Selenium toxicity to invertebrates: will proposed thresholds for toxicity to fish and birds also protect their prey? *Environmental Science & Technology*, 41, 1766-1770.
- Delcourt, P. A., & Delcourt, H. R. (1998). The influence of human-set fires on oak-chestnut forests in the southern Appalachians. *Castanea*, 63, 337-345.
- Delucchi, C. M. (1988). Comparison of community structure among streams with different temporal flow regimes. *Canadian Journal of Zoology*, 66, 579-586.
- Diamond, J., & Daley, C. (2000). What is the relationship between whole effluent toxicity results and biological condition? *Environmental Toxicology and Chemistry*, 19, 158-168.
- Diamond, J., Daley, C., & Moore, T. (1999). *Evaluating whole effluent toxicity testing as an indicator of instream biological conditions*. Alexandria (VA): Water Environment Research Foundation. 95-HHE-1. Available from [www.werf.org](http://www.werf.org)
- Dickens, P. S., Minear, R. A., & Tschantz, B. A. (1989). Hydrologic alteration of mountain watersheds from surface mining. *Research Journal of the Water Pollution Control Federation*, 61, 1249-1260.

- Diehl, S.F., Goldhaber, M.B., Koenig, A.E., Tuttle, M.L.W., & Ruppert, L.F. (2005) *Concentration of arsenic, selenium, and other trace elements in Appalachian coals of Alabama and Kentucky*. Presented at the 2005 National Meeting of the American Society of Mining and Reclamation, June 19-23, 2005. Published by ASMR, Lexington, KY.
- Doppelt, B. (1993). *Entering the Watershed: a New Approach to Save America's River Ecosystems*. Washington, D.C.: Island Press.
- Douglass, J.E. & Swank, W.T. (1975). Effects of management practices on water quality and quality: Coweeta Hydrologic Laboratory, North Carolina. *Northeastern Forest Experiment Station (USDA For. Serv. Gen. Tech. Rep. NE-13)*.
- Eaton, S. W. (1958). A life history study of the Louisiana Waterthrush. *Wilson Bulletin*, 70, 210-235. *Ecological Applications* 17(4): 1156-1167.
- Easton, R.S., Orth, D.J., & Voshell, J.R. (1996). Spatial and annual variation in the diets of juvenile smallmouth bass, *Micropterus dolomieu*, *Environmental Biology of Fishes* 46(4) 383-392
- Echols, B. S., Currie, R. J., & Cherry, D. S. (2009, November 4). Preliminary results of laboratory toxicity tests with the mayfly, *Isonychia bicolor* (Ephemeroptera: Isonychiidae) for development as a standard test organism for evaluating streams in the Appalachian coalfields of Virginia and West Virginia. *Environmental Monitoring and Assessment*. [Electronic version].
- Echternacht, A. C., & Harris, L. D. (1993). The fauna and wildlife of the southeastern United States. In W. H. Martin, G. Boyce & A. C. Echternacht (Eds.), *Biodiversity of the Southeastern United States, Volume 1*. (pp. 81-116). New York: John Wiley & Sons.
- Elmore, A. J., & Kaushal, S. S. (2008). Disappearing headwaters: patterns of stream burial due to urbanization. *Frontiers in Ecology and the Environment*, 6, DOI:10.1890/070101.
- Ensign, S. H., and Doyle, M.W. (2006). Nutrient spiraling in streams and river networks. *Journal of Geophysical Research*, Vol. 111, G04009.
- Ewers, R. M. and **Didham, R. K.** (2006). Confounding factors in the detection of species responses to habitat fragmentation. *Biological Reviews* **81**: 117-142.
- Fausch, K.D., J.R. Karr, and P.R. Yant. 1984. Regional application of and index of biotic integrity based on stream fish communities. *Transactions of the American Fisheries Society* 113: 39-55.
- Feminella, J. W. (1996). Comparison of benthic macroinvertebrate assemblages in small streams along a gradient of flow permanence. *Journal of the North American Benthological Society*, 15, 651-669.
- Ferrari, J. R., Lookingbill, T. R., McCormick, B., Townsend, P. A., & Eshelman, K. N. (2009). Surface mining and reclamation effects on flood response of watersheds in the central Appalachian Plateau region. *Water Resources Research* 45:W04407.
- Ferreri, C. P., Stauffer, J. R., & Stecko, T. D. (2004). *Evaluating impacts of mountain top removal/valley fill coal mining on stream fish populations*: Proceedings of a joint conference, 21st Annual Meeting of the American Society of Mining and Reclamation and 25th West Virginia Surface Mine Drainage Task Force, April 18-24, 2004. Morgantown, WV. Published by ASMR, p. 576-592.
- Février L., Martin-Garin, A., & Leclerc, E. (2007). Variation of the distribution coefficient (K<sub>d</sub>) of selenium in soils under various microbial states. *Journal of Environmental Radioactivity*, 97 (2-3), 189-205.
- Fischenich, J. C. (2006). *Functional Objectives for Stream Restoration*. EMRPP Technical Notes Collection. U.S. Army Corps of Engineers, Engineer Research and Development Center. Vicksburg, MS.

- Ford, C.R., & Vose, J.M. (2007). *Tsuga canadensis* (L.) Carr. mortality will impact hydrologic processes in southern Appalachian forest ecosystems. *Ecological Applications*, 17(4), 1156–1167.
- Freeman M. C., Pringle, C. M., & Jackson, C. R. (2007). Hydrologic connectivity and the contribution of stream headwaters to ecological integrity at regional scales. *Journal of the American Water Resources Association*, 43, 5–14.
- Freund, J. G, & Petty, J. T. (2007). Response of fish and macroinvertebrate bioassessment indices to water chemistry in a mined Appalachian watershed. *Environmental Management*. DOI 10.1007/s00267-005-0116-3.
- Fritz, K. M., Fulton, S., Johnson, B. R., Barton, C. D., Jack, J. D., Word, D. A., & Burke, R. A. (2010). Structural and functional characteristics of natural and constructed channels draining a reclaimed mountaintop removal and valley fill coal mine. *Journal of the North American Benthological Society*, 29(2), 673–689.
- Fritz, K. M., Johnson, B. R., & Walters, D. W. (2006). Field Operations Manual for Assessing the Hydrological Permanence and Ecological Condition of Headwater Streams. EPA/600/R-06/126. U.S. Environmental Protection Agency, Office of Research and Development, Washington, D.C.  
<http://www.epa.gov/eerd/manual/headwater/HISSmanual-cover.pdf>
- Fritz, K. M., Johnson, B. R., & Walters, D. M. (2008). Physical indicators of hydrologic permanence in forested headwater streams. *Journal of the North American Benthological Society*, 27, 690-704.
- Fry, J. A., Coan, M. J., Homer, C. G., Meyer, D. K., & Wickham, J. D. (2009). *Completion of the National Land Cover Database (NLCD) 1992–2001 Land Cover Change Retrofit product*. U.S. Geological Survey Open-File Report 2008–1379, 18 p.
- Fukui, D., Murakami, M., Nakano, S., & Aoi, T. (2006). Effect of emergent aquatic insects on bat foraging in a riparian forest, *Journal of Animal Ecology* 75, 1252–1258
- Fulk, F., Autrey, B., Hutchens, J., Gerritsen, J., Burton, J., Cresswell, C., & Jessup, B. (2003). *Ecological assessment of streams in the coal mining region of West Virginia using data collected by the U.S. EPA and environmental consulting firms*. National Exposure Research Laboratory, US Environmental Protection Agency, Cincinnati, Ohio. Appendix in Mountaintop mining/valley fills in Appalachia. Final programmatic environmental impact statement. US Environmental Protection Agency, Philadelphia, Pennsylvania.  
<http://www.epa.gov/region03/mtntop/eis2003appendices.htm#appd>
- Gerritsen, J., Burton, J., & Barbour, M. T. (2000). *A stream condition index for West Virginia wadeable streams*. Tetra Tech, Inc., Owings Mills, MD.
- Gill, J. D. (1975). Comparison of seven forest types for game in West Virginia. *Journal of Wildlife Management*, 39, 762-768.
- Gingerich, G. A. (2009). *Quantifying changes in ecological function of headwater catchments following large-scale surface mining in southern West Virginia*. M.S. Thesis. West Virginia University, Morgantown WV.
- Goetsch, P. A., & Palmer, C. G. (1997). Salinity tolerances of selected macroinvertebrates of the Sabie River, Kruger National Park, South Africa. *Archives of Environmental Contamination and Toxicology*, 32, 32-41.
- Gosz, J. R., Holmes, R. T., Likens, G. E., & Bormann, F. H. (1978). The flow of energy in a forest ecosystem. *Scientific American*, 283, 92-102.
- Grant, E. H. C., Green, L. E., & Lowe, W. H. (2009). Salamander occupancy in headwater stream networks. *Freshwater Biology*, 54, 1370–1378.

- Green, J., & Passmore, M. (1999). *Field survey report: an estimate of perennial stream miles in the area of the 1997 proposed Hobet Mining Spruce No. 1 Mine*. US Environmental Protection Agency. Wheeling, WV.
- Green, J., Passmore, M., & Childers, H. (2000). *A survey of the condition of streams in the primary region of mountaintop mining/valley fill coal mining*. US Environmental Protection Agency. Wheeling, WV. Appendix in *Mountaintop mining/valley fills in Appalachia*. Final programmatic environmental impact statement. US Environmental Protection Agency, Philadelphia, Pennsylvania.  
<http://www.epa.gov/region03/mtntop/eis2003appendices.htm#appd>
- Green, N. B., & Pauley, T. K. (1987). *Amphibians and reptiles in West Virginia*. Pittsburgh: University of Pittsburgh Press.
- Guerold, F. (2000). Influence of taxonomic determination level on several community indices. *Water Research*, 34, 487–492.
- Hairston, Jr., N. G. (1987). Diapause as a predator avoidance adaptation. In W. C. Kerfoot and A. Sih (Eds.), *Predation: Direct and indirect impacts on aquatic communities* (p. 281-290.) New England.
- Hambright, K. D. (unpublished). Golden algae data results, “WV-PA-Summary-qPCR results.xls. 2009 samples collected by WVDEP, PADEP and USEPA and sent to Dr. David Hambright's laboratory.
- Hambright, K. D. (2010) *Prymnesium parvum* Growth studies using the Dunkard Creek isolate (WANA strain). Report submitted to: West Virginia Department of Environmental Protection Division of Water and Waste Management. Charleston, WV. Department of Zoology University of Oklahoma, Norman, OK.
- Hamel, P. B. (2000). Cerulean warbler (*Dendroica cerulea*), The birds of North America online. (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology. <http://bna.birds.cornell.edu/bna/species/511>
- Hamilton, S. J. (2003). Review of residue-based selenium toxicity thresholds for freshwater fish. *Ecotoxicology and Environmental Safety*, 56, 201–210.
- Handel, S. N. (2003). Mountaintop removal mining/valley fill environmental impact statement technical study project report for terrestrial studies: terrestrial plant (spring herbs, woody plants) populations of forested and reclaimed sites. *Draft Report 2001. Final 2003*. New Brunswick: Rutgers University. Appendix in *Mountaintop mining/valley fills in Appalachia*. Final programmatic environmental impact statement. US Environmental Protection Agency, Philadelphia, Pennsylvania. <http://www.epa.gov/region03/mtntop/eis2003appendices.htm#appd>
- Harding, J. S., Benfield, E. F., Bolstad, P. V., Helfman, G. S., & Jones E. B. D. III. (1998). Stream biodiversity: The ghost of land use past. *Proceedings of the National Academy of Sciences*, 95, 14843–14847.
- Harper, C. A., & Guynn Jr., D. C. (1999). Factors affecting salamander density and distribution in four forest types in the southern Appalachian mountains. *Forest Ecology and Management*, 114, 245-252.
- Harper, K.A., Macdonald, E., Burton, P.J., Chen, J., Brososke, K.D., Saunders, S.C., Euskirchen, E.S., Roberts, D., Jaiteh, M.S., & Esseen, P. (2005). Edge Influence on Forest Structure and Composition in Fragmented Landscapes. *Conservation Biology*, 19:3, 768-782.
- Harpole, D. N., & Haas, C. A. (1999). Effects of seven silvicultural treatments on terrestrial salamanders. *Forest Ecology and Management*, 114, 349–356.
- Harrison, R. L. (1992). Toward a theory of inter-refuge corridor design. *Conservation Biology*, 6, 293-295.
- Hartman, K. J., Kaller, M. D., Howell, J. W., & Sweka, J. A. (2005). How much do valley fills influence headwater streams? *Hydrobiologia*, 532, 91-102.



- Hassell, K. L., Kefford, B. J., & Nuggeoda, D. (2006). Sub-lethal and chronic salinity tolerances of three freshwater insects: *Cloeon* sp. and *Centropilum* sp. (Ephemeroptera: Baetidae) and *Chironomus* sp. (Diptera: Chironomidae). *Journal of Experimental Biology*, 209, 4024-4032.
- Hawkins, C. P., Norris, R. H., Hogue, J.N., & Feminella, J. W. (2000). Development and evaluation of predictive models for measuring the biological integrity of streams. *Ecological Applications*, 10, 1456-1477.
- Hawkins, C. P. (2006a). Maintaining and restoring the ecological integrity of freshwater ecosystems: refining biological assessments. *Ecological Applications*, 16, 1249-1250.
- Hawkins, C. P. (2006b). Quantifying biological integrity by taxonomic completeness: Its utility in regional and global assessments. *Ecological Applications*, 16(4), 1277-1294.
- Hawkins, C. P., Olson J. R., & Hill R. A. (2010). The reference condition: predicting benchmarks for ecological and water-quality assessments. *Journal of the North American Benthological Society*, 29(1), 312-343.
- Heinz, G. H., & Hoffman D. J. (1996). Comparison of the effects of seleno-L-methionine, seleno-DL-methionine, and selenized yeast on reproduction of mallards. *Environmental Pollution*, 91, 169-175.
- Heinz, G. H., & Hoffman, D. J. (1998). Methylmercury chloride and selenomethionine interactions on health and reproduction in mallards. *Environmental Toxicology and Chemistry*, 17, 139-145.
- Heinz, G. H., Hoffman D. J., Krynitsky, A. J., & Weller, D. M. G. (1987). Reproduction in mallards fed selenium. *Environmental Toxicology and Chemistry*, 6, 423-433.
- Heinz, G. H., Hoffman D. J., and L. G. Gold. (1989). Impaired reproduction of mallards fed an organic form of selenium. *Journal of Wildlife Management*, 53, 418-428.
- Hendryx, M. (2009). Mortality from heart, respiratory, and kidney disease in coal mining areas of Appalachia. *International Archives of Occupational and Environmental Health*, 82, 243-249.
- Hendryx, M., & Ahern, M. M. (2008). Relations between health indicators and residential proximity to coal mining in West Virginia. *American Journal of Public Health*, 98(4), 669-670.
- Hendryx, M., & Ahern, M. M. (2009). Mortality in Appalachian coal mining regions: The value of statistical life lost. *Public Health Reports*, 124, 541-550.
- Hendryx, M., Ahern, M. M., & Nurkiewicz, T. R. (2007). Hospitalization patterns associated with Appalachian coal mining. *Journal of Toxicology and Environmental Health, Part A*, 70, 2064-2070.
- Hendryx, M., Fedorko, E., & Anesetti-Rothermel, A. (2010). A geographical information system-based analysis of cancer mortality and population exposure to coal mining activities in West Virginia, United States of America. *Geospatial Health*, 4(2), 243-256.
- Hendryx, M., O'Donnell, K., & Horn, K. (2008). Lung cancer mortality is elevated in coal-mining areas of Appalachia. *Lung Cancer*, 62, 1-7.
- Hendryx, M., & Zullig, K. J. (2009). Higher coronary heart disease and heart attack morbidity in Appalachian coal mining regions. *Journal of Preventive Medicine*, 49(5), 355-359.
- Herkert, J. R., Szafoni, R. E., Kleen, V. M., & Schwegman, J. E. (1993). *Habitat establishment, enhancement and management for forest and grassland birds in Illinois*. Natural Heritage Technical Publication #1. Illinois Department of Conservation, Division of Natural Heritage.  
<http://www.npwrc.usgs.gov/resource/birds/manbook/index.htm>

- Hinkle, C. R., McComb, W. C., & Marcus Jr., S. J. (1993.) Mixed Mesophytic Forest. pp. 203-254, 373, In W. H. Martin (Ed.). *Biodiversity of the Southeastern United States: Upland Terrestrial Communities*. New York: John Wiley & Sons, Inc.
- Hitt, N. P., & Hendryx, M. (2010). Ecological integrity of streams related to human cancer mortality rates. *Ecohealth*, 7(1), 91-104.
- Holl, K. D. (2002). Long-term vegetation recovery on reclaimed coal surface mines in the eastern USA. *Journal of Applied Ecology*, 39, 960-970.
- Hornick, L. E. (1981). Periphyton production in an Appalachian mountain trout stream. *American Midland Naturalist*, 106, 22-36.
- Howard, H. S., Berrang B., Flexner M., Pond G., & Call S. (2001). *Kentucky mountaintop mining benthic macroinvertebrate survey*. US Environmental Protection Agency. Athens, GA. Appendix in Mountaintop mining/valley fills in Appalachia. Final programmatic environmental impact statement. US Environmental Protection Agency, Philadelphia, Pennsylvania. <http://www.epa.gov/region03/mnttop/eis2003appendices.htm#appd>
- Hu, M., Yang, Y., Martin, J.-M., Yin, K. & Harrison, P.J. (1997). Preferential Uptake of Se(IV) Over Se(VI) and the Production of Dissolved Organic Se by Marine Phytoplankton, *Marine Environmental Research*. Vol. 44. No. 2, pp. 225-231.
- Hufford, M. (2003.) Reclaiming the Commons: Narratives of Progress, Preservation, and Ginseng. In B. J. Howell, (Ed.), *Culture, Environment, and Conservation in the Appalachian South*, Urbana: University of Illinois Press.
- Hynes, H. B. N. (1970). The Ecology of Running Waters. *Liverpool: Liverpool University Press*.
- Illinois Wildlife Habitat Commission. (1985). *The Crisis of Wildlife Habitat*. (p. 26) Illinois Department of Conservation.
- Intergovernmental Panel on Climate Change (IPCC). (2001). IPCC special report on land use, land-use change and forestry. [http://www.grida.no/publications/other/ipcc\\_sr/?src=/climate/ipcc/land\\_use/index.htm](http://www.grida.no/publications/other/ipcc_sr/?src=/climate/ipcc/land_use/index.htm)
- James, F. C., McCulloch, C. E., & Wiedenfeld, D. A. (1996). New approaches to the analysis of population trends in land birds. *Ecology* 77, 13-27.
- Jardine T.D. & Kidd K.A. (2010). Low concentrations of selenium in stream food webs of eastern Canada. *Science of the Total Environment*. DOI:10.1016/j.scitotenv.2010.11.013
- Jeffres, C. A., Opperman, J. J., & Moyle, P. B. (2008). Ephemeral floodplain habitats provide best growth conditions for juvenile Chinook salmon in a California river. *Environmental Biology of Fishes*, 83, 449–458.
- Johnson, B. R., Fritz, K. M., Blocksom, K. A., & Walters, D. M. (2009). Larval salamanders and channel geomorphology are indicators of hydrologic permanence in forested headwater streams. *Ecological Indicators*, 9, 150-159.
- Johnson, B. R., Haas, A., & Fritz, K. M. (2010). Use of spatially explicit physicochemical data to measure downstream impacts of headwater stream disturbance. *Water Resources Research*, 46. W09526, DOI:10.1029/2009WR008417.
- Jones, K. B. (1997). *An ecological assessment of the United States mid-Atlantic Region: a landscape atlas*. U.S. EPA. EPA/600/R-97/130. Washington, D.C.
- Jorgensen, C. B. (1966). *Biology of Suspension Feeding*. Pergamon: Oxford.

- Kaplan, D.I., & Serne, R. J. (1995). *Distribution coefficient values describing iodine, neptunium, selenium, technetium, and uranium sorption to Hanford sediments*. Prepared for the U.S. Department of Energy Pacific Northwest Laboratory, Richland, Washington. DOI 10.2172/158466
- Kefford, B. E. (1998). The relationship between electrical conductivity and selected macroinvertebrate communities in four riversystems of south-west Victoria, Australia. *International Journal of Salt Lake Research*, 7, 153–170.
- Kefford, B. E. (1998). Is salinity the only water quality parameter affected when saline water is disposed in rivers? *International Journal of Salt Lake Research*, 7, 285–300.
- Kefford, B. J., Papas, P. J., & Nuggeoda, D. (2003). Relative salinity tolerance of macroinvertebrates from the Barwon River, Victoria, Australia. *Marine and Freshwater Research*, 54, 755-765.
- Kefford, B. J., Papas, P. J., Metzeling, L., & Nuggeoda, D. (2004). Do laboratory salinity tolerances of freshwater animals correspond with their field salinity? *Environmental Pollution* 129, 355–362.
- Kefford, B. J., Dalton, A., Palmer, C. G., & Nuggeoda, D. (2004). The salinity tolerance of eggs and hatchlings of selected aquatic macroinvertebrates in south-east Australia and South Africa. *Hydrobiologia*, 517, 179-192.
- Kefford, B. J., Palmer, C.G., Jooste, S., Warne, M.St.J., Nuggeoda, D. (2005). What is meant by “95% of species”? An argument for the inclusion of rapid tolerance testing. *Human and Ecological Risk Assessment*, 11, 1025-1046.
- Kennedy, A. J., Cherry, D. S., & Currie, R. J. (2003). Field and laboratory assessment of a coal processing effluent in the Leading Creek watershed, Meigs County, Ohio. *Archives of Environmental Contamination and Toxicology*, 44, 324-331.
- Kennedy, A. J., Cherry, D. S., & Currie, R. J. (2004). Evaluation of ecologically relevant bioassays for a lotic system impacted by a coal-mine effluent, using *Isonychia*. *Environmental Monitoring and Assessment*, 95, 37-55.
- Kennedy, A. J., Cherry, D. S., & Zipper, C. E. (2005). Evaluation of ionic contribution to the toxicity of a coal-mine effluent using *Ceriodaphnia dubia*. *Archives of Environmental Contamination and Toxicology* 49, 155-162.
- Kiffney, P. M., & Clements, W. H. (1996). Size-dependent response of macroinvertebrates to metals in experimental streams. *Environmental Toxicology and Chemistry*, 15, 1352-1356.
- Kirk, E. (1999a). An evaluation of the aquatic habitats provided by sediment control ponds and other aquatic enhancement structures located on mine permitted areas in southern West Virginia. *Conducted for Pen Coal Corporation, by R.E.I. Consultants, Inc. Beaver, WV. Appendix in Mountaintop mining/valley fills in Appalachia. Final programmatic environmental impact statement. US Environmental Protection Agency, Philadelphia, Pennsylvania.* <http://www.epa.gov/region03/mtntop/eis2003appendices.htm#appd>
- Kirk, E. (1999b). Benthic macroinvertebrate study of Honey Branch, its sediment control ponds, and its influence on the East Fork of Twelvepole Creek, conducted 10/08/09. *Conducted for Pen Coal Corporation, by R.E.I. Consultants, Inc. Beaver, WV.*
- Klemm, D. J., Blocksom, K.A., Fulk, F.A., Herlihy, A.T., Hughes, R.M., Kaufmann, P.R., Peck, D.V., Stoddard, J.L., Thoeny, W.T., Griffith, M.B., & Davis, W.S. (2003). Development and evaluation of a Macroinvertebrate Biotic Integrity Index (MIBI) for regional assessing Mid-Atlantic Highland streams. *Environmental Management* 31, 656-669.
- Knapp, S. M., Haas, C. A., Harpole, D. N., & Kirkpatrick, R. L. (2003). Initial effects of clearcutting and alternative silvicultural practices on terrestrial salamander abundance. *Conservation Biology*, 7(3), 752–762.
- Laughlin, S. B., & Kibbe, D. P. (1985). *The atlas of breeding birds of Vermont*. Hanover, NH: University Press of New England.

- Lemly, A. D. (1985). Ecological basis for regulating aquatic emissions from the power industry: the case with selenium. US Forest Service. *Ecotoxicology and Environmental Safety*, 5, 465-486.
- Lemly, A. D. (1992). *Guidelines for evaluating selenium data from aquatic monitoring and assessment studies*. US Forest Service. Virginia Tech University, Blacksburg, VA.
- Lemly, A. D. (1993). Guidelines for evaluating selenium data from aquatic monitoring and assessment. *Environmental Monitoring and Assessment*, 28, 83-100.
- Lemly, A. D. (1999). *Selenium transport and bioaccumulation in aquatic ecosystems: a proposal for water quality criteria based on hydrological units*. US Forest Service, Virginia Tech University, Blacksburg, VA.
- Lemly, A. D. (2002a). Selenium assessment in aquatic ecosystems: *a guide for hazard evaluation and water quality criteria*. New York: Springer.
- Lemly, A. D. (2002b). Symptoms and implications of selenium toxicity in fish: the Belews Lake case example. *Aquatic Toxicology*, 57, 39-49.
- Lemly, A. D. (2002c). Tracking the Pathways and Anticipated Impact of Selenium. *News Release, USDA Southern Research Station*.
- Lemly, A. D. (2004). Aquatic Selenium Pollution is a Global Environmental Safety Issue. *Ecotoxicology and Environmental Safety*, 59, 44-56.
- Lemly, A. D. (2007). A procedure for NEPA assessment of selenium hazards associated with mining. *Environmental Monitoring and Assessment*, 125, 361-375.
- Lemly, A.D. (2008). Declaration on Aquatic Hazard of Selenium Releases From Coal Mining in the Mud River Ecosystem, West Virginia. Filed in Case #3:08-cv-00088 (S.D. Wv.).
- Lemly, A. D. (2009). White paper on “Aquatic hazard of selenium pollution from mountaintop removal coal mining.” Submitted for the record in Senate hearing.
- Lenat, D. R., & Resh, V. H. (2001). Taxonomy and stream ecology—the benefits of genus- and species-level identifications. *Journal of the North American Benthological Society*, 20, 287-298.
- Leopold, L.B., Wolman, M.G., and Miller, J.P. (1964). *Fluvial Processes in Geomorphology*. W.H. Freeman and Company, San Francisco.
- Lotrich, V. A. (1973). Growth, production, and community composition of fishes inhabiting a first-, second-, and third-order stream of eastern Kentucky. *Ecological Monographs*, 43(3), 377-397.
- MacDonald, L. & Drew, C. (2007). Influence of headwater streams on downstream reaches in forested areas. *Forest Science*, 53(2) .
- Maggard, R., & Kirk, E. (1998). Downstream impacts of surface mining and valley fill construction. Report for Pen Coal Corporation. Appendix in *Mountaintop mining/valley fills in Appalachia. Final programmatic environmental impact statement*. US Environmental Protection Agency, Philadelphia, Pennsylvania. <http://www.epa.gov/region03/mtntop/eis2003appendices.htm#appd>
- Maier, K. J., & Knight, A. W. (1994). Ecotoxicology of selenium in freshwater ecosystems. *Reviews of Environmental Contamination & Toxicology*, 134, 31-48.
- Martin, W. H., Boyce, S. G., & Echternacht, A. C. (Eds.) (1973). Biodiversity of the Southeastern United States: Upland Terrestrial Communities. (pp. 1-34, 373) *New York: John Wiley & Sons, Inc.*

- Mattsson, B. J., & Cooper, R. J. (2006). Louisiana Waterthrushes (*Seiurus motacilla*) and habitat assessments as cost-effective indicators of instream biotic integrity. *Freshwater Biology*, 51(10), 1941-1958.
- Mattsson, B. J., & Cooper, R. J. (2009). Multiscale analysis of the effects of rainfall extremes on reproduction by an obligate riparian bird in urban and rural landscapes. *Auk*, 126(1), 64-76.
- Mattsson, B. J., Master, T. L., Mulvihill, R. S., & Robinson, W. D. (2009). Louisiana Waterthrush (*Seiurus motacilla*), In (A. Poole, (Ed.), *The birds of North America online*. Ithaca: Cornell Lab of Ornithology. <http://bna.birds.cornell.edu/bna/species/151>
- May, T.W., Walther, M.J., Petty, J.D., Fairchild, J.F., Lucero, J., Delvaux, M., Manring, J., Armbruster, M. & Hartman, D. (2001). An evaluation of selenium concentrations in water, sediment, invertebrates, and fish from the Republican River Basin: 1997-1999. *Environmental Monitoring and Assessment*, 72, 179-206.
- McComb, W. C., McGarigal, K., Fraser, J. D., & Davis, W. H. (1991). Planning for basin-level cumulative effects in the Appalachian coal field. In J. E. Rodiek & E. G. Bolen (Eds.), *Wildlife and habitats in managed landscapes*. Washington, D.C.: Island Press.
- McCormick, F. H., Hughes, R. M., Kaufmann, P. R., Peck, D. V., Stoddard, J. L., & Herlihy, A. T. (2001). Development of an Index of Biotic Integrity for the Mid-Atlantic Highlands Region. *Transactions of the American Fisheries Society*. 130, 857-877.
- Meffe, G. K., & Carroll, C. R., (Eds.). (1994). Principles of conservation biology. *Sunderland: Sinauer Associates*.
- Mengel, R. M. (1965). The birds of Kentucky. *American Ornithological Monograph* 3.
- Merriam, E. R. (2009). Mining and Residential Development Interact to Produce Highly Impaired Stream Conditions in an Intensively Mined Appalachian Watershed. *M.S. thesis. West Virginia University. Morgantown, WV*.
- Merriam, E., Merovich, G. T., & Petty, J. T. In Review. Additive effects of mining and residential development on stream conditions in an intensively mined Appalachian watershed. *M.S. thesis. West Virginia University. Morgantown, WV*.
- Merricks, T. C., Cherry, D. S., Zipper, C. E., Currie, R. J., & Valenti, T. W. (2007). Coal mine hollow fill and settling pond influences on headwater streams in southern West Virginia, USA. *Environmental Monitoring and Assessment* 129, 359-378.
- Merritt, J. F. (1987). Guide to the mammals of Pennsylvania. *Pittsburgh: University of Pittsburgh*.
- Merritt, R. W., Cummins, K. W., & Burton, T. M. (1984). The role of aquatic insects in the processing and cycling of nutrients. In V.H. Resh and D.M. Rosenberg (Eds.), *The ecology of aquatic insects*, (pp. 134-163.). *New York: Praeger Publishers*.
- Merritt, R. W., & Cummins, D. W. (Eds.) (1996). An introduction to the aquatic insects of North America 3<sup>rd</sup> Edition. *Dubuque: Kendall-Hunt*.
- Messinger, T. (2003). Comparison of storm response of streams in small, unmined and valley-filled watersheds, 1999–2001, Ballard Fork, West Virginia. *Water-Resources Investigations Report 02-4303, Department of the Interior, U.S. Geological Survey, Charleston, WV*.
- Messinger, T., & Paybins, K. S. (2003). Relations between precipitation and daily and monthly mean flows in gaged, unmined and valley-filled watersheds, Ballard Fork, West Virginia, 1999–2001. *Water-Resources Investigations Report 03-4113, Department of the Interior, U.S. Geological Survey, Charleston, WV*.

- Meyer J. L., Strayer, D. L., Wallace, J. B., Eggert, L., Helfman, G. S., & Leonard, N. E. (2007). The contribution of headwater streams to biodiversity in river networks. *Journal of the American Water Resources Association*, 43, 86–103.
- Michel, N., Desante, D. F., Kaschube, D. R., & Nott, M. P. (2006). The Monitoring Avian Productivity and Survivorship (MAPS) Program Annual Reports, 1989–2003. *NBII/MAPS Avian Demographics Query Interface*. [Online.] <http://www.birdpop.org/nbii/NBIIHome.asp>
- Moler, P. E., & Franz, R., (1987). Wildlife values of small, isolated wetlands in the Southeastern coastal plain. In R. R. Odom, K. A. Riddleberger, J.C. Ozier, (Eds.), *Proceedings of the Third Nongame and Endangered Wildlife Symposium*. Georgia Department of Natural Resources, Social Circle, GA.
- Moog, O., Bauernfeind, E., & Weichselbaumer, P. (1997). Use of Ephemeroptera as saprobic indicators in Austria. In P. Landolt and M. Sartori (Eds.), *Ephemeroptera and Plecoptera: biology-ecology-systematics* (pp. 254-260). Fribourg: MTL.
- Mordecai, R.S., Mattsson, B.J., Caleb J. Tzilkowski, C.J., & Robert J. Cooper, R.J. (2011). Addressing challenges when studying mobile or episodic species: hierarchical Bayes estimation of occupancy and use, *Journal of Applied Ecology*, 48, 56–66
- Morse, J. C., Stark, B. P., & McCafferty, W. P. (1993). Southern Appalachian streams at risk: implications for mayflies, stoneflies, caddisflies, and other aquatic biota. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 3, 293-303.
- Morse, J. C., Stark, B. P., McCafferty, W. P., & Tennessen, K. J. (1997). Southern Appalachian and other southeastern streams at risk: implications for mayflies, dragonflies, stoneflies, and caddisflies. In G.W. Benz and D. E. Collins (Eds). *Aquatic fauna in peril: the southeastern perspective* (pp.17-42, 554). *Special Publication 1, Southeastern Aquatic Research Institute*. Decatur, GA: Lenz Design and Communications.
- Mount, D. R., Gulley, D. D., Hockett, J. R., Garrison, T. D., & Evans, J. M. (1997). Statistical models to predict the toxicity of major ions to *Ceriodaphnia dubia*, *Daphnia magna*, and *Pimephales promelas* (fathead minnows). *Environmental Toxicology and Chemistry*, 16, 2009-2019.
- Mueller, A. J., Loesch, C. R., & Twedt, D. J. (1999). Development of management objectives for breeding birds in the Mississippi Alluvial Valley. *Proceedings of Partners in Flight International Workshop. 1-5 Oct 1995, Cape May, NJ*.
- Mulvihill, R. S. (1999). Effects of stream acidification on the breeding biology of an obligate riparian songbird, the Louisiana Waterthrush (*Seiurus motacilla*). In W. E. Sharpe & J. R. Drohan, (Eds.), *The effects of acidic desposition on aquatic ecosystems in Pennsylvania, Proceedings of the 1998 Pennsylvania Acidic Deposition Conference, Vol. 2* (pp. 51-61) University Park, PA: Environmental Resources Research Institute.
- Mulvihill, R.S. (2010). Personal communication via U.S. FWS
- Mulvihill, R. S., Newell, F. L., & Latta, S. C. (2008). Effects of acidification on the breeding ecology of a stream-dependent songbird, the Louisiana Waterthrush (*Seiurus motacilla*). *Freshwater Biology*, 53(11), 2158-2169.
- Murray, N. L., & Stauffer, D. F. (1995). Nongame bird use of habitat in central Appalachian riparian forests. *Journal of Wildlife Management*, 59, 78-88.
- Nadeau, T. L., & Rains, M. C. (2007). Hydrological connectivity between headwater streams and downstream waters: How science can inform policy. *Journal of the American Water Resource Association*, 43, 118-133.
- Nakano and Murakami 2001: Reciprocal subsidies: dynamic interdependence between terrestrial and aquatic food webs. *Proceedings of the National Academy of Science*, 98:166-170;

- Napier B.A., & Snyder, S.F. 2002. Recommendations for User Supplied Parameters for the RESRAD Computer Code for Application to the Hanford Reach National Monument. *PNNL-14041, Pacific Northwest National Laboratory, Richland, Washington.*
- National Drought Mitigation Center. US Drought Monitor. <http://www.drought.unl.edu/dm/>
- National Oceanic and Atmospheric Association (NOAA) Satellite and Information Services. West Virginia climate summary. <http://www.ncdc.noaa.gov/oa/climate/research/cag3/wv.html>
- National Oceanic and Atmospheric Association (NOAA) National Weather Service Climate Prediction Center. Drought monitoring. [http://www.cpc.noaa.gov/products/monitoring\\_and\\_data/drought.shtml](http://www.cpc.noaa.gov/products/monitoring_and_data/drought.shtml)
- NatureServe. (2010). Conservation Status of the Eastern Small-footed Myotis (*Myotis leibii*) as published in *NatureServe Explorer online*.  
[http://www.natureserve.org/explorer/servlet/NatureServe?sourceTemplate=tabular\\_report.wmt&loadTemplate=species\\_RptComprehensive.wmt&selectedReport=RptComprehensive.wmt&summaryView=tabular\\_report.wmt&elKey=106402&paging=home&save=true&startIndex=1&nextStartIndex=1&reset=false&offPageSelectedElKey=106402&offPageSelectedElType=species&offPageYesNo=true&post\\_processes=&radiobutton=radiobutton&selectdIndexes=106402](http://www.natureserve.org/explorer/servlet/NatureServe?sourceTemplate=tabular_report.wmt&loadTemplate=species_RptComprehensive.wmt&selectedReport=RptComprehensive.wmt&summaryView=tabular_report.wmt&elKey=106402&paging=home&save=true&startIndex=1&nextStartIndex=1&reset=false&offPageSelectedElKey=106402&offPageSelectedElType=species&offPageYesNo=true&post_processes=&radiobutton=radiobutton&selectdIndexes=106402)
- Neal, R. H., & G. Sposito, G. (1989). Selenate adsorption on alluvial soils. *Soil Sci. Soc. Am. J.* 53:70-74.
- Negley, T. L., & Eshleman, K. N. (2006). Comparison of stormflow responses of surface-mined and forested watersheds in the Appalachian Mountains, USA. *Hydrological Processes*, 20, 3467-3483.
- Neuzil, S.G., Dulong, F.T, Cecil, C.B., Fedorko, N.F., Renton, J.J., and Bhumbla, D.K. (2007). Selenium Concentrations in Middle Pennsylvanian Coal-Bearing Strata in the Central Appalachian Basin. *USGS Open-File Report 2007-1090*
- Neuzil, S. G., Dulong, F. D., & Cecil, C. B. (2005). Spatial trends in ash yield, sulfur, selenium, and other selected trace element concentrations in coal beds of the Appalachian Plateau Region, U.S.A. *Open-File Report 2005-1330 Department of the Interior, U.S. Geological Survey, Reston, VA.*
- North American Bird Conservation Initiative (NABCI). (2000). Bird Conservation Regions Map. *U.S. NABCI Committee, U.S. Fish and Wildlife Service, Division of Bird Habitat Conservation, Arlington, VA.*  
<http://www.nabci-us.org/map.html>
- Nickens, T. E. (2004). Small is beautiful. *National Wildlife*, 42, 28-31.
- North Carolina Department of Environment and Natural Resources. (2005). Identification methods for the origins of intermittent and perennial streams. Version 3.1. *Division of Water Quality. Raleigh, NC.*  
[http://h2o.enr.state.nc.us/ncwetlands/documents/NC\\_Stream\\_ID\\_Manual.pdf](http://h2o.enr.state.nc.us/ncwetlands/documents/NC_Stream_ID_Manual.pdf)
- Nowacki, G. J., & Abrams, M. D. (1991). Community and edaphic analysis of mixed oak forests in the ridge and valley province of central Pennsylvania. In L.H. McCormick and K.W. Gottschalk (Eds.), *8<sup>th</sup> Central Hardwood Forest Conference: Proceedings*, (pp. 247-260). *Pennsylvania State University.*
- O'Connell, T. J., Brooks, R. P., Laubscher, S. E., Mulvihill, R. S., & Master, T. L. (2003). *Using bioindicators to develop a calibrated index of regional ecological integrity for forested headwater ecosystems. Final Report to U.S. Environmental Protection Agency, STAR Grants Program. Penn State Cooperative Wetlands Center, Pennsylvania State University, University Park, PA.*
- O'Connell, T. J., Jackson, L. E., & Brooks, R. P. (1998). A bird community index of biotic integrity for the Mid-Atlantic highlands. *Environmental Monitoring and Assessment*, 51, 145-156.

- Ohio Environmental Protection Agency. (2002). Technical report: Ohio's primary headwater streams-macroinvertebrate assemblages. *Division of Surface Water, Columbus, OH*.  
[http://www.epa.ohio.gov/portals/35/wqs/headwaters/TechRep\\_Macroinvert\\_2002.pdf](http://www.epa.ohio.gov/portals/35/wqs/headwaters/TechRep_Macroinvert_2002.pdf)
- Ohio Environmental Protection Agency. (2009). Field evaluation manual for Ohio's primary headwater habitat streams, Version 2.3. *Division of Surface Water, Columbus, OH*.  
[http://www.epa.ohio.gov/portals/35/wqs/headwaters/PHWHManual\\_2009.pdf](http://www.epa.ohio.gov/portals/35/wqs/headwaters/PHWHManual_2009.pdf)
- Ohlendorf, H.M. (2003). Ecotoxicology of selenium. In D.J. Hoffman *et al.* (Eds.), *Handbook of Ecotoxicology*, 2<sup>nd</sup> Edition, (pp. 465-500). Boca Raton, FL: Lewis Publishers
- Ohlendorf, H. A., Hoffman, D. J., Rattner, B. A., Burton Jr., G. A., & Cairns Jr., J. (2003). Ecotoxicology of selenium. In *Handbook of Ecotoxicology, Second Edition*, (pp. 465-500). *Boca Ration: Lewis*.
- Orr, P. L., Guiguer, K. R., & Russel, C. K. (2005). Food chain transfer of selenium in lentic and lotic habitats of a western Canadian watershed. *Ecotoxicology and Environmental Safety*, 63, 175-188.
- Orwig, D. A., Cobb, R. C., D'Amato, A. W., Kizlinski, M. L., and Foster, D. R. (2008). Multi-year ecosystem response to hemlock woolly adelgid infestation in southern New England forests. *Canadian Journal of Forest Research- Revue Canadienne De Recherche Forestiere*, 38(4), 834-843.
- Owen, S.F., Menzel, M.A., Edwards, J.W., Ford, M.W., Menzel, J.M., Chapman, B.R., Wood, P.B., & Miller, K.V. (2004). Bat Activity in harvested and intact forest stands in the Allegheny mountains, *J. of Applied Forestry Vol. 21, No.3*
- Palmer, M. A., & Bernhardt, E. S. (2009). White paper on “*Mountaintop Mining Valley Fills and Aquatic Ecosystems: A Scientific Primer on Impacts and Mitigation Approaches*” Submitted for the record in Senate hearing.  
[www.http://epw.senate.gov/public/index.cfm?FuseAction=Files.View&FileStore\\_id=66fea6d0-9bce-4a9b-be47-aa264a471a89](http://epw.senate.gov/public/index.cfm?FuseAction=Files.View&FileStore_id=66fea6d0-9bce-4a9b-be47-aa264a471a89)
- Palmer, M. A., Bernhardt, E.S., Schlesinger, W.H., Eshleman, K.N., Foufoula-Georgiou, E., Hendryx, M.S., Lemly, A.D., Likens, G.E., Loucks, O.L., Power, M.E., White, P.S., Wilcock, P.R. (2010). Mountaintop mining consequences. *Science*, 327, 149-149.
- Palmer, M.A., Menninger, H.L., and E. Bernhardt, E.S. (2009) River restoration, habitat heterogeneity and biodiversity: a failure of theory or practice? *Freshwater Biology*, 55:1-18.
- Parker, C. R., Flint, O. S., Jacobus, L. M., Kondratieff, B. C., McCafferty, W. P., & Morse, J. C. (2007). Ephemeroptera, Plecoptera, Megaloptera, and Trichoptera of Great Smoky Mountains National Park. *Southeastern Naturalist Special Issue 1*, 159-174.
- Pate, R. L. (2008). Soil Survey of Logan and Mingo Counties, West Virginia. *USDA, Natural Resources Conservation Service*. 317 pp.
- Patnode, K., Kain, C., Ramsey, D., Rhodes, J., & Evans, B. (2005). Salamander assemblage survey of mercury and selenium contaminated headwater sites in the Appalachian Mountains of Pennsylvania, Virginia, and West Virginia. *USFWS Region 5 Environmental Contaminants Program Final Report*. 21 pp.
- Patrick, R. (1948). Factors effecting the distribution of diatoms. *Botanical Review*, 14, 473-524.
- Patton, P. W. C. (1994). The effect of edge on avian nest success: how strong is the evidence? *Conservation Biology*, 8, 17-26.
- Pauley, T. Personal communication. Marshall University



- Paybins, K. S. (2003). Flow origin, drainage area, and hydrologic characteristics for headwater streams in the mountaintop coal-mining region of southern West Virginia. *Water Resources Investigation Report 02-4300*. U.S. Geological Survey. Charleston, WV.
- Paybins, K. S., Messinger, T., Eychaner, J. H., Chambers, D. B., & Kozar, M. D. (2000). Water quality in the Kanawha-New River Basin. West Virginia, Virginia, and North Carolina, 1996-98: *U.S. Geological Survey Circular 1204*. 32 pp. <http://pubs.water.usgs.gov/circ1204/>
- Pennsylvania Department of Environmental Protection. (PADEP) (unpublished). Golden algae data, "Algae.xls." CONSOL & PADEP 2009 compilation by PADEP from PADEP and CONSOL data
- Peterjohn, B. G., Sauer, J. R., & Robbins, C. S. (1995). Population trends from the North American breeding bird survey. In T. E. Martin & D. M. Finch, (Eds.), *Ecology and Management of Neotropical Migratory Birds* (pp 3-39). New York: Oxford University Press.
- Peterman, W. E., Crawford, J. A., & Semlitsch, R. D. (2008). Productivity and significance of headwater streams: population structure and biomass of the black-bellied salamander (*Desmognathus quadramaculatus*). *Freshwater Biology* 53, 347–357.
- Peterson, S. A., Ralston, N. V., Peck, D.V., Van Sickle, J., Robertson, J. D., Spate, V. L., & Morris, J. S. (2009). How might selenium moderate the toxic effects of mercury in stream fish of the western U.S.? *Environmental Science and Technology*, 43, 3919–3925.
- Petranka, J. W. (1998). Salamanders of the United States and Canada. Washington, D.C.: Smithsonian Institution Press.
- Petranka, J. W., Eldridge, M. E., & Haley, K. E. (1993). Effects of timber harvesting on southern Appalachian salamanders. *Conservation Biology*, 7(2), 363-370.
- Petty, J. T., Fulton, J. B., Strager, M. P., Merovich, Jr., G. T., Stiles, J. M., & Ziemkiewicz, P. F. (2010). Landscape indicators and thresholds of stream ecological impairment in an intensively mined Appalachian watershed. *Journal of the North American Benthological Society*, 29(4), 1292–1309.
- Phillips, J. D. (2004). Impacts of surface mine valley fills on headwater floods in eastern Kentucky. *Environmental Geology*, 45, 367-380.
- Pond, G. J. (2004). Effects of surface mining and residential land use on headwater stream biotic integrity in the eastern Kentucky coalfield region. *Kentucky Department for Environmental Protection, Division of Water, Frankfort, KY*.
- Pond, G. J. (2010). Patterns of Ephemeroptera taxa loss in Appalachian headwater streams (Kentucky, USA). *Hydrobiologia DOI 10.1007/s10750-009-0081-6*.
- Pond, G. J., & Passmore, M. (2008). Issue Paper: Recommendations for determining Relatively Permanent Waters (RPWs) in headwater streams in the Huntington and Pittsburgh Corps Districts. July 19, 2007; Finalized May 19, 2008. US Environmental Protection Agency. Wheeling, WV.
- Pond, G. J., Passmore, M. E., Borsuk, F. A., Reynolds, L. & Rose, C. J. (2008). Downstream effects of mountaintop coal mining: comparing biological conditions using family- and genus-level macroinvertebrate bioassessment tools. *Journal of the North American Benthological Society*, 27 (3), 717-737.
- Pond, G. J., & McMurray, S. E. (2002). A macroinvertebrate bioassessment index for headwater streams in the eastern coalfield region, Kentucky. *Kentucky Department for Environmental Protection, Division of Water, Frankfort, KY*.
- Potesta & Associates, Inc. (2010). The status of selenium criteria development in mining influenced watersheds in West Virginia (Presenter: Mindy Armstead, Ph.D.). 2010 West Virginia Water Conference. West Virginia's Water Resources: Threats and Opportunities. October 6-7, 2010. Morgantown, WV.

- Pough, F. H., Smith, E. M., Rhodes, D. H. & Collazo, A. (1987). The abundance of salamanders in forest stands with different histories of disturbance. *Forest Ecology and Management*, 20, 1-9.
- Riedel, G.F., Ferrier, D.P., & Sanders, J.G. (1991). Uptake of Selenium by freshwater phytoplankton, *Water, Air, and Soil Pollut.* 57/58: 23-30
- Robbins, C. S. (1979). Effect of forest fragmentation on bird populations. In R. M. DeGraaf and K. E. Evans, (Comps.) Management of North Central and Northeastern Forests for Nongame Birds. *General Technical Report Nc-51* (pp. 198-212). USDA, Forest Service, North Central Forest Experiment Station, St. Paul, MN.
- Robbins, C. S., Sauer, J. R., Greenburg, R. S., & Droege, S. (1989). Population declines in North American birds that migrate to the neotropics. *Proceedings of the National Academy of Sciences*, 86, 7658-7662.
- Robbins, C. S. (1980). Effect of forest fragmentation on breeding bird populations in the piedmont of the Mid-Atlantic Region. *Atlantic Naturalist*, 33, 31-36.
- Robbins, C. S., Fitzpatrick, J. W. & Hamel, P. B. (1992). A warbler in trouble: *Dendroica cerulea*. In J. M. Hagan III, & D. W. Johnston (Eds.), *Ecology and conservation of neotropical migrant landbirds*. (pp. 549-562). Washington, D.C.: Smithsonian Institution Press.
- Robinson, W. D. (1990). Louisiana Waterthrush foraging behavior and microhabitat selection in southern Illinois. *Master's Thesis, Southern Illinois University, Carbondale, IL.*
- Rocco, G. L., Brooks, R. P., & Hite, J. T. (2004). Stream Plethodontid Assemblage Response (SPAR) index: development, application, and verification in the MAHA. *Penn State Cooperative Wetlands Center, Pennsylvania State University Report No. 2004-1 prepared for US EPA Star Grant Program.*
- Rodgers, J. H., Johnson, B. M., & Bishop, W. M. (In Press.) Comparison of three algaecides for controlling the density of *Prymnesium parvum*. *Journal of the American Water Resources Association Golden Algae Featured Collection.*
- Roelke, D. L., Grover, J.P., Brooks, B.W., Glass, J., Buzan, D., Southard, G.M., Fries, L., Gable, G.M., Schwierzke-Wade, L., Byrd, M., Nelson, J. (2010). A decade of fish-killing *Prymnesium parvum* blooms in Texas: roles of inflow and salinity. *Journal of Plankton Research* DOI: 10.1093/plankt/fbq079, available online at [www.plankt.oxfordjournals.org](http://www.plankt.oxfordjournals.org). First published online: July 6, 2010
- Rosenberg, K.V., Barker, S.E., and Rohrbaugh, R.W. (2000). *An Atlas of Cerulean Warbler Populations. Final Report to the U.S. Fish and Wildlife Service: 1997-2000 Breeding Seasons.* Cornell Lab of Ornithology, Ithaca, NY 14850.
- Rosenberg, K. V., & Wells, J. V. (2002). Global perspectives on neotropical migratory bird conservation in the northeast: Long-term responsibility vs. immediate concern. [http://birds.cornell.edu/pifcapemay/rosenberg\\_wells.htm](http://birds.cornell.edu/pifcapemay/rosenberg_wells.htm).
- Ross, R. M., Redell, L. A., Bennett, R. M., & Young, J. A. (2004). Mesohabitat use of threatened hemlock forests by breeding birds of the Delaware River Basin in northeastern United States. *Natural Areas Journal*, 24(4), 307-315.
- Sager, D. R., Barkoh, A, Buzan, D.L., Fries, L.T., Glass, J.A., Kurten, G.L., Ralph, J.J., Singhurst, E.J., Southard, G.M., and Swanson, E. (2008). Toxic *Prymnesium parvum*: A potential threat to U.S. reservoirs. *American Fisheries Society Symposium*, 62, 261-273.
- Samuel, D. E., & Whitmore R. C. (1979). Reclamation and management of surface mines for game and non-game birds. pp. 578-582 in J. Luchok, J. D. Cawthon and M. J. Breslin (Eds.). *Hill Lands: proceedings of an international symposium held in Morgantown WV, 3-9 October 1976.* West Virginia University, Morgantown, WV.

- Sauer, J. R. (1993). Monitoring goals and programs of the U.S. Fish and Wildlife Service. In D.M. Finch & P. W. Stangel (Eds.), *Status and management of neotropical migratory birds; 21-25 Sep 1992*(pp. 245-251). Estes Park, CO, U.S. Department of Agriculture, Forest Service General Technical Report RM-229, Fort Collins, CO.
- Sauer, J. R., Hines, J. E., & Fallon, J. (2004). *The North American Breeding Bird Survey, Results and Analysis 1966-2003*. Version 2004.1. USGS Patuxent Wildlife Research Center, Laurel, MD. [Online.] <http://www.mbr-pwrc.usgs.gov/bbs/>
- Sauer, J. R., Hines, J.E., & Fallon, J. (2005). The North American breeding bird survey, results and analysis 1966 - 2005. Version 6.2.2006. *USGS Patuxent Wildlife Research Center, Laurel, MD.*
- Science Advisory Board (SAB). (2010). *Review of a field-based aquatic life benchmark for conductivity in central Appalachian streams*. SAB Draft Report dated December 28, 2010 for Quality Review. 47 pp.
- Science Applications International Corporation. (1998). Analysis of valley fill impacts using benthic macroinvertebrates. *Draft final report to U.S. EPA, 30 Sept. 1998.*
- Seby, F., Gautier, M.P., Lespks, G., Astruc, M. 1997. Selenium speciation in soils after alkaline extraction. *The Science of the Total Environment* 207 (1997) 81-90
- Seby, F., Potin-Gautier, M., Giffaut, E., & Donard, O. F. X., 1998. Assessing the speciation and the biogeochemical processes affecting the mobility of selenium from a geological repository of radioactive wastes to the biosphere. *Analisis* 26, 193-198.
- Serne, R.J. (2007). Kd Values for Agricultural and Surface Soils for Use in Hanford Site Farm, Residential, and River Shoreline Scenarios, *Technical Report for Groundwater Protection Project --Characterization of Systems Task, Prepared for Fluor Hanford, Inc. and the U.S. Department of Energy under Contract DE-AC05-76RL01830, PNNL-16531*
- Silva, P. S., & Giles, C. (2010). Memorandum: Detailed Guidance: Improving EPA Review of Appalachian Surface Coal Mining Operations under the Clean Water Act, National Environmental Policy Act, and the Environmental Justice Executive Order. USEPA, Washington D.C.
- Simmons, J. A., Currie, W.S., Eshleman, K.N., Kuers, K., Monteleone S., Negley, T.L., Pohlad, B.R. (2008.) Forest to reclaimed mine land use change leads to altered ecosystem structure and function. *Ecological Applications*, 18, 104-118.
- Simon, A., Doyle, M., Kondolf, M., F.D. Shields Jr., F.D. , Rhoads, B., and McPhillips, M. (2007). Critical evaluation of how the rosgen classification and associated “natural channel design” methods fail to integrate and quantify fluvial processes and channel response. *Journal of the American Water Resources Association*, 43, 15–25.
- Skorupa, J. P. (1998). Selenium poisoning of fish and wildlife in nature: Lessons from twelve real-world examples. In W. T. Frankenberger, & R. A. Engberg (Eds), *Environmental chemistry of selenium* (pp. 315–354). New York: Marcel Dekker.
- Skousen, J., Ziemkiewicz, P., & Venable, C. (2006). Tree recruitment and growth on 20-year-old, unreclaimed surface mined lands in West Virginia. *International Journal of Mining, Reclamation and Environment*, 20, 142–154.
- Slate, L.O., Shields, F.D., Schwartz, J.S., Carpenter, D.D., & Freeman, G.E. (2007) Engineering Design Standards and Liability for Stream Channel Restoration. *Journal of Hydraulic Engineering*, 133(10): 1099-1102.
- Smith, W.P., Hamel, P.B., & Ford, R.P. (1996). Mississippi Alluvial Valley forest conversion: implications for eastern North American avifauna. *Proceedings of the Annual Conference Southeastern Association Fish and Wildlife Agencies*, 47, 460-469.

- Snyder, C. D., Young, J. A., Lemarie, D. P., & Smith, D. R. (2002.) Influence of eastern hemlock (*Tsuga canadensis*) forests on aquatic invertebrate assemblages in headwater streams. *Canadian Journal of Fisheries and Aquatic Sciences*, 59(2), 262-275.
- Soucek, D. J., & Kennedy, A. J. (2005). Effects of hardness, chloride, and acclimation on the acute toxicity of sulfate to freshwater invertebrates. *Environmental Toxicology and Chemistry*, 24, 1204-1210.
- Soucek, D.J. (2007a). Bioenergetic effects of sodium sulfate on the freshwater crustacean, *Ceriodaphnia dubia*. *Ecotoxicology*, 16(3), 317-325.
- Soucek, D. J. (2007b). Comparison of hardness- and chloride-regulated acute effects of sodium sulfate on two freshwater crustaceans. *Environmental Toxicology and Chemistry*, 26,773-779.
- Soule, M. E. (1991). Conservation: tactics for a constant crisis. *Science*, 253, 744-750.
- Strager, M. P., Petty, J. T., Strager, J. M., & Fulton, J. B. (2009). A spatially explicit framework for quantifying downstream hydrologic conditions. *Journal of Environmental Management*, 90, 1854-1861.
- Stanley, T.R., Jr., Spann, J.W., Smith, G. J., & Rosscoe, R. (1994). Main and interactive effects of arsenic and selenium on mallard reproduction and duckling growth and survival. *Archives of Environmental Contamination and Toxicology*, 26, 444-451.
- Stanley, T.R., Jr., Smith, G. J., Hoffman, D. J., Heinz, G. H., and R. Rosscoe, R. (1996). Effects of boron and selenium on mallard reproduction and duckling growth and survival. *Environmental Toxicology and Chemistry*, 15, 1124-1132.
- Stauffer, J., & Ferreri, C. P. (2002). Characterization of stream fish assemblages in selected regions of mountain top removal/valley fill coal mining. *Pennsylvania State University, State College, PA. Appendix in Mountaintop mining/valley fills in Appalachia. Final programmatic environmental impact statement. US Environmental Protection Agency, Philadelphia, Pennsylvania. <http://www.epa.gov/region03/mtntop/eis2003appendices.htm#appd>*
- Stein, B. A., Kutner, L., & Adams, J. (Eds.). (2000). Precious Heritage, The status of biodiversity in the United States. *Oxford: Oxford University Press*.
- Stephan, C.E., Mount, D.I., Hansen, D.J., Gentile, J.R., Chapman, G.A., & William A. Brungs, W.A. 1985. Guidelines for Deriving Numerical National Water Quality Criteria for the Protection of Aquatic Organisms and their Uses. *U.S. EPA, 822-R-85-100. Office of Research and Development, Duluth, MN.*
- Stephenson, S. L., Ash, A. N., & Stauffer, D. F. (1993). Appalachian oak forests. In W. H. Martin, S. G. Boyce and A. C. Echternacht (Eds.), *Biodiversity of the Southeastern United States: Upland terrestrial communities* (pp. 255-304). *New York: John Wiley & Sons Inc.*
- Stoddard, J. L., Larsen, D. P., Hawkins, C. P. Johnson, R. K., & Norris, R. H. (2006). Setting expectations for the ecological condition of streams: the concept of reference condition. *Ecological Applications*, 16(4), 1267–1276.
- Stottlemyer, D., WVDEP (2011). Personal communication
- Stout, B., & Wallace, J. B. (2003). A survey of eight major aquatic insect orders associated with small headwater streams subject to valley fills from mountaintop mining. *Appendix in Mountaintop mining/valley fills in Appalachia. Final programmatic environmental impact statement. US Environmental Protection Agency, Philadelphia, PA. <http://www.epa.gov/region03/mtntop/eis2003appendices.htm#appd>*
- Strausbaugh, P. D., & Core, E. L. (1978). *Flora of West Virginia. 2<sup>nd</sup> Edition.*

- Stein, B. A., Kutner, L. S., & Adams, J. S. (2000). *Precious Heritage: The Status of Biodiversity in the United States*. Oxford University Press.
- Stucker, H. S. (2000). Biodiversity of Southeastern Minnesota forested streams: relationships between trout habitat improvement practices, riparian communities and Louisiana Waterthrushes. *Master's Thesis, University of Minnesota, St. Paul, MN*.
- Sudduth, E.B., Meyer, J. L., & Bernhardt, E. S. (2007). Stream restoration practices in the Southeastern United States. *Restoration Ecology*, 15(3), 572-583.
- Svec, J. R., Kolka, R. K., & Stringer, J. W. (2005). Defining perennial, intermittent, and ephemeral channels in eastern Kentucky: Application to forestry best management practices. *Forest Ecology and Management*, 214, 170-182.
- Swift, M. C. (2002). Stream ecosystem response to and recovery from, experimental exposure to selenium. *Journal of Aquatic Ecosystem Stress and Recovery*, 9, 159-184.
- Timpano, A.J., Schoenholtz, S.H., Zipper, C.E., Soucek, D.J. (2010). Isolating effects of total dissolved solids on aquatic life in central appalachian coalfield streams. *Paper presented at the 2010 National Meeting of the American Society of Mining and Reclamation, Pittsburgh, PA, Bridging Reclamation, Science and the Community June 5 - 11, 2010. R.I. Barnhisel (Ed. Published by ASMR, 3134 Montavesta Rd., Lexington, KY 40502*
- Tullos, D. D., Penrose, D. L., Jennings, G. D., & Cope, W. G. (2009). Analysis of functional traits in reconfigured channels: implications for the bioassessment and disturbance of river restoration *Journal of the North American Benthological Society*, 28(1), 80–92.
- U.S. Army Corps of Engineers Huntington District. (2006a). Spruce No. 1 Draft Environmental Impact Statement, March 2006. Huntington, WV.
- U.S. Army Corps of Engineers Huntington District. (2006b ). Spruce No. 1 Mine Final Environmental Impact Statement, September 2006. Huntington, WV.
- U.S. Census Bureau; Census 2000; generated 22 December 2009 by Matthew T. Lee using American FactFinder; <http://factfinder.census.gov>
- U. S. Department of Agriculture (USDA). (1998). *Stream Visual Assessment Protocol*. National Water and Climate Center Technical Note 99–1. Natural Resources Conservation Service (NRCS) Aquatic Assessment Workgroup.
- U.S. EPA. (1991). Technical Support Document for Water Quality-based Toxics Control. *EPA/505/2-90-001*. Washington, D.C.
- U.S. Environmental Protection Agency (U.S. EPA). (1998). *Guidelines for ecological risk assessment*. *EPA/630/R-95/002F*. Washington, D.C.
- U.S. EPA. (2000). Stressor identification guidance document. *EPA/822B-00/025*. Washington, D.C.
- U.S. EPA. (2003). Draft Programmatic Environmental Impact Statement on Mountaintop Mining/Valley Fills in Appalachia. <http://www.epa.gov/region3/mtntop/eis2003.htm>
- U.S. EPA. (2004). Draft aquatic life water quality criteria for selenium 2004. *EPA-822-D-04-001*. Washington, D.C.
- U.S. EPA. (2005a). Interagency mountaintop mining/valley fills in Appalachia Programmatic Environmental Impact Statement (PEIS) finalized in October 2005. [www.epa.gov/Region3/mtntop/eis2005.htm](http://www.epa.gov/Region3/mtntop/eis2005.htm).
- U.S. EPA. (2005b). Partition coefficients for metals in surface water, soil, and waste. *EPA/600/R-05/074*. Washington, D.C.

- USEPA. (2006). Wadeable streams assessment: a collaborative survey of the nation's streams. *EPA 841-B-06-002*. U.S. Environmental Protection Agency, Office of Research and Development, Washington, D.C.
- U.S. EPA. (2009). The Effects of Mountaintop Mines and Valley Fills on Aquatic Ecosystems of the Central Appalachian Coalfields. *Office of Research and Development, National Center for Environmental Assessment, Washington, D.C. EPA/600/R-09/138A*.
- U.S. EPA Region 3 Toxicity Testing Database. Accessed January 2010. *Wheeling, WV*.
- U.S. EPA. (2010a). A Field-based Aquatic Life Benchmark for Conductivity in Central Appalachian Streams. *EPA/600/R-10/023A*. Office of Research and Development, National Center for Environmental Assessment, Washington, D.C.
- U.S. EPA. (2010b). Inferring causes of biological impairment in the Clear Fork Watershed, West Virginia. *EPA/600/R-08/146*. Washington, D.C.
- USEPA (unpublished) Golden algae data, "Bailey\_P parvum summary" compiled by USEPA from 2010 CONSOL data.
- U.S. EPA. Notes taken by Scott Fulton at an April 2010 U.S. EPA meeting with Arch Coal. US Environmental Protection Agency (USEPA). (1992). Behavior of Metals in Soils. *EPA/540/S-92/018*
- U.S. Fish and Wildlife Service (U.S. FWS). (1978). Preliminary assessment of the fish and wildlife resources of the Tug Fork River sub-basin. *Elkins, WV*.
- U.S. FWS. (1998a). Permitted stream losses due to valley filling in Kentucky, Pennsylvania, Virginia, and West Virginia: a partial inventory. *Kentucky/Tennessee, Pennsylvania, Southwestern Virginia, and West Virginia. USFWS Ecological Services Field Offices. Sept. 1998*.
- U.S. FWS. (1998b). A survey of aquatic life and terrestrial wildlife habitats on the proposed Spruce No. 1 Surface Mine in Logan County, West Virginia. *Pennsylvania Field Office, State College, PA. 23 pp*.
- U.S. FWS. (2000). The value of headwater streams: results of a workshop. *State College, PA. April 13, 1999*. Sponsored by the Pennsylvania Field Office. April 2000.
- U.S. FWS. (2002). Louisiana Waterthrush habitat model. [http://www.fws.gov/r5gomp/gom/habitatstudy/metadata2/Louisiana\\_waterthrush\\_model.htm](http://www.fws.gov/r5gomp/gom/habitatstudy/metadata2/Louisiana_waterthrush_model.htm).
- U.S. FWS. (2004). 2003 results of sediment, water, and fish tissue analyses for selenium and mercury. A letter with the data and a discussion of the results was sent to Allyn Turner, Director, Division of Water and Waste Management, West Virginia Department of Environmental Protection from David Densmore, U.S. FWS Supervisor on January 16, 2004.
- U.S. FWS. (2005). Public comment package in response to the U.S. Environmental Protection Agency's (EPA) request for scientific information, data, and views pertaining to the "Draft Aquatic Life Criteria Document for Selenium" (Federal Register 69(242):75541-75546; December 17, 2004).
- U.S. FWS. (2007a). A conservation action plan for the Cerulean Warbler (*Dendroica cerulea*). *U.S. Fish and Wildlife Service, Division of Migratory Bird Management, Focal Species Program. Revised version – 30 June 2007. 14 pp*.
- U.S. FWS. (2007b). Indiana Bat (*Myotis sodalis*) Draft Recovery Plan: First Revision. *Fort Snelling, MN. 258 pp*.
- U.S. FWS. (2008). Birds of Conservation Concern 2008. *United States Department of Interior, Fish and Wildlife Service, Division of Migratory Bird Management, Arlington, VA. 85 pp. http://www.fws.gov/migratorybirds/*

- Van Kirk, R.W., & Hill, S. L. (2006). Modeling predicts trout population response to selenium based on individual-level toxicity. *Idaho State University, Final Project Report to Greater Yellowstone Coalition. Idaho Falls, ID.*
- Vannote, R. L., Minshall, G. W., Cummins, K.W., Sedell, J. R., & Cushing, C. E. (1980). The river continuum concept. *Canadian Journal of Fisheries and Aquatic Sciences*, 37, 130-137.
- Van Sickle, J., Hawkins, C. P., Larsen, D. P., & Herlihy, A. T. (2005). A null model for the expected macroinvertebrate assemblage in streams. *Journal of the North American Benthological Society*, 24(1), 178-191.
- Van Sickle, J., Larsen, D. P., & Hawkins, C. P. (2007). Exclusion of rare taxa affects performance of the O/E index in bioassessments. *Journal of the North American Benthological Society*, 26(2), 319-331.
- Vesper, D., Bryant, G., & Ziemkiewicz, P. F. (2004). A preliminary study on the speciation of selenium in a West Virginia watershed. *Paper presented at the 2004 National Meeting of the American Society of Mining and Reclamation and the 25th West Virginia Surface Mine Drainage Task Force, April 18-24, 2004, Morgantown, WV. Published by ASMR, p.1950-1959.*
- Vesper, D.J., Roy, M., & Rhoads, C. L. (2008). Selenium distribution and mode of occurrence in the Kanawha Formation, southern West Virginia, U.S.A. *International Journal of Coal Geology*, 73, 237-249.
- Vesper, D.J., & Roy, M. (2006). Selenium Location and Mode of Occurrence in the Kanawha Formation Rocks in West Virginia. *West Virginia Mine Drainage Task Force Symposium, April 18-19, 2006*
- Villard, M. A., & Maurer, B. A. (1996). Geostatistics as a tool for examining hypothesized declines in migratory songbirds. *Ecology*, 77, 59-68.
- Von Schiller, D., Martí, E., Riera, J. L., Ribot, M., Argerich, A., Fonollà P., & Sabater, F. (2008). Inter-annual, annual, and seasonal variation of P and N retention in a perennial and an intermittent stream. *Ecosystems*, 11, 670–687.
- Wallace, J. B., Webster, J. R., & Lowe, R. L. (1992). High-gradient streams of the Appalachians. In Hackney, C.T., S. M. Adams & W. H. Martin (Eds.), *Biodiversity of the Southeastern United States: Aquatic Communities*. New York: John Wiley & Sons, Inc.
- Wallace, J. B., & Gurtz, M. E. (1986). Response of *Baetis* mayflies (Ephemeroptera) to catchment logging. *American Midland Naturalist*, 115, 25-41.
- Wallace, J. B., & Merritt, R. W. (1980). Filter-feeding ecology of aquatic insects. *Annual Review of Entomology*, 25, 103-132.
- Walters, D. M. (2006). Development of a Fish Index of Biotic Integrity to assess the condition of West Virginia streams: Technical Support Document. *EPA/600/R-06/010. U.S. Environmental Protection Agency, Cincinnati, OH.*
- Warner, R. E. (1979). Fish and wildlife resource needs in riparian ecosystems. *Proceedings of a Workshop. May 30-31, 1979: 53. USFWS, EELUT, Kearneysville, WV.*
- Wayland, M., & Crosley, R. (2006). Selenium and other trace elements in aquatic insects in coal mine-affected streams in the Rocky Mountains of Alberta, Canada. *Archives of Environmental Contamination and Toxicology*, 50, 511-522.
- Weakland, C. A., & Wood, P. B. (2005). Cerulean warbler (*Dendroica cerulea*) microhabitat and landscape-level habitat characteristics in southern West Virginia. *The Auk* 122, 497-508.
- Webster, J. R. (1983). Stability of stream ecosystems. In J.R. Barnes & G.W. Minshall (Eds.), *Stream Ecology* (pp. 355-395). *New York: Plenum.*

- Webster, J. R. (1990). Effects of forest disturbance on particulate organic matter budgets of small streams. *Journal of the North American Benthological Society*, 9, 120-140.
- Webster, J. R. & Benfield, E. F. (1986). Vascular plant breakdown in freshwater ecosystems. *Annual Review of Ecology, Evolution, and Systematics* 17, 567-594.
- Webster, J. R., Wallace, J. B., & Benfield, E. F. (1995). Organic processes in streams of the Eastern United States. In C.E. Cushing, G.W. Minshall, & K.W. Cummins (Eds.), *River and Stream Ecosystems (Ecosystems of the World)*. Amsterdam: Elsevier Science.
- Wells, J. V. (2007). Birder's Conservation Handbook: 100 North American Birds at Risk. Princeton: Princeton University Press,. 464 pp.
- West Virginia Department Environmental Protection (WVDEP) 303(d) lists.  
[http://www.dep.wv.gov/WWE/watershed/IR/Pages/303d\\_305b.aspx](http://www.dep.wv.gov/WWE/watershed/IR/Pages/303d_305b.aspx)
- Williams, D.D. 1987. A laboratory study of predator-prey interactions of running water stoneflies and mayflies. *Freshwat. Biol.* 17: 471-490.
- Williams, D.D. 1996. Environmental constraints in temporary waters and their consequences for the insect fauna. *J. N. Amer. Benth. Soc.* 15: 634-650.
- Williams, D. D., & Hynes, H.B.N. 1977. The ecology of temporary streams II. General remarks on temporary streams. *Internat. Rev. Hydrobiologie.* 62:53-61.
- WVDEP. (1997a). An ecological assessment of the Coal River Watershed. Report number - 05050009 – 1997. Watershed Assessment Program, Division of Water Resources, West Virginia Department of Environmental Protection, Charleston WV.
- WVDEP. (1997b). An ecological assessment of the Elk River Watershed. Charleston, WV.
- WVDEP. (1997c). An ecological assessment of the Upper Kanawha River Watershed. Charleston, WV.
- WVDEP. (2003). Water quality monitoring data for 2002-2003.
- WVDEP. (2006). Maximum Daily Loads for selected streams in the Coal River Watershed, West Virginia, Final Approved Report, and Appendices. Watershed Assessment Program, Division of Water Resources, West Virginia Department of Environmental Protection, Charleston WV.
- WVDEP. (2007a). West Virginia integrated water quality monitoring and assessment report 2006. Division of Water and Waste Management, Charleston, WV.
- WVDEP. (2007b). Selenium Implementation Guidance. In section 32 of the Permitting Handbook.  
<http://www.dep.wv.gov/dmr/handbooks/Documents/Permitting%20Handbook/sect32.pdf>
- WVDEP. (2007c). Tug Fork Watershed: a summary of the Watershed Assessment Section's 1998 and 2003 Monitoring Efforts. Charleston, WV.
- WVDEP. (2008). West Virginia integrated water quality monitoring and assessment report 2008. Division of Water and Waste Management. Charleston, WV  
[http://www.dep.wv.gov/WWE/watershed/IR/Documents/IR\\_2008\\_Documents/WV\\_IR\\_2008\\_Supplements\\_Complete\\_Version\\_EPA\\_Approved.pdf](http://www.dep.wv.gov/WWE/watershed/IR/Documents/IR_2008_Documents/WV_IR_2008_Supplements_Complete_Version_EPA_Approved.pdf)
- WVDEP. (2009a). Selenium bioaccumulation among select stream and lake fishes in West Virginia. Watershed Assessment Program, Division of Water Resources. Charleston WV.



- WVDEP (2009b). Dunkard Creek 2009 Fish Kill Information. Data and links regarding golden algae and Dunkard Creek. <http://www.dep.wv.gov/WWE/watershed/wqmonitoring/Pages/DunkardCreekFishKillInformation.aspx>
- WVDEP, Division of Mining and Reclamation. (2009). Mining permit boundaries. *Data retrieved December 28, 2009, from <http://gis.wvdep.org/data/omr.html>.*
- WVDEP. (2010). Selenium-induced developmental deformities among fishes in West Virginia. *Watershed Assessment Program, Division of Water Resources, West Virginia Department of Environmental Protection, Charleston WV. <http://www.dep.wv.gov/WWE/watershed/wqmonitoring/Documents/Selenium/Se%20Larvae%202010%20final.pdf>*
- WVDEP. Technical Applications and Geographic Information Systems (TAGIS). *WVDEP Equis Trend Station Explorer. <http://gisonline.dep.wv.gov/equis/equis.html>*
- WVDEP. Division of Water and Wastewater Management. Ambient sampling data. *Accessed December 2009.*
- WVDEP (2010a). *Selenium-induced developmental effects among fishes in select West Virginia waters.* Watershed Assessment Program, Division of Water Resources. Charleston WV. The report and data regarding selenium can be accessed at: <http://www.dep.wv.gov/WWE/watershed/wqmonitoring/Pages/SpecialStudiesonwaterquality.aspx>
- WVDEP (2010b), 2010 West Virginia Integrated Water Quality Monitoring and Assessment Report (*pending approval from EPA*) (“2010 Section 303(d) List”)
- WVDEP . Unpublished data, (2010a) “Valley Fill Study” water quality data collected September and October 2009.
- WVDEP. Unpublished data. (2010b). Preliminary report from Dr. David Hambright to WVDEP.
- WVDEP Basic guidelines for biological thresholds to certain water quality constituents used by WVDEP’s TMDL program in stressor identification. *Charleston, WV.*
- WVDEP. Watershed assessment (WAB) database. *Division of Water and Wastewater Management. Accessed December 2009. Charleston, WV.*
- WVDEP Office of Hydrologic Protection Unit (OMR), Division of Mining and Reclamation, Discharge Monitoring Reports from WV1017021, 9/30/2009 and 08/31/2009. *Charleston, WV.*
- WVDEP (2009). Watershed assessment program database. *Accessed December 2009. Charleston, WV.*
- WVDEP.( 2005). The occurrence of selenium in the Upper Kanawa Formation of the Pennsylvania System in the Southern West Virginia Coal Fields, *George Jenkins and Nick Schaer*
- West Virginia Department of Natural Resources (WVDNR). (1976). Basin water quality management plan for the Big Sandy-Tug Fork River Basin. *Charleston, WV.*
- WVDNR, Division of Wildlife Resources. (1977). *Today’s Plan for Tomorrow’s Wildlife: A Strategic Plan for Fish, Game, and Non-Game Management, 1975-1985. Charleston, WV.*
- WVDNR Fish Database. (2009). *Accessed December 2009. Elkins, WV.*
- WVDNR, Division of Natural Resources, Wildlife Resources Section. (2009). Federal Assistance Performance Report: Endangered Species (Animals), Project E-1, Segment 26 (1 October 2008 – 30 September 2009). 47 pp. + *Appendices.*
- West Virginia Geological and Economic Survey (WVGES). (2002) Trace Elements in West Virginia Coals. (*Last revised: March 2, 2002.*) <http://www.wvgs.wvnet.edu/www/datastat/te/SeHome.htm>;  
<http://www.wvgs.wvnet.edu/www/datastat/te/index.htm>

- West Virginia GIS Technical Center, West Virginia University. Department of Geology & Geography. Morgantown, WV. <http://wvgis.wvu.edu/data/dataset.php?ID=230>
- West Virginia Highlands Conservancy, et al. v. McClung, Appeal Nos. 07-10-EQB & 07-12-EQB, West Virginia Environmental Quality Board Final Order, slip op. at 28 (June 12, 2008)
- West Virginia Natural Heritage Program. (2007). Rare, Threatened and Endangered Animals. 14 pp. <http://www.wvdnr.gov/Wildlife/documents/Animals2007.pdf>
- West Virginia University. (2010). Influence of selenium on bluegill in Mud River Reservoir, West Virginia. (Presenter: Alison Anderson, WVU and Patricia Mazik, Ph.D. USGS). 2010 West Virginia Water Conference. West Virginia's Water Resources: Threats and Opportunities. October 6-7, 2010. Morgantown, WV. s
- Whittaker, R. H. (1956). Vegetation of the Great Smoky Mountains. *Ecological Monograph*, 22, 1-44.
- Whittaker, R. H. (1975). *Communities and Ecosystems 2<sup>nd</sup> Edition*. New York: Macmillan.
- Wickham, J. D., Riitters, K. H., Wade, T. G., Coan, M., & Homer, C. (2007). The effect of Appalachian mountaintop mining on interior forest. *Landscape Ecology*, 22, 179-187.
- Wiley, J. B., & Brogan, F. D. (2003). Comparison of peak discharges among sites with and without valley fills for the July 8–9, 2001, flood in the headwaters of Clear Fork, Coal River basin, mountaintop coal-mining region, southern West Virginia. *Open-File Report 03-133, Department of the Interior, U.S. Geological Survey, Charleston, WV*. <http://pubs.usgs.gov/of/2003/ofr03-133/pdf/ofr03133.pdf>
- Wiley, J. B., Evaldi, R. D., Eychaner, J. H., & Chambers, D. B. (2001). Reconnaissance of stream geomorphology, low streamflow, and stream temperature in the mountaintop coal-mining region, southern West Virginia, 1999-2000. *U.S. Department of the Interior, U.S. Geological Survey, Charleston, WV. Water-Resources Investigations Report 01-4092*. [http://pubs.usgs.gov/wri/wri014092/pdf/wri01-4092.book\\_new.pdf](http://pubs.usgs.gov/wri/wri014092/pdf/wri01-4092.book_new.pdf).
- Williams, J. M. (2003). Impacts on terrestrial and streamside herpetofauna by mountaintop removal mining in southern West Virginia. *Ph. D. Dissertation, West Virginia University, Morgantown, WV*.
- Williams, J. M., & Wood, P. B. (2004). Streamside salamanders in valley fill and reference streams in southern West Virginia. *Paper presented at the 2004 National Meeting of the American Society of Mining and Reclamation and The 25th West Virginia Surface Mine Drainage Task Force, April 18-24, 2004. ASMR: Lexington, KY*.
- Wilson, D. E., & Ruff, S. (1999). *The Smithsonian book of North American mammals*. Washington D.C.: Smithsonian Institution Press.
- Wilson, H. L. (1951). Cover mapping and habitat analysis. *Unpublished final report, Federal Aid in Wildlife Restoration Project 21-R. Conservation Commission of West Virginia: 221 & 116 of supplement*.
- Winter, T. C. (2007). The role of groundwater in generating streamflow in headwater areas and in maintaining base flow. *Journal of the American Water Resources Association*, 43, 15–25.
- Wipfli, M. S., Richardson, J. S., & Naiman, R. J. (2007). Ecological linkages between headwaters and downstream ecosystems: transport of organic matter, invertebrates, and wood down headwater channels. *Journal of the American Water Resources Association*, 43, 72–85.
- Wollheim, W. M., Vorosmarty, C.J., Bouwman, A.F., Green, P., Harrison, J., Linder, E., Peterson, B.J., Seitzinger, S.P., Syvitski, J.P.M. (2008). Global N removal by freshwater aquatic systems: a spatially distributed, within-basin approach. *Global Biogeochemical Cycles*, 22, GB2026.

- Wood, P. B., & Edwards, J. W. (2002). Mountaintop removal mining/valley fill Environmental Impact Statement technical study project report for terrestrial studies: terrestrial vertebrate (breeding songbird, raptor, small mammal, herpetofaunal) populations of forested and reclaimed sites. *Draft Report 2001. Final 2002. West Virginia University, Morgantown WV. Appendix in Mountaintop mining/valley fills in Appalachia. Final programmatic environmental impact statement. US Environmental Protection Agency, Philadelphia, Pennsylvania.*  
<http://www.epa.gov/region03/mtntop/eis2003appendices.htm#appd>
- Wood, P. B., Bosworth, S. B., & Dettmers, R. (2006). Cerulean warbler abundance and occurrence relative to large-scale edge and habitat characteristics. *Condor*, 108, 154-165.
- Woods, A. J., Omernik, J. M., Brown, D. D. , & Kiilsgaard, C. W. (1996). Level III and IV ecoregions of Pennsylvania and the Blue Ridge Mountains, the Ridge and Valley, and Central Appalachians of Virginia, West Virginia, and Maryland. *EPA/600/R-96/077. National Health and Environmental Effects Research Laboratory, US Environmental Protection Agency, Corvallis, OR.*
- Wunderle Jr., J. M., & Waide, R. B. (1994). Future prospects for nearctic migrants wintering in Caribbean forests. *Bird Conservation International*, 4, 191-207.
- Wunsch, D. R., Dinger, J. S., Taylor, P. B., Carey, D. I., & Graham C. D. R. (1996). Hydrogeology, hydrogeochemistry, and spoil settlement at a large mine-spoil area in eastern Kentucky: *Star Fire Tract. Kentucky Geological Survey, University of Kentucky, Lexington, KY.*
- Wunsch, D. R., Dinger, J. S., & Graham, C. D. R. (1999). Predicting ground-water movement in large mine spoil areas in the Appalachian Plateau. *International Journal of Coal Geology*, 41, 73-106.
- Yahner, R. H. (1988). Changes in wildlife communities near edges. *Conservation Biology*, 2, 333-339.
- Yuill, C. (2001). Land use assessment: mountaintop mining and the Mountaintop Mining Region of West Virginia. *Draft Report. West Virginia University.*
- Ziegler, C. G., Suter II, G. W., Kefford, B. J., Schofield, K. A., & Pond, G. J. (2007). Common candidate cause: ionic strength. *In U.S. EPA Causal Analysis and Diagnosis Decision Information System.*  
[http://cfpub.epa.gov/caddis/candidate.cfm?section=138&step=24&parent\\_section=132](http://cfpub.epa.gov/caddis/candidate.cfm?section=138&step=24&parent_section=132)
- Ziemkiewicz, P. (2009). Identifying Sources and Effects of Selenium in the Mud River Basin. *West Virginia Water Research Institute Progress Report to Patriot Coal, November 30, 2009.*