

CRITERIA SUMMARY

Lake Erie Basin			
Tier I HNC ($\mu\text{g/l}$)		Tier I HCC ($\mu\text{g/l}$)	
Drinking	Nondrinking	Drinking	Nondrinking
130	3,100	ID	ID

EXPOSURE AND TOXICITY DATA

Human health trophic level 3 bioaccumulation factor ($\text{BAFH}_{\text{TL}3}$) = 5.4 l/kg (MDEQ)
Human health trophic level 4 bioaccumulation factor ($\text{BAFH}_{\text{TL}4}$) = 5.4 l/kg (MDEQ)
Acceptable daily exposure (ADE) = $5.0\text{E-}3$ mg/kg/day (IRIS RfD, last revised 09/01/91)
Carcinogen assessment: Class D; not classifiable (IRIS, last revised 07/01/93)
Cancer slope factor (q_1^*) = Not available (IRIS, last revised 07/01/93)
Body weight of average human (BW) = 70 kg (OAC 3745-1-38)
Relative source contribution factor (RSC) = 0.8 (OAC 3745-1-38)
Per capita water consumption (WC) = 2.0 l/day for drinking water criteria (OAC 3745-1-38)
= 0.01 l/day for nondrinking water criteria (OAC 3745-1-38)
Mean consumption of trophic level three fish ($\text{FC}_{\text{TL}3}$) = 0.0036 kg/day (OAC 3745-1-38)
Mean consumption of trophic level four fish ($\text{FC}_{\text{TL}4}$) = 0.0114 kg/day (OAC 3745-1-38)

REFERENCES

- Integrated Risk Information System. USEPA Office of Research and Development, National Center for Environmental Assessment.
- Michigan Department of Environmental Quality, Surface Water Quality Division. 1997. Bioaccumulation Factor Worksheet for Selenium. Verification Date: 4/25/97.
- Ohio Administrative Code rule 3745-1-38: Methodologies for Development of Human Health Criteria and Values for the Lake Erie Drainage Basin. Effective 10/31/97.

CALCULATION OF HUMAN NONCARCINOGENIC CRITERION (HNC) ^a

$$\text{HNC} = \frac{\text{ADE} \times \text{BW} \times \text{RSC}}{\text{WC} + [(\text{FC}_{\text{TL3}} \times \text{BAFH}_{\text{TL3}}) + (\text{FC}_{\text{TL4}} \times \text{BAFH}_{\text{TL4}})]}$$

$$\begin{aligned} \text{Drinking Water HNC} &= \frac{5.0\text{E-}3 \text{ mg/kg/day} \times 70 \text{ kg} \times 0.8}{2.0 \text{ l/day} + [(0.0036 \text{ kg/day} \times 5.4 \text{ l/kg}) + (0.0114 \text{ kg/day} \times 5.4 \text{ l/kg})]} \\ &= 0.13 \text{ mg/l} = 130 \text{ }\mu\text{g/l} \end{aligned}$$

$$\begin{aligned} \text{Nondrinking Water HNC} &= \frac{5.0\text{E-}3 \text{ mg/kg/day} \times 70 \text{ kg} \times 0.8}{0.01 \text{ l/day} + [(0.0036 \text{ kg/day} \times 5.4 \text{ l/kg}) + (0.0114 \text{ kg/day} \times 5.4 \text{ l/kg})]} \\ &= 3.1 \text{ mg/l} = 3,100 \text{ }\mu\text{g/l} \end{aligned}$$

CALCULATION OF HUMAN CARCINOGENIC CRITERION (HCC) ^a

$$\text{HCC} = \frac{\text{RAD} \times \text{BW}}{\text{WC} + [(\text{FC}_{\text{TL3}} \times \text{BAFH}_{\text{TL3}}) + (\text{FC}_{\text{TL4}} \times \text{BAFH}_{\text{TL4}})]}$$

Insufficient data (no q_1^*).

^aSee Ohio Administrative Code 3745-1-38 effective October 31, 1997.