

CRITERIA SUMMARY

Lake Erie Basin			
Tier I HNC ($\mu\text{g/l}$)		Tier I HCC ($\mu\text{g/l}$)	
Drinking	Nondrinking	Drinking	Nondrinking
31,000	83,000	ID	ID

EXPOSURE AND TOXICITY DATA

Human health trophic level 3 bioaccumulation factor (BAFH_{TL3}) = 54.77 l/kg (MDEQ)
Human health trophic level 4 bioaccumulation factor (BAFH_{TL4}) = 87.69 l/kg (MDEQ)
Acceptable daily exposure (ADE) = 1.79 mg/kg/day (IRIS RfD, last revised 09/30/87)
Carcinogen assessment: Class D; not classifiable (IRIS, last revised 03/01/91)
Cancer slope factor (q_1^*) = Not available (IRIS, last revised 03/01/91)
Body weight of average human (BW) = 70 kg (OAC 3745-1-38)
Relative source contribution factor (RSC) = 0.8 (OAC 3745-1-38)
Per capita water consumption (WC) = 2.0 l/day for drinking water criteria (OAC 3745-1-38)
= 0.01 l/day for nondrinking water criteria (OAC 3745-1-38)
Mean consumption of trophic level three fish (FC_{TL3}) = 0.0036 kg/day (OAC 3745-1-38)
Mean consumption of trophic level four fish (FC_{TL4}) = 0.0114 kg/day (OAC 3745-1-38)

REFERENCES

- Integrated Risk Information System. USEPA Office of Research and Development, National Center for Environmental Assessment.
- Michigan Department of Environmental Quality, Surface Water Quality Division. 1997. Bioaccumulation Factor Worksheet for Xylene. Verification Date: 5/12/97.
- Ohio Administrative Code rule 3745-1-38: Methodologies for Development of Human Health Criteria and Values for the Lake Erie Drainage Basin. Effective 10/31/97.

CALCULATION OF HUMAN NONCARCINOGENIC CRITERION (HNC) ^a

$$\text{HNC} = \frac{\text{ADE} \times \text{BW} \times \text{RSC}}{\text{WC} + [(\text{FC}_{\text{TL3}} \times \text{BAFH}_{\text{TL3}}) + (\text{FC}_{\text{TL4}} \times \text{BAFH}_{\text{TL4}})]}$$

$$\begin{aligned} \text{Drinking Water HNC} &= \frac{1.79 \text{ mg/kg/day} \times 70 \text{ kg} \times 0.8}{2.0 \text{ l/day} + [(0.0036 \text{ kg/day} \times 54.77 \text{ l/kg}) + (0.0114 \text{ kg/day} \times 87.69 \text{ l/kg})]} \\ &= 31 \text{ mg/l} = 31,000 \text{ } \mu\text{g/l} \end{aligned}$$

$$\begin{aligned} \text{Nondrinking Water HNC} &= \frac{1.79 \text{ mg/kg/day} \times 70 \text{ kg} \times 0.8}{0.01 \text{ l/day} + [(0.0036 \text{ kg/day} \times 54.77 \text{ l/kg}) + (0.0114 \text{ kg/day} \times 87.69 \text{ l/kg})]} \\ &= 83 \text{ mg/l} = 83,000 \text{ } \mu\text{g/l} \end{aligned}$$

CALCULATION OF HUMAN CARCINOGENIC CRITERION (HCC) ^a

$$\text{HCC} = \frac{\text{RAD} \times \text{BW}}{\text{WC} + [(\text{FC}_{\text{TL3}} \times \text{BAFH}_{\text{TL3}}) + (\text{FC}_{\text{TL4}} \times \text{BAFH}_{\text{TL4}})]}$$

Insufficient data (no q₁*).

^aSee Ohio Administrative Code 3745-1-38 effective October 31, 1997.