

AL CRIT-00187
-00188

ALH CRIT-00167
-00168

BAF = CRIT-00794

TIER II ACUTE AND CHRONIC AQUATIC LIFE VALUES FOR NAPHTHALENE

Standard:

The procedures described in the Tier II methodology indicate that, except possibly where a locally important species is very sensitive, aquatic organisms should not be affected unacceptably if the four (4) day average concentration of naphthalene does not exceed 26.3 µg/L more than once every three (3) years on the average and if the one (1) hour average concentration does not exceed 201 µg/L more than once every three (3) years on the average.

Calculations:

Acute Aquatic Life:

$$SAV = \text{lowest GMAV}/SAF$$

$$\text{Lowest GMAV} = 3216 \mu\text{g/L}$$

$$SAF = 8.0$$

$$SAV = 3216/8.0 = 401.9 \mu\text{g/L}$$

$$SMC = SAV/2 = 401.9/2 = \mathbf{201 \mu\text{g/L}}$$

Chronic Aquatic Life:

$$SCV = SAV/SACR$$

$$SACR = 15.27 \text{ (Geometric mean of 18, 18, 11)}$$

$$SCV = 401.9/15.27 = \mathbf{26.3 \mu\text{g/L}}$$

Notes:

NONE

DRAFT

DRAFT

Table 1. GMAVs and SMAVs for naphthalene

<u>Genus Mean Acute Value ($\mu\text{g/L}$)</u>	<u>Species</u>	<u>Species Mean Acute Value ($\mu\text{g/L}$)</u>	<u>Acute- Chronic Ratio</u>	<u>Reference Number</u>
5219	Fathead Minnow <u>Pimephales promelas</u>	5219	11	1,3,5,7,8
3216	Rainbow Trout <u>Salmo gairdneri</u>	3216		1,2
3524	Cladoceran <u>Daphnia magna</u>	3441		6,7,9
	Cladoceran <u>Daphnia pulex</u>	3610		4,10

DRAFT

DRAFT

References:

1. DeGraeve, G.M., R.G. Elder, D.C. Wood 1982. Effects of naphthalene and benzene on fathead minnow and rainbow trout. Arch. Environ. Contam. Toxicol. 11: 487-490.
2. Edsall, C.C. 1991. Acute toxicities to larval rainbow trout of representative compounds detected in Great Lakes Fish. Bull. Environ. Contam. Toxicol. 46: 173-178.
3. Geiger, D.L., C.E. Northcott, D.J. Call 1985. Acute Toxicities of Organic Chemicals to Fathead Minnows (Pimephales promelas). Vol II. Center for Lake Superior Environmental Studies University of Wisconsin-Superior. 326 pp.
4. Geiger, J.G. and M. Buikema, Jr. 1982. Hydrocarbons depress growth and reproduction of D. Rilex. Can. J. Fish. Aquat. Sci. 39: 830-836.
5. Holcombe, G.W., G.L. Phipps, M.L. Knuth 1984. The acute toxicity of selected substituted phenols, benzenes and benzoic acid esters to fathead minnows. Environ. Pollut. (Series A) 35: 367-381.
6. LeBlanc, G.A. 1980. Acute toxicity of priority pollutants to water flea (Daphnia magna). Bull. Environ. Contam. Toxicol. 24: 684-691.
7. Milliman, R.E., W.J. Birge, and J.A. Black 1984. Comparative acute toxicity to aquatic organisms of components of coal-derived synthetic fuels. Trans. Am. Fish. Soc. 113: 74-85.
8. USEPA 1980. Ambient water quality criteria for naphthalene. EPA 440/5-80-059.
9. Munoz, M.J. and J.V. Tarazona 1993. Synergistic effects of 2 and four component combinations of polycyclic aromatic hydrocarbons: Phenanthrene, anthracene, naphthalene, and acenaphthene on Daphnia magna. Bull. Environ. Toxicol. 50: 363-368.
10. Smith, S.B., J.F. Savino, and M.A. Blouin 1988. Acute toxicity to Daphnia pulex of six classes of chemical compounds potentially hazardous to Great Lakes aquatic biota. J. Great Lakes Res. 14(4): 394-404.

Last Modified:
January 9, 1997