

Indiana Department of Environmental Management Office of Water Quality - Water Quality Standards Section

TIER II ACUTE AND CHRONIC AQUATIC LIFE VALUES

VANADIUM

CAS RN: 7440-62-2

Water Solubility:

Log K_{ow}:

Standard

The procedures described in the Tier II methodology indicate that, except possibly where a locally important species is very sensitive, aquatic organisms should not be affected unacceptably if the four (4) day average concentration of vanadium does not exceed 12 μ g/L more than once every three (3) years on the average and if the one (1) hour average concentration does not exceed 110 μ g/L more than once every three (3) years on the average.

Calculations

Acute Aquatic Life:

SAV = lowest GMAV/SAF

Lowest GMAV = 1,519 μ g/L SAF = 7.0

 $SAV = 1,519/7.0 = 217 \mu g/L$

 $SMC = SAV/2 = 217/2 = 110 \mu g/L$

Chronic Aquatic Life:

$$SCV = SAV/SACR$$

$$SACR = 18$$

$$SCV = 217/18 = 12 \mu g/L$$

Notes

Cowgill et al. (1985) was not used since the data obtained in this study varied by greater than 10 from Pillard (1995) and ancillary information from another study.

Data

Table 1. GMAVs and SMAVs for vanadium

| Genus Mean Acute Value (µg/L) | <u>Species</u> | | Species Mean Acute Value (µg/L) | Acute- Chronic Ratio | Refei Nur | |
|-------------------------------------|---------------------------------------|-------|---------------------------------------|-------------------------|--------------|---|
| 1,519 | Cladoceran Daphnia magna | | 1,580 | | | 1 |
| | Cladoceran Daphnia magna | | 1,460 | | | 1 |
| 3,903 | Bonytail Chub Gila elegans | 5,300 | | | 2 | |
| | Bonytail Chub Gila elegans | 2,200 | | | 2 | |
| | Bonytail Chub Gila elegans | 5,100 | | | 2 | |
| 4,727 | Razorback Sucker Xyrauchen texanus | | 8,800 | | | 2 |
| | Razorback Sucker Xyrauchen texanus | | 4,000 | | | 2 |

| | Razorback Sucker Xyrauchen texanus | 3,000 | 2 |
|--------|--|--------|---|
| 5,033 | Colorado Squawfish Pychocheilus lucius | 7,800 | 2 |
| | Colorado Squawfish Pychocheilus lucius | 3,800 | 2 |
| | Colorado Squawfish Pychocheilus lucius | 4,300 | 2 |
| 10,247 | Brook Trout Salvelinus fontinalis | 7,000 | 3 |
| | Brook Trout | 15,000 | 3 |
| 16,500 | Salvelinus fontinalis Chinook Salmon Oncorhynchus tschawytscha | 16,500 | 4 |
| | Rainbow Trout Oncorhynchus mykiss | 7,925 | 5 |

References

- 1. Alexander, H.C., W.M. McCarty and E.A. Bartlett 1978. Toxicity of perchloroethylene, trichloroethylene, 1,1,1-trichloroethane and methylene chloride to fathead minnows. Bull. Environ. Contam. Toxicol. 20: 334-352.
- 2. Buccafusco, R.J., S.J. Ells, and G.A. LeBlanc 1981. Acute toxicity of priority pollutants to bluegill (Lepomis macrochirus). Bull. Environ. Contam. Toxicol. 24(5): 446-452.
- 3. Canton, H., and D.M.M. Adema 1978. Reproducibility of short-term and reproduction toxicity experiments with Daphnia magna and comparison of the sensitivity of Daphnia magna with Daphnia pulex and Daphnia cucullata in short-term experiments. Hydrobiologia 59(2): 135-140.
- 4. LeBlanc, G.A. 1980. Acute toxicity of priority pollutants to water flea (<u>Daphnia magna</u>). Bull Environ. Contam. Toxicol. 24(5): 684-691.
- 5. Sloof, W. 1983. Benthic macroinvertebrates and water quality assessment. Some toxicological considerations. Aquatic Toxicol. 4: 73-82.

- 6. Smith, A.D., A. Bharath, C. Mallard 1991. The acute and chronic toxicity of ten chlorinated organic compounds to the American Flagfish (Jordanella floridae). Arch. Environ. Toxicol. 20: 94-102.
- 7. Veith, G., D.J. Call, L.T. Brodie 1983. Structure-toxicity relationships for the fathead minnow, Pimephales promelas: Narcotic industrial chemicals. Can. J. Fish. Aq. Sci. 40(6): 743-748.

Acronyms/Abbreviations

| CAS RN | Chemical Abstract Service Registry Number | | |
|-----------------|---|--|--|
| K _{ow} | Octanol-Water Partition Coefficient | | |
| P (superscript) | Predicted value | | |
| SAV | Secondary Acute Value | | |
| GMAV | Genus Mean Acute Value | | |
| SAF | Secondary Acute Factor | | |
| SMC | Secondary Maximum Concentration | | |
| SCC | Secondary Continuous Concentration | | |
| SACR | Secondary Acute-Chronic Ratio | | |
| FT | Flow-through | | |
| S | Static | | |
| U | Unmeasured | | |
| M | Measured | | |
| EVISTRA | Evaluation and Interpretation of Suitable Test Results in AQUIRE (EPA quality checking method/database) | | |

Revision History

February 24, 1999 Values first developed September 21, 2001 New search for data. No studies added.

Contact Information

David B. Kallander Water Quality Standards Section Indiana Department of Environmental Management 100 North Senate Ave., P.O. Box 6015 Indianapolis, IN 46206-6015 (317) 233-2472

Email: dkalland@dem.state.in.us