




TIER II ACUTE AND CHRONIC AQUATIC LIFE VALUES

4-METHYLPHENOL

CAS RN: 106-44-5
Water Solubility: <0.1 g/100 mL at 21 C
Log K_{ow}:



Standard

The procedures described in the Tier II methodology indicate that, except possibly where a locally important species is very sensitive, aquatic organisms should not be affected unacceptably if the four (4) day average concentration of 4-methylphenol does not exceed 53 µg/L more than once every three (3) years on the average and if the one (1) hour average concentration does not exceed 480 µg/L more than once every three (3) years on the average.

Calculations

Acute Aquatic Life:

$$\text{SAV} = \text{lowest GMAV/SAF}$$

$$\begin{aligned}\text{Lowest GMAV} &= 7,700 \text{ } \mu\text{g/L} \\ \text{SAF} &= 8.0\end{aligned}$$

$$\text{SAV} = 7,700/8.0 = 962.5 \text{ } \mu\text{g/L}$$

$$\text{SMC} = \text{SAV}/2 = 962.5/2 = \mathbf{480 \text{ } \mu\text{g/L}}$$

Chronic Aquatic Life:

$$SCV = SAV/SACR$$

$$SACR = 18$$

$$SCV = 962.5/18 = \mathbf{53 \mu g/L}$$

Data

Table 1. GMAVs and SMAVs for 4-methylphenol

<u>Genus Mean Acute Value ($\mu\text{g/L}$)</u>	<u>Species</u>	<u>Species Mean Acute Value ($\mu\text{g/L}$)</u>	<u>Acute- Chronic Ratio</u>	<u>Reference Number</u>
11,681	Fathead Minnow <u>Pimephales promelas</u>	28,600		2
	Fathead Minnow <u>Pimephales promelas</u>	16,500		3
2,775	Cladoceran <u>Daphnia magna</u>	7,700		1
1,120	Rainbow Trout <u>Oncorhynchus mykiss</u>	7,900		2

References

1. Kuhn,R., M.Pattard, K.Pernak, and A.Winter 1989. Results of the Harmful Effects of Selected Water Pollutants (Anilines, Phenols, Aliphatic Compounds) to *Daphnia magna* Water Res. 23(4):495-499.
2. Degraeve,G.M., D.L.Geiger, J.S.Meyer, and H.L.Bergman 1980. Acute and Embryo-Larval Toxicity of Phenolic Compounds to Aquatic Biota. Arch. Environ. Contam. Toxicol. 9(5):557-568.
3. Geiger,D.L., S.H.Poirier, L.T.Brooke, and D.J.Call 1986. Acute Toxicities of Organic

Acronyms/Abbreviations

CAS RN	Chemical Abstract Service Registry Number
K _{ow}	Octanol-Water Partition Coefficient
P (superscript)	Predicted value
SAV	Secondary Acute Value
GMAV	Genus Mean Acute Value
SAF	Secondary Acute Factor
SMC	Secondary Maximum Concentration
SCC	Secondary Continuous Concentration
SACR	Secondary Acute-Chronic Ratio
FT	Flow-through
S	Static
U	Unmeasured
M	Measured
EVISTRA	Evaluation and Interpretation of Suitable Test Results in AQUIRE (EPA quality checking method/database)

Revision History

June 24, 1999	Values first developed
May 15, 2001	New search for data. No new studies added.

Contact Information

David B. Kallander
Water Quality Standards Section
Indiana Department of Environmental Management
100 North Senate Ave., P.O. Box 6015
Indianapolis, IN 46206-6015
(317) 233-2472
Email: dkalland@dem.state.in.us