

RECEIVED
UNIT CINC

2006 APR 28 AM 9:52

201-16248B

I U C L I D

Data Set

Existing Chemical	: ID: 68953-70-a
CAS No.	: 68953-70-a
Producer related part	
Company	: ToxWorks
Creation date	: 01.11.2005
Substance related part	
Company	: ToxWorks
Creation date	: 01.11.2005
Status	:
Memo	:
Printing date	: 01.12.2005
Revision date	:
Date of last update	: 01.12.2005
Number of pages	: 22
Chapter (profile)	: Chapter: 1, 2, 3, 4, 5, 6, 7, 8, 10
Reliability (profile)	: Reliability: without reliability, 1 , 2, 3, 4
Flags (profile)	: Flags: without flag, confidential, non confidential, WGK (DE), TA-Luft (DE), Material Safety Dataset , Risk Assessment, Directive 67/548/EEC , SIDS

1. General Information

Id 68953-70-8
Date 01.12.2005

1.0.1 APPLICANT AND COMPANY INFORMATION

Type : manufacturer
Name : Huntsman Corp
Contact person : Mr. Ray Papciak
Date :
Street : 10003 **Woodloch** Forest Drive
Town : 77380 The Woodlands, Texas
Country : United States
Phone : 281-719-6094
Telefax :
Telex :
Cedex :
Email :
Homepage :

01.11.2005

Type : manufacturer
Name : **INEOS** Americas LLC
Contact person : Ms. Dana Morisse-Arnold
Date :
Street : 2305 **Brazosport** Blvd.
Town : 77541 Freeport, TX
Country : United States
Phone : 979-415-8511
Telefax :
Telex :
Cedex :
Email :
Homepage :

01.11.2005

1.0.2 LOCATION OF PRODUCTION SITE, IMPORTER OR FORMULATOR

1.0.3 IDENTITY OF RECIPIENTS

1.0.4 DETAILS ON CATEGORY/TEMPLATE

1.1.0 SUBSTANCE IDENTIFICATION

IUPAC Name : Oxirane, reaction products with ammonia, distn, residue
Smiles Code :
Molecular formula :
Molecular weight :
Petrol class :

01.11.2005

1. General Information

Id 68953-70-8
Date 01.12.2005

1.1.1 GENERAL SUBSTANCE INFORMATION

Purity type :
Substance type : organic
Physical status : liquid
Purity :
Colour : amber to dark brown
Odour : slight ammonia

Remark : Product is a distillation residue which contains at least 80% triethanolamine, less than 1% diethanolamine, and the remainder being higher amines.

01.11.2005

1.1.2 SPECTRA

1.2 SYNONYMS AND TRADENAMES

Alkanolamine 5503

01.12.2005

Amine 1-N

01.11.2005

1.3 IMPURITIES

Purity : typical for marketed substance
CAS-No :
EC-No :
EINECS-Name :
Molecular formula :
Value :

01.11.2005

1.4 ADDITIVES

1.5 TOTAL QUANTITY

1.6.1 LABELLING

1.6.2 CLASSIFICATION

1.6.3 PACKAGING

1.7 USE PATTERN

1.7.1 DETAILED USE PATTERN

1.7.2 METHODS OF MANUFACTURE

1.8 REGULATORY MEASURES

1.8.1 OCCUPATIONAL EXPOSURE LIMIT VALUES

1.8.2 ACCEPTABLE RESIDUES LEVELS

1.8.3 WATER POLLUTION

1.8.4 MAJOR ACCIDENT HAZARDS

1.8.5 AIR POLLUTION

1.8.8 LISTINGS E.G. CHEMICAL INVENTORIES

1.9.1 DEGRADATION/TRANSFORMATION PRODUCTS

1.9.2 COMPONENTS

08. 11. 2005

1.10 SOURCE OF EXPOSURE

1.11 ADDITIONAL REMARKS

1.12 LAST LITERATURE SEARCH

1.13 REVIEWS

2. Physico-Chemical Data

Id 68953-70-8
Date 01.12.2005

2.1 MELTING POINT

2.2 BOILING POINT

Value : ca. 372 °C at
Decomposition :
Method : other
Year :
GLP : no data
Test substance :

Remark : from MSDS
01.11.2005

2.3 DENSITY

Type : density
Value : ca. 1.12 g/cm³ at 20°C
Method : other
Year :
GLP : no data
Test substance :

Remark : from MSDS
01.11.2005

2.3.1 GRANULOMETRY

2.4 VAPOUR PRESSURE

Value : < .1 hPa at 20 °C
Decomposition :
Method : other (measured)
Year :
GLP : no data
Test substance :

Remark : from MSDS
23.11.2005

2.5 PARTITION COEFFICIENT

Partition coefficient :
Log pow : < at °C
pH value

23.11.2005

2.6.1 SOLUBILITY IN DIFFERENT MEDIA

2. Physico-Chemical Data

Id 68953-70-8

Date 01.12.2005

Solubility in : Water
Value : at °C
pH value :
concentration : at °C
Temperature effects :
Examine different pol. :
pKa : at 25 °C
Description :
Stable :

Remark : completely soluble in water; from MSDS
01.11.2005

2.6.2 SURFACE TENSION

2.7 FLASH POINT

2.6 AUTO FLAMMABILITY

2.9 FLAMMABILITY

2.10 EXPLOSIVE PROPERTIES

2.11 OXIDIZING PROPERTIES

2.12 DISSOCIATION CONSTANT

2.13 VISCOSITY

2.14 ADDITIONAL REMARKS

3.1.1 PHOTODEGRADATION**3.1.2 STABILITY IN WATER**

Type	:	abiotic
	:	at °C
	:	> 7 day(s) at 25 °C
	:	at °C
Deg. product	:	
Method	:	
Year	:	1992
GLP	:	yes
Test substance	:	other TS: TEA
Method	▪	As part of biodegradation study of TEA in two river water samples, control samples containing TEA and formaldehyde to deactivate microorganisms. Samples were analyzed for TEA over 7 or 10 days.
Result	:	Recovery of added radioactive TEA was greater than 90% up to 7 days in a water sample from one river and greater than 95% in the sample from another river after 10 days. pH in both river samples was 7.7
Reliability	:	(2) valid with restrictions Control samples from biodegradation study; conducted under GLP, but not according to standard water stability study.

01.12.2005

(14)

3.1.3 STABILITY IN SOIL**3.2.1 MONITORING DATA****3.2.2 FIELD STUDIES****3.3.1 TRANSPORT BETWEEN ENVIRONMENTAL COMPARTMENTS**

Type	:	fugacity model level III
Media	:	
Air	:	.0266 % (Fugacity Model Level I)
Water	:	45.9 % (Fugacity Model Level I)
Soil	:	54 % (Fugacity Model Level I)
Biota	:	% (Fugacity Model Level II/III)
Soil	:	.0766 % (Fugacity Model Level II/III)
Method	:	
Year	:	2005
Remark	:	Model based on TEA.
Reliability	:	(2) valid with restrictions EPA model calculations

01.12.2005

(2)

3.3.2 DISTRIBUTION

3. Environmental Fate and Pathways

Id 68953-70-8

Date 01.12.2005

3.4 MODE OF **DEGRADATION** IN ACTUAL USE

3.5 **BIODEGRADATION**

Type : aerobic
Inoculum : industrial sewage
Concentration : .6 **mg/l** related to Test substance
5.4 **mg/l** related to Test substance
Contact time :
Degradation : (±) % after
Result : inherently biodegradable
Deg. product : yes
Method :
Year : 1992
GLP : yes
Test substance : other TS: TEA

Method : The biodegradation of TEA was tested in the presence of activated sewage sludge from a wastewater treatment plant. The sludge was aerated for 48 hours before use. After the addition of the activated sludge mixed liquor and TEA, the sample tubes were sealed with rubber stoppers with carbon dioxide traps. Cultures were analyzed periodically for the disappearance of TEA and formation of degradation products, including carbon dioxide. Recovery of applied radioactive TEA was between 74 and 92%. Cultures were sampled until all TEA had disappeared.

Result : Half-life = 0.02 to 0.1 day. The rate of degradation was 1500 mg **TEA/mg** solids/ hr for the low TEA concentration and 4800 mg **TEA/mg** solids/ hr for the high TEA concentration.

Also tested biodegradation in sandy loam surface soil • half life = 0.5 to 1.8 days for concentrations of 1.4 to 2000 mg/kg.

Biodegradation in two river waters: average half-life = 1.2 days for TEA concentrations of 0.099 and 0.489 **mg/l**.

Reliability : (1) valid without restriction
Standard methodology; well reported.

01.12.2005

(14)

3.6 **BOD5, COD OR BOD5/COD RATIO**

3.7 **BIOACCUMULATION**

3.8 **ADDITIONAL REMARKS**

4.1 ACUTE/PROLONGED TOXICITY TO FISH

4.2 ACUTE TOXICITY TO AQUATIC INVERTEBRATES

Type : static
Species : Ceriodaphnia sp. (Crustacea)
Exposure period : 48 hour(s)
Unit : mg/l
EC50 : = 610 calculated
Analytical monitoring : no
Method : other: NSW EPA procedure
Year : 1999
GLP : no data
Test substance : other TS: TEA

Reliability : (1) valid without restriction
Well-reported and uses standard procedure

01.12.2005

(13)

4.3 TOXICITY TO AQUATIC PLANTS E.G. ALGAE

Species : Scenedesmus subspicatus (Algae)
Endpoint : growth rate
Exposure period : 48 hour(s)
Unit : mg/l
EC10 : = 110 calculated
EC50 : = 750 calculated
Limit test :
Analytical monitoring : no
Method : other: Din 38 412, Part 9, 1988
Year : 1988
GLP : no data
Test substance : other TS: TEA

Method : Culture medium was seeded with 1000 algae cells/ml and incubated at 25 C for 48 hours under controlled lighting. TEA was added at 16-2000 mg/l.

Reliability : (2) valid with restrictions
No data on GLP; method and results described briefly.

01.12.2005

(5)

4.4 TOXICITY TO MICROORGANISMS E.G. BACTERIA

4.5.1 CHRONIC TOXICITY TO FISH

4.5.2 CHRONIC TOXICITY TO AQUATIC INVERTEBRATES

4.6.1 TOXICITY TO SEDIMENT DWELLING ORGANISMS

4.6.2 TOXICITY TO TERRESTRIAL PLANTS

4.6.3 TOXICITY TO SOIL **DWELLING** ORGANISMS

4.6.4 TOX. TO OTHER NON **MAMM.** TERR. SPECIES

4.7 BIOLOGICAL EFFECTS MONITORING

4.6 BIOTRANSFORMATION AND **KINETICS**

4.9 **ADDITIONAL** REMARKS

5.0 TOXICOKINETICS, METABOLISM AND DISTRIBUTION

5.1.1 ACUTE ORAL TOXICITY

Type	• LD50
Value	• > 5000 mg/kg bw
Species	• rat
Strain	• Sprague-Dawley
Sex	• male/female
Number of animals	10
Vehicle	
Doses	• 5000 mg/kg
Method	• OECD Guide-line 401 "Acute Oral Toxicity"
Year	• 1990
GLP	• yes
Test substance	• as prescribed by 1.1 - 1.4
Method	: Five male (209-229 g) and 5 female (145-177 g) Sprague-Dawley rats were fasted overnight and dosed by gavage with 1000 mg/kg of the undiluted test material. They were observed for signs of toxicity at 1 and 4 hours after dosing and once daily for the next 14 days. Body weights were recorded on days 0, 7, and 14. After 14 days all rats were euthanized by carbon dioxide inhalation and necropsied.
Result	: Signs of toxicity included piloerection, salivation and diarrhea in 1 to 3 rats on the day of dosing. Poor grooming was seen in up to 5 rats through day 3. No rats died during the 14 days of observation. Body weights on days 7 and 14 post-dosing appeared normal. Mottled kidneys were observed in males at necropsy; no visible lesions were seen in females.
Reliability	• (1) valid without restriction Meets guidelines and GLP requirements.

23.11.2005

(7)

Type	• LD50
Value	• > 2000 mg/kg bw
Species	: rat
Strain	: Fischer 344
Sex	female
Number of animals	• 3
Vehicle	
Doses	• 2000 mg/kg
Method	•
Year	: 1995
GLP	: no data
Test substance	• as prescribed by 1.1 - 1.4

Method : The study used common methodology, but did not describe in sufficient detail to ascribe to a published standard. Animals were administered the test material by gavage and observed for 14 days.

Result : There was no effect on body weight and no clinical signs of toxicity were seen. There were no deaths.

Reliability (2) valid with restrictions
The report does not identify a Guideline or GLP status, but corresponds with another GLP study.

01.12.2005

(4)

5.1.2 ACUTE INHALATION TOXICITY

5.1.3 ACUTE DERMAL TOXICITY

Type	LD50
Value	: > 3000 mg/kg bw
Species	: rabbit
Strain	: New Zealand white
Sex	: male/female
Number of animals	: 10
Vehicle	:
Doses	: 3000 mg/kg
Method	: OECD Guide-line 402 "Acute dermal Toxicity
Year	: 1990
GLP	: yes
Test substance	• as prescribed by 1 .1 • 1.4
Method	: Approximately 24 hours before dosing, the fur was removed from the trunks of 5 male and 5 female rabbits weighing between 2147 and 2586 grams. The test substance was spread evenly (area not specified) on intact skin and covered with gauze. The trunk was wrapped with rubber dam and an elastic bandage for 24 hours.
Result	• Diarrhea was observed in 3 of 10 rabbits on days 1 and 2 after dosing; no other signs of toxicity were seen. Moderate to severe erythema was present at the test site in all animals upon removal of the test site coverings. One female had necrosis of the skin. No erythema was present by day 3. There were no deaths during the 14 days of observation. Body weights on days 7 and 14 post-dosing appear normal. No lesions were seen during necropsy.
Reliability	: (1) valid without restriction Meets guideline and GLP requirements

03.11.2005

(6)

5.1.4 ACUTE TOXICITY, OTHER ROUTES

5.2.1 SKIN IRRITATION

Species	• rabbit
Concentration	• undiluted
Exposure	• Occlusive
Exposure time	: 24 hour(s)
Number of animals	• 6
Vehicle	:
PDII	: 1.62
Result	• slightly irritating
Classification	• not irritating
Method	: EPA OPP 81-5
Year	: 1990
GLP	• yes
Test substance	as prescribed by 1.1 • 1.4
Method	: Trunks of 3 male and 3 female rabbits (2246 • 2645 grams) were clipped free of hair. The test material was applied to 1 abraded and 2 intact sites on the dorsal trunk of each rabbit. Exposure at 1 intact site was for 4 hours; exposure at the other intact site and the abraded site was for 24 hours. Sites were scored for erythema and edema at 24, 48 and 72 hours after the

5. Toxicity

Id 68953-70-8

Date 01.12.2005

Result : end of the exposure periods. The 4 hour exposure was in compliance with OECD method 404 and the 24 hour exposure in compliance with EPA 81-5.
: Following 4 hours of exposure, there **was** very slight erythema on 2 of 6 rabbits, but no edema. After 72 hours, no erythema or edema was seen, The **PII** for 4 hours of exposure to intact sites was 0.25. Following 24 hours of exposure, slight to severe erythema was observed on all rabbits; there was slight edema on 5 of 6. Erythema and edema was gone from the intact sites by 48 hours, but did not completely clear from the abraded sites in all animals until day 9. The **PII** from 24 hours of exposure (intact and abraded combined) was 1.62.

Reliability : (1) valid without restriction
Meets guideline and GLP requirements

23.11.2005

(8)

Species : rabbit
Concentration : undiluted
Exposure : Occlusive
Exposure time : 24 hour(s)
Number of animals : 1
Vehicle :
PDII :
Result : slightly irritating
Classification : not irritating
Method :
Year : 1995
GLP : no data
Test substance : as prescribed by 1.1 - 1.4

Method : Test material applied to intact skin for 5 consecutive days and to abraded skin 3 times. Sites were observed for irritation.

Result : No erythema or edema was seen at the intact site and slight erythema was seen at the abraded site.

Reliability : (2) valid with restrictions
The report does not provide **sufficient** details to determine if a Guideline or GLP were followed

23.11.2005

(3)

5.2.2 EYE IRRITATION

Species : rabbit
Concentration : undiluted
Dose : .1 ml
Exposure time :
Comment : not rinsed
Number of animals : 6
Vehicle : none
Result : moderately irritating
Classification : irritating
Method : EPA OPP 81-4
Year : 1991
GLP : yes
Test substance : as prescribed by 1.1 - 1.4

Method : 0.1 ml test material instilled in lower eyelid; eyelids held closed for approximately 1 second. Eyes were not washed. Scored after **24, 48** and 72 hours according to Draize scale. Number of rabbits scored as positive according to US Federal Hazardous substances Act Regulations at 16 CFR 1500.42 was recorded. Report includes Draize score and number positive.

Result : There was no **corneal** opacity in any rabbit at any time. Iris irritation was

5. Toxicity

Id 68953-70-8

Date 01.12.2005

observed in 4 of 6 rabbits 1 hour after instillation, but was not present in any rabbits after 24 hours. Conjunctival redness, grade 2 or 3, was observed in all 6 rabbits after 1 hour, in 5 rabbits after 24 hours, in 1 rabbit after 48 hours, but not after 72 hours. Conjunctival swelling, grade 2 or higher, was observed in all six rabbits 1 hour after dosing, but was not present after 24 hours. The mean score using the Draize scale was 18.2 of possible 110.

Reliability : (1) valid without restriction
Meets guideline and GLP requirements

01.12.2005

(9)

Species : rabbit
Concentration : undiluted
Dose : .1ml
Exposure time : 1 hour(s)
Comment :
Number of animals : 1
Vehicle : none
Result : slightly irritating
Classification : not irritating
Method :
Year : 1995
GLP : no data
Test substance : as prescribed by 1 .1 - 1.4

Method : Instilled into both eyes, one washed after 30 seconds; the other was washed after 1 hr.

Result : Slight conjunctival redness and swelling for first 24 hours. No irritation after 48 or 72 hours.

Reliability : (2) valid with restrictions
This was a limited study that did not meet guidelines and no data on GLP

01.12.2005

(3)

5.3 SENSITIZATION

Type : Buehler Test
Species : guinea pig
Concentration : 1st: Induction undiluted occlusive epicutaneous
2nd: Challenge undiluted occlusive epicutaneous
3rd:
Number of animals : 35
Vehicle :
Result : not sensitizing
Classification : not sensitizing
Method : EPA OPP 81-6
Year : 1990
GLP : yes
Test substance : as prescribed by 1.1 - 1.4

Method : Male and female Hartley Guinea pigs (300 to 500 grams) were obtained from USDA registered dealer and housed according to NRC Guidelines. The left shoulder of each animal was clipped free of hair 24 hrs before weekly application of 0.3 ml of the undiluted test material or DNCB (1-chloro-2,4-dinitrobenzene) and the animals held in restrainers during exposure. The test material was applied to 10 males and 10 females beneath a 25 mm Hill Top Chamber (Hill Top Research, Inc., Cincinnati, OH) and covered with a dental dam. DNCB, positive control at 0.3% in 80% in ethanol/20% water, was applied similarly to 2 males and 3 females. The patch and dam were held in place with clips attached to the sides of the animal restrainer. After 6 hours, the dams and patches were removed.

5. Toxicity

Id 68953-70-a

Date 01.12.2005

Thirteen days after the third induction exposure, each animal was challenged by 0.3 ml undiluted test material, or 0.3% DNCB in acetone on a naive site on the left flank as described above. 24 hours after the challenge exposure, the test sites were depilated with Neet Cream Hair Remover for 30 minutes. After 2 hours, the sites were graded. The grading was repeated 24 hours later.

Scores greater than slight patchy erythema were regarded as evidence of sensitization.

Result : All five positive control (DNCB) guinea pigs showed evidence of sensitization (severe erythema). No erythema was evident in any guinea pig following 3 induction and 1 challenge exposure to undiluted test material. Thus, the test material is considered not a sensitizer.

Reliability : (1) valid without restriction
meets guideline and GLP requirements

03.11.2005

(1)

5.4 REPEATED DOSE TOXICITY

5.5 GENETIC TOXICITY 'IN VITRO

Type : Ames test
System of testing : TA98, TA100, TA1538
Test concentration : 167, 500, 1670, 5000, 7500, and 10,000 ug/plate
Cycotoxic concentr. : >10,000 ug/plate
Metabolic activation : with and without
Result : negative
Method : EPA OPPTS 870.5265
Year : 1990
GLP : yes
Test substance : as prescribed by 1 .1 - 1.4

Method : The test material or positive control was mixed with top agar and poured over minimal glucose plates in triplicate for each dose tested. Plates were incubated in the dark at 37 C for 48 hr. Plates were scored for revertant colonies. Positive controls were as follows: -S9: TA100 and TA1538 - sodium azide; TA1537 - 9-aminoacridine; TA98 and TA1538 - 2-nitrofluorene;

A positive result was defined as a dose-dependent statistically significant increase in revertants with at least one concentration that is double the value of the negative control.

Result : All positive controls produced more than double the mutant frequencies of the negative controls. Statistically significant increases that were less than double and not dose dependent were seen with the test material in strains TA100 and TA1535 without activation and in TA1538 with activation. In repeated tests, they did not produce increases. Thus, the laboratory concluded that the test material was not mutagenic.

Reliability : (1) valid without restriction
meets guideline and GLP requirements

01.12.2005

(12)

Type : Unscheduled DNA synthesis
System of testing : Rat hepatocyte primary culture
Test concentration : 0.5, 5, 25, 50, and 5000 ug/ml
Cycotoxic concentr. : 750 ug/ml
Metabolic activation : without

5. Toxicity

Id 68953-70-8

Date 01.12.2005

Result : negative
Method : EPA OPPTS 870.5550
Year : 1991
GLP : **yes**
Test substance : as prescribed by 1.1 - 1.4

Method : The liver of an anesthetized male F344 rat was perfused with EGTA, followed by collagenase. The liver was excised, the capsular membrane opened and hepatocytes detached by gently brushing with a camel's hair brush. Cells were centrifuged and suspended in calf serum. Aliquots of 1×10^5 viable hepatocytes were **inoculated** into 12 well cluster dishes and allowed to attach to coverslips for 2 hours. The cultures were incubated for 18-20 hours with the test material and **[3H]-thymidine**. Cells were swollen and washed; then dipped in NTB-2 photographic emulsion in the dark and stored in the dark for 1 week. Autoradiographs were developed in **D19** and stained with Harris Alum hematoxylin. Slides were evaluated for a net increase in black silver grains over the nucleus by counting 150 hepatocytes.

Result : The dose levels of 750 to 5000 **ug/ml** were not scored as there was excess cytotoxicity. At 100 and 500 **ug/ml**, the net nuclear grain (NNG) counts were slightly greater than the negative control values: **0.5+/-6.2** and **2.4+/-6.1**, respectively. NNG between 0-5 are generally considered a marginal response. The test was repeated at concentrations between 50 and 600 **ug/ml**. None had increased NNG counts. The investigators concluded that the test was negative.

Reliability : (1) valid without restriction
meets guideline and GLP requirements

03.11.2005

(11)

5.6 GENETIC TOXICITY 'IN VIVO'

Type : Micronucleus assay
Species : mouse
Sex : male/female
Strain : CD-1
Route of admin. : i.p.
Exposure period : single dose; assayed at 24, 48 and 72 hours post dosing
Doses : 1000 **mg/kg**
Result : negative
Method : EPA OTS 798.5395
Year : 1990
GLP : **yes**
Test substance : as prescribed by 1.1 - 1.4

Method : A preliminary toxicity test was conducted using 2 male and 2 female CD-1 mice at 500, 750, 1000, 1500 and 5000 **mg/kg** ip in distilled water. 1 of 4 dosed at 1500 and 3 of 4 dosed at 5000 **mg/kg** died. Fifteen males and 15 female 8-week old CD-1 mice were dosed once ip at 1000 **mg/kg**; 5 of each sex were sacrificed after 24, 48, and 72 hours. Five males and 5 females were also exposed to 0.5 **mg/kg** triethylenemelamine in saline by ip injection, as positive controls. Following sacrifice, both femurs were removed from each mouse and marrow removed into fetal bovine serum. The suspension was centrifuged and a small drop of the cell pellet was smeared on a glass slide, dried and stained with modified Wrights Stain Pak 4481. Slides were evaluated for micronuclei in 1000 polychromatic erythrocytes.

Result : The positive control resulted in 3.5% of **PCEs** with micronuclei. Treatment of mice with the test material did not result in increased micronuclei, compared to control. Incidences of micronuclei were 0.04% for negative control mice and **0.02%**, **0.03%**, and 0.04% for mice treated with the test

5. Toxicity

Id 68953-70-8

Date 01.12.2005

Reliability

01.12.2005

material at 1000 mg/kg and sacrificed after 24, 48, and 72 hours, respectively. The test material did not change the ratio of PCEs to NCEs.
: (1) valid without restriction
meets guideline and GLP requirements

(10)

5.7 CARCINOGENICITY

5.8.1 TOXICITY TO FERTILITY

5.8.2 DEVELOPMENTAL TOXICITY/TERATOGENICITY

5.8.3 TOXICITY TO REPRODUCTION, OTHER STUDIES

5.9 SPECIFIC INVESTIGATIONS

5.10 EXPOSURE EXPERIENCE

5.11 ADDITIONAL REMARKS

6.1 ANALYTICAL METHODS

6.2 DETECTION AND IDENTIFICATION

7.1 FUNCTION

7.2 EFFECTS ON ORGANISMS TO BE CONTROLLED

7.3 ORGANISMS TO BE PROTECTED

7.4 USER

7.5 RESISTANCE

8.1 METHODS HANDLING AND STORING

8.2 FIRE GUIDANCE

8.3 EMERGENCY MEASURES .

8.4 POSSIB. OF RENDERING SUBST. HARMLESS

8.5 WASTE MANAGEMENT

8.8 SIDE-EFFECTS DETECTION

8.7 SUBSTANCE REGISTERED AS DANGEROUS FOR GROUND WATER

8.8 REACTIVITY TOWARDS **CONTAINER** MATERIAL

- (1) Armondi, S.E., (1990). Delayed Contact **Hypersensitivity** in Guinea Pigs: PH **424-TX-004-90** (Amine N-I). Pharmakon Research International, Inc., Waverly, PA (dated 11/30/1990), pp. 1-24.
- (2) **EPIWIN** version 3.1, 2000. US EPA
- (3) Gilbert, K.S. (1995). Alkanolamine 5503: Acute Toxicological Properties. Dow Chemical Company, Final Report, Midland, MI, pp. 1-4.
- (4) Gilbert, K.S., (1995). Alkanolamine 5503: Acute Toxicological Properties. Dow Chemical Company, Final Report, Midland, MI, pp. 1-4.
- (5) Kuhn, R., Pattard, M. (1990). Results of the harmful effects of water pollutants to green algae (*Scenedesmus subspicatus*) in the cell multiplication inhibition test. Wat. Res. 24: 31-38.
- (6) **Mallory, V.T.**, (1990). Acute Exposure Dermal Toxicity: PH **422-TX-008-90** (Amine N-I). Pharmakon Research International, Inc., Waverly, PA (dated 10/26/90), pp. 1-5.
- (7) Mallory, V.T. (1990). Acute Exposure Oral Toxicity: PH **402-TX-008-90** (Amine N-I). Pharmakon Research International, Inc., Waverly, PA (dated 10/26/90), pp. 1-19.
- (8) Mallory, V.T. (1992). Primary Dermal Irritation Study: PH **420-TX-007-90:Amended** Report (Amine N-I). Pharmakon Research International, Inc., Waverly, Pa (dated 2/19/92), pp. 1-16.
- (9) Mallory, V.T., (1991). Primary Eye Irritation: PH **421-TX-005-91** (Amine N-I). Pharmakon Research International, Inc, Waverly, PA (dated 2/4/91), pp. 1-19.
- (10) SanSebastian, J.R., (1990). Micronucleus Test: PH **309-TX-003-90** (Amine N-I). Pharmakon Research International, Inc., Waverly, PA (dated 08/31/90), pp. 1-13.
- (11) SanSebastian, **J.R.**, (1991). Rat Hepatocyte Primary Culture/DNA Repair Test: PH 31 I-TX-004-90 (Amine N-I). Pharmakon Research International, Inc., Waverly, Pa (dated 01/24/91), pp. 1-13.
- (12) Stankowski, L.F., Jr. (1990). Ames/Salmonella Plate Incorporation Assay: PH **301-TX-005-90** (Amine N-I). Pharmakon Research International, Inc., Waverly, Pa (dated 07/30/90), pp. 1-13.
- (13) Warne, M. St. J., Schiffko, A. D., (1999). Toxicity of laundry detergent components to a freshwater cladoceran and their contribution to detergent toxicity. Ecotoxicol. Environ. Safety 44: 196-206.
- (14) West, R.J., Gonsior, S.J. (1996). Biodegradation of triethanolamine. Environ. Toxicol. Chem. 15: 472-480.

10.1 END POINT SUMMARY

10.2 HAZARD SUMMARY

10.3 RISK ASSESSMENT