

FOREWORD

INTRODUCTION

NEOPENTYL GLYCOL
CAS N°: 126-30-7

2006 DEC 20 AM 7:39

RECEIVED
OPT/ONIC

Substance

<i>End Point</i>	:	IDENTIFIERS, PHYSICAL AND CHEMICAL PROPERTIES
<i>Chemical Name</i>	:	1,3-Propanediol, 2,2-dimethyl
<i>Common Name</i>	:	Neopentyl glycol
<i>CAS Number</i>	:	126-30-7
<i>RTECS Number</i>	:	TY5775000

Synonyms

2,2-Dimethyl-1,3-propanediol
Neopentyl glycol

Dimethyltrimethylene glycol

Properties & Definitions

<i>Molecular Formula</i>	:	C5H12O2
<i>Molecular Weight</i>	:	104.15
<i>Melting Point</i>	:	127C
<i>Boiling Point</i>	:	208C
<i>Vapour Pressure</i>	:	30 mmHg (140C), 760 mmHg (211C)
<i>Octanol/Water Partition Coefficient</i>	:	log Pow = 0.12 at 25C (measured)
<i>Water Solubility</i>	:	190g/100 ml at 20C (65%)
<i>Impurities</i>	:	Neopentyl glycol formic acid ester and neopentyl glycol isolactic acid ester
<i>General Comments</i>	:	Thermal decomposition occurs at higher than 120C in strong base. Thermal decomposition products: methanol, isobutanol, isobutyl aldehyde, formaldehyde etc.

Overall Evaluation

SIDS INITIAL ASSESSMENT

This substance is presently of low priority for further work.

SHORT SUMMARY OF THE REASONS WHICH SUPPORT THE RECOMMENDATION:

2,2-Dimethyl-1,3-propanediol is stable solid, and the production volume is 12,000 tonnes for 1991 in Japan. This chemical is stable in neutral, acidic or alkaline solutions, and is classified as "not readily biodegradable" by the results of the biodegradation test conducted as SIDS testing. The chemical is non-toxic to fish, daphnids and algae. The chemical showed no genotoxic effects, and NOAEL for repeated dose toxicity was 100 mg/kg/day and NOAEL for reproductive toxicity was 1000 mg/kg/day. Estimated dose of low concern (EDCL) was calculated as 0.1 mg/kg/day and 10.0 mg/kg/day for repeated dose toxicity and reproductive toxicity, respectively. Daily intake of the chemical was estimated as 1.11E mg/day from calculation using MNSEM 145J exposure model. In conclusion, although 2,2-dimethyl-1,3-propanediol is persistent and toxicological test showed moderate toxicity, no further testing is needed at present considering its exposure levels.

However, international information on exposure is needed for consideration of more realistic analysis.

Production-Trade

Chemical Name : **1,3-Propanediol, 2,2-dimethyl**
CAS Number : **126-30-7**
Geographic Area : **JPN**

Production

<u>Quantity</u>	<u>Year</u>
14000 T/Y - P	1985
12000 T/Y - P	1991
4000 T/Y - IM	1991

References

!SIDSP*

OECD/SIDS. Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, (1993)

Uses

Chemical Name : **1,3-Propanediol, 2,2-dimethyl-**
CAS Number : **126-30-7**
Geographic Area : **JPN**

Use

<u>Quantity</u>	<u>Year</u>	<u>Comments</u>
7300 T		Raw material for alkyd resins
5900 T		Raw material for unsaturated polyester resins
1800 T		Raw material for powder paint resin
1000 T		Other uses-unspecified

References

Primary References : **#MITIR***
Chemical Report submitted by the Ministry of International Trade and Industry, Japan

Secondary References : **!SIDSP***
OECD/SIDS. Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, (1993)

Study

End Point : **CONCENTRATION**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl**
CAS Number : **126-30-7**
Study type : **MODEL**
Geographic Area : **JPN**

Test Subject

Organism Medium Specification Lifestage Sex

AIR
AQ
SOIL

Test Method and Conditions

Test method description : Multi-phase, non-steady state equilibrium model (MNSEM 145J) for evaluation of fate of chemicals in environment consisting of air, water, soil and sediment phases and food. Version 145J. All values are calculated.

Test Results

Matrix Concentrations Spec. Date

AIR 1.79E-10 ppm
 Steady state (SS) mass = 1.53E+0g

AQ 5.08E-04 mg/l
 In water SS-mass = 1.02E+07g was also given.

SOIL 3.85E-05 mg/l
 In soil; a second value of SS-mass = 6.16E+04g was also given.

SED 1.53E-03 mg/l
 In sediment SS-mass = 1.53E+05g was also given.

FOOD 2.37E-10 mg/l
 In meat.

FOOD 2.24E-10 mg/l
 In milk.

PLANT 2.42E-04 mg/l
 In vegetation.

References

- Primary Reference* : **#EAMIT***
 MITI ENVIRON. Agency. Exposure Estimation conducted by MITI and Environmental Agency (EA), Japan
- Secondary Reference* : **!SIDSP***
 OECD/SIDS. Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, (1993)
-

Study

End Point : **CONCENTRATION**
Chemical Name : **1-3 Propanediol, 2,2-dimethyl**
CAS Number : **126-30-7**
Geographic Area : **JPN**

Test Subject

Organism Medium Specification Lifestage Sex

AQ **SURF**
SOIL
SED

Species/strain/system : Two areas in Japan

Test Results

<u>Matrix</u>	<u>Concentrations</u>	<u>Spec.</u>	<u>Date</u>
AQ	ND		1977-
Not detected in surface water.(Detection limit:0.2-0.4mg/l)			
SOIL	ND		1977-
Not detected in soil or sediment.(Detection limit:0.002mg/l)			

References

- Primary Reference* : **#MOREA***
 E. A. Environmental Monitoring of Chemicals, Environmental Survey Report (Office of Health Studies, Department of Environmental Health), Japan, (1977)
- Secondary Reference* : **!SIDSP***
 OECD/SIDS. Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, 6-7, (1993)
-

Study

End Point : **HUMAN INTAKE AND EXPOSURE**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl**
CAS Number : **126-30-7**
Geographic Area : **JPN**

Test Subject

Organism Medium Specification Route Lifestage Sex

FOOD

Species/strain/system : Fish, meat, milk and vegetables

Test Method and Conditions

Test method description : Multi-phase, non-steady state equilibrium model (MNSEM 145J) for evaluation of fate of chemicals in environment consisting of air, water, soil and sediment phases. Version 145J (presented by Kikuo Yoshuda). All values are calculated

Test Results

Intake Spec. Date

1.11E-3 mg/d

Total exposure dose calculated.

1.45E-08 mg/d

From inhalation of air.

1.02E-03 mg/d

From drinking water.

4.89E-06 mg/d

From ingestion of fish.

1.76E-11 mg/d

From ingestion of meat.

2.74E-11 mg/d

From ingestion of milk.

9.05E-05 mg/d

From ingestion of vegetable.

General Comments : Consumer exposure seems to be low because this chemical is used as raw material, and processed in closed system except packaging.

References

Secondary Reference : **!SIDSP***
 OECD/SIDS. Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, (1993)

Study

End Point : **BIODEGRADATION**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl**
CAS Number : **126-30-7**
Study type : **LAB**
Geographic Area : **JPN**

Test Subject

Organism Medium Specification

AQ SLUDG

Species/strain/system : Activated sludge 30mg/l as suspended solid

Test Substance

Purity Grade : **99.4%**

Test Method and Conditions

Test method description : OECD Guideline 301 C. The sludge samples were mixed by stirring in a single container and then cultured at 25C for 1 month. GLP: yes
Temperature : **25C**

(An)aerobic : **AEROB**

Exposure

Exposure Period : **1 mo**
Dose / Concentration : **100 mg/l**

Test Results

<i>Quantity</i>	<i>Time</i>	<i>Comments on result</i>
0.6 %	LOSS 14 d	Degree of biodegradation from BOD 14
1 %	LOSS	Degree of biodegradation from DOC
0 %		Degree of biodegradation from GC

Total oxygen demand (TOD) = 64.5 mg

General Comments : These results indicate that neopentyl glycol should be classified as "not readily biodegradable".

References

Primary Reference : **#MITIT***
 Test conducted by the Ministry of International Trade and Industry (MITI), Japan

Secondary Reference : **!SIDSP***
 OECD/SIDS. Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, 4, (1993)

Study

End Point : **PHOTODEGRADATION**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl**
CAS Number : **126-30-7**
Study type : **LAB**

Test Results

Quantity Time Comments on result

Photochemical degradation rate reported as 0.00. T/2 = infinitude.

References

Primary Reference : **#MITIT***
Test conducted by the Ministry of International Trade and Industry (MITI), Japan

Secondary Reference : **!SIDSP***
OECD/SIDS. Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, (1993)

Study

End Point : **HYDROLYSIS**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl**
CAS Number : **126-30-7**
Study type : **LAB**

Test Substance

Purity Grade : **99.4%**

Test Method and Conditions

Test method description : OECD Test Guideline 111, GLP: yes.
Temperature : **25 C**
pH : **4-9**

Test Results

<u>Quantity</u>	<u>Time</u>	<u>Comments on result</u>
50 %	1 y	T/2 of test compound in pH 4.0, 7.0 and 9.0 at 25C.

References

Primary Reference : **#MITIT***
Test conducted by the Ministry of International Trade and Industry (MITI), Japan, (1993)

Secondary Reference : **!SIDSP***
Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, (1993)

Study

End Point : **BIOCONCENTRATION**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl-**
CAS Number : **126-30-7**

Test Subject

Organism Medium Specification Route Lifestage Sex Number exposed Number controls

FISH

Species/strain/system : Japanese carp*

Test Substance

Description of the test substance : Neopentyl glycol
Purity Grade : >98%

Test Method and Conditions

Test method description : Exposure period = 8 weeks. OECD Test Guideline 305C. Flow-through test. GLP: yes.

Exposure

Exposure comments : Level 1 exposure means low exposure level. Level 2 exposure means higher exposure level and is 10x higher in concentration than the low one.

Test Results

<i>Organ</i>	<i>Bioconcent. Factor</i>	<i>Calc Basis</i>	<i>Time</i>	<i>State</i>	<i>Comments on result</i>
	0				log BCF for level 1 exposure.
	1				log BCF for level 2 exposure.

General Comments : * Specific details on the lifestage of the test organism and test conditions were not given.

References

Primary Reference : **#MITIT***
Test conducted by the Ministry of International Trade and Industry (MITI), Japan

Secondary Reference : **!SIDSP***
Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, (1993)

Study

End Point : **MAMMALIAN ACUTE TOXICITY**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl**
CAS Number : **126-30-7**

Exposure Type : **ACUTE**
Dose / Concentration : **3.200 mg/kg**

Test Results

<u>Organism</u>	<u>Medium</u>	<u>Spec.</u>	<u>Route</u>	<u>Lifestage</u>	<u>Sex</u>	<u>Effect</u>	<u>Effect Comments</u>
RAT			ORL			LD50	

References

Primary Reference : **#URKOD***
 Eastman Kodak Company Reports, (1993)

Secondary Reference : **!SIDSP***
 Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, (1993)

Study

End Point : **MAMMALIAN ACUTE TOXICITY**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl**
CAS Number : **126-30-7**

Species/strain/system : Strain not specified

Test Results

<u>Organism</u>	<u>Medium</u>	<u>Spec.</u>	<u>Route</u>	<u>Lifestage</u>	<u>Sex</u>	<u>Effect</u>	<u>Effect Comments</u>
RAT			ORL	ADULT		LD50	Oral acute toxicity dose was reported as 3200mg/kg.

References

Primary Reference : **#URKOD***
 Eastman Kodak Company Reports, (1971)

Secondary Reference : **!SIDSP***
 OECD/SIDS. Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, 10, (1993)

Study

End Point : **MAMMALIAN TOXICITY**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl-**
CAS Number : **126-30-7**

Test Subject

<u>Organism</u>	<u>Medium</u>	<u>Specification</u>	<u>Route</u>	<u>Lifestage</u>	<u>Sex</u>	<u>Number exposed</u>	<u>Number controls</u>
RAT			ORL		M F		

Species/strain/system : Slc: SD strain

Test Substance

Description of the test substance : Neopentyl glycol
Purity Grade : **99.1%**
Vehicle - Solvent : Distilled water

Test Method and Conditions

Test method description : OECD Combined Repeated Dose and Reproductive/Developmental Toxicity Screening Test. Killing day: male/day 43; female/day 4 of lactation. GLP: yes.

Exposure

Dose / Concentration : **100-1000 mg/kg /d**
Exposure comments : Per gavage to 0 (vehicle), 100, 300, 1000mg/kg/day. Administration period: male: 42 days; female: from 14 day before mating to day 3 of lactation.

Test Results

<u>Organ</u>	<u>Effect</u>	<u>Rev.</u>	<u>OnSet</u>	<u>Sex</u>	<u>Affected in Exposed - Controls</u>
	NEF				

There were no dead or no abnormal animals with clinical signs suggested to be relating to the treatment. Bodyweight and food consumption did not reveal consistent or apparently treatment-related differences with the control groups. No observed effects on haematology of the treated male rats.

BLOOD BIOCH

Blood chemical examination revealed an elevation in values of total protein, total bilirubin and albumin for male rats receiving 300 and 1,000mg/kg. Moreover, glucose values were depressed for male rats receiving 1,000mg/kg.

**LIVER SIZE
KIDNEY SIZE**

Absolute and relative weights of the liver and kidney of both male and female rats receiving 300 and 1,000mg/kg were elevated.

LIVER SIZE

Necropsy revealed hypertrophy of the liver in 2 males receiving 1,000mg/kg. No definite lesion was found histologically. Histopathological examination revealed high incidence of protein casts, hyaline droplet and basophilic change of the renal tubules in males at 1,000mg/kg.

NOAEL

Dose or concentration at which no toxic effects were observed: NOAEL: 100mg/kg/day.

General Comments : Estimated Dose of Low Concern: EDLC = 0.1mg/kg/day.

References

- Primary Reference* : **#URMHW***
 Unpublished Report on Combined Repeated Dose and Reproductive/
 Developmental Toxicity Screening Test conducted by the Ministry of Health
 and Welfare (MHW), Japan
- Secondary Reference* : **!SIDSP***
 Screening Information Data Set (SIDS) of OECD High Production Volume
 Chemicals Programme, (1993)

Study

- End Point* : **MAMMALIAN TOXICITY**
Chemical Name : **Neopentyl glycol**
CAS Number : **126-30-7**
Study type : **LAB**

Test Subject

<u>Organism</u>	<u>Medium</u>	<u>Specification</u>	<u>Route</u>	<u>Lifestage</u>	<u>Sex</u>	<u>Number exposed</u>	<u>Number controls</u>
RAT			ORL	ADULT	M F		

Species/strain/system : Slc:SD strain

Test Substance

Purity Grade : **99%**

Test Method and Conditions

Test method description : OECD Combined Repeated Dose and Reproductive/Developmental Toxicity Screening Test. GLP: yes.

Exposure

Exposure Period : **100-1000 mg**
Exposure comments : The doses 0, 100, 300, 1000 mg/kg/day were administered in oral gavage for 42 days to the males and for 14 days before mating to the females and continued through 3-rd day of lactation.

Test Results

<i>Organ</i>	<i>Effect</i>	<i>Rev.</i>	<i>OnSet</i>	<i>Sex</i>	<i>Affected in Exposed - Controls</i>
BLOOD	BIOCH				
Chemical examination of blood revealed elevated values of: total protein, total bilirubin and albumin for male rats receiving 300 and 1000 mg/kg of neopentyl glycol. The glucose values were depressed for male rats receiving 100 mg/kg of the test substance.					
LIVER	SIZE				
Absolute and relative weights of liver and kidneys of both males and females receiving 300 and 1000mg/kg were elevated.					
KIDNY	SIZE				
Absolute and relative weights of liver and kidneys of both males and females receiving 300 and 1000mg/kg were elevated.					
LIVER	STRUC				
Necropsy revealed hypertrophy of the liver in 2 rats receiving the dose of 1000mg/kg but there was not definite lesions found on microscopic examination.					
KIDNY	STRUC				
Histopathological examination revealed high incidence of protein casts, hyaline droplet and basophilic change in renal tubules in male rats on 1000mg/kg dose.					
NOAEL					
Dose of 100mg/kg/day was the dose at which no toxic effects were observed.					
EDCL					
Estimated dose of low concern was calculated as 0.1mg/kg/day .					

References

- Primary Reference* : **#URMHW***
 Unpublished Report on Combined Repeated Dose and Reproductive/ Developmental Toxicity Screening Test conducted by the Ministry of Health and Welfare (MHW), Japan
- Secondary Reference* : **!SIDSP***
 OECD/SIDS. Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, 11-12, (1993)

Study

End Point : **MUTAGENICITY**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl-**
CAS Number : **126-30-7**

Test Subject

Organism Medium Specification Route Lifestage Sex Number exposed Number controls

BACT

Species/strain/system : Salmonella typhimurium /TA100, TA1535, TA98,TA1537; Escheri
chia coli WP2 uvrA

Test Substance

Description of the test substance : Neopentyl glycol
Purity Grade : **99.1%**
Vehicle - Solvent : Distilled water

Test Method and Conditions

Test method description : Japanese Guideline for Screening Mutagenicity Testing of Chemicals.
 Procedure: Plate method. Positive control: * without S9: AF-2 (TA100, WP2 uvrA, TA98), sodium azide (TA1525) and 9-aminoacridine (TA1537); * with S9: 2-aminoanthracene (all strains). GLP: yes.

Exposure

0-5000 ug/ plate
Exposure comments : The exposure doses used: 0, 312.5, 625, 1250, 2500, 5000ug/plate.

Test Results

<u>Organ</u>	<u>Effect</u>	<u>Rev.</u>	<u>OnSet</u>	<u>Sex</u>	<u>Affected in Exposed - Controls</u>
NEF					

The test material was classified as "negative" under the experimental condition used.

General Comments : Minimum concentration of test substance at which toxicity to bacteria was observed: with and without metabolic activation: >5000ug/plate.

References

- Primary Reference* : **#URMMT***
 Unpublished Report on Mutagenicity Test conducted by the Ministry of Health and Welfare (MHW), Japan, (1993)
- Secondary Reference* : **!SIDSP***
 Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, (1993)

Study

End Point : **MUTAGENICITY**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl-**
CAS Number : **126-30-7**
Study type : **LAB**

Test Subject

Organism Medium Specification Route Lifestage Sex Number exposed Number controls

HAMST

VTR

Species/strain/system : Chinese hamster CHL cells

Test Substance

Description of the test substance : Neopentyl glycol
Purity Grade : **99.1%**
Vehicle - Solvent : Distilled water

Test Method and Conditions

Test method description : Japanese Guideline for Screening Mutagenicity Testing of Chemicals. Positive control: mitomycin C and cyclophosphamide S-9: induced by phenobarbital and 5,6-benzoflavone. GLP: YES.

Exposure

0-1.00 mg/ml
Exposure comments : The exposure doses are: 0, 0.25, 0.50, 1.00mg/ml.

Test Results

<u>Organ</u>	<u>Effect</u>	<u>Rev.</u>	<u>OnSet</u>	<u>Sex</u>	<u>Affected in Exposed - Controls</u>
-----	-----	-----	-----	-----	-----
	NEF				

The test material was classified as "negative" under the experimental condition used.

References

- Primary Reference* : **#URMMT***
 Unpublished Report on Mutagenicity Test conducted by the Ministry of Health and Welfare (MHW), Japan, (1993)
- Secondary Reference* : **!SIDSP***
 Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, (1993)

Study

End Point : **REPRODUCTION**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl-**
CAS Number : **126-30-7**
Study type : **LAB**

Test Subject

Organism Medium Specification Route Lifestage Sex Number exposed Number controls

RAT **ORL**

Species/strain/system : Slc: SD strain

Test Substance

Description of the test substance : Neopentyl glycol
Purity Grade : **99.1%**
Vehicle - Solvent : Distilled water

Test Method and Conditions

Test method description : OECD Combined Repeated Dose and Reproductive/Developmental Toxicity Screening Test. Killing day: male: day 43; female: day 4 of lactation. GLP: yes.

Exposure

0-1000 mg/kg /day
Exposure comments : The exposure doses are: 0(vehicle), 100, 300, 1000mg/kg/day. Administration period: male: 42 days; female: from 14 day before mating to day 3 of lactation.

Test Results

<u>Organ</u>	<u>Effect</u>	<u>Rev.</u>	<u>OnSet</u>	<u>Sex</u>	<u>Affected in Exposed - Controls</u>
REPRO	NEF				

There were no effect of test substance on copulation, fertility and estrus cycle of rats. Delivery was normal for dams except for one animal of control group. No effects of test substance on dams during the lactation period were observed.

OFSPR **NEF**

No increase in appearance of abnormal pups to be caused by test substance. Body weight gain of pups was normal up to day 4 of lactation. Stillborn, dead pups and pups killed at day 4 of lactation showed no abnormal gross finding suggested to be attributable to the treatment with test substance.

NOAEL

For P generation: 1000mg/kg

NOAEL

For F1 generation: 1000mg/kg

General Comments : Estimated Doses of Low Concern: EDLC = NOAEL/UF = 1000/100 = 10.0mg/kg/day.

References

Primary Reference : **#URMHW***
Unpublished Report on Combined Repeated Dose and Reproductive/
Developmental Toxicity Screening Test conducted by the Ministry of Health
and Welfare (MHW), Japan, (1993)

Secondary Reference : **!SIDSP***
Screening Information Data Set (SIDS) of OECD High Production Volume
Chemicals Programme, (1993)

Study

End Point : **TERATOGENICITY**
Chemical Name : **Neopentyl glycol**
CAS Number : **126-30-7**
Study type : **LAB**

Test Subject

Organism Medium Specification Route Lifestage Sex Number exposed Number controls

RAT **ORL**

Species/strain/system : Slc: SD strain

Test Method and Conditions

Test method description : OECD Reproduction/Developmental Toxicity Screening Test

Exposure

Dose / Concentration : **100-1000 mg/kg BW**
Exposure comments : In utero exposure to the maternal doses of 0, 100, 300, 1000mg/kg body weight/day of neopentyl glycol for assessment of teratogenic potential.

Test Results

<u>Organ</u>	<u>Effect</u>	<u>Rev.</u>	<u>OnSet</u>	<u>Sex</u>	<u>Affected in Exposed - Controls</u>
-----	-----	-----	-----	-----	-----
FETUS	NEF				

Stillborn, dead pups and pups sacrificed at day 4 of lactation showed no abnormal gross findings suggesting any influence on fetal development from the test substance.

General Comments : External examination of pups revealed no increase in appearance of abnormal pups to be caused by the test substance. Body weight gain of pups was normal up to day 4 of lactation. In the final comment the author stated that no effect on developmental toxicity was observed.

References

- Primary Reference* : **#URMHW***
 Unpublished Report on Combined Repeated Dose and Reproductive/ Developmental Toxicity Screening Test conducted by the Ministry of Health and Welfare (MHW), Japan
- Secondary Reference* : **!SIDSP***
 OECD/SIDS. Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, 14-15, (1993)

Study

End Point : **AQUATIC ACUTE TOXICITY**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl**
CAS Number : **126-30-7**

Species/strain/system : Orange-red Killifish (*Oryzias latipes*)
Exposure Period : **24-96 h**
Exposure comments : The same doses were also tested for 48h and 72h.

Test Method and Conditions

Test method description : Semi-static

Test Results

<u>Organism</u>	<u>Medium</u>	<u>Spec.</u>	<u>Route</u>	<u>Lifestage</u>	<u>Sex</u>	<u>Effect</u>	<u>Effect Comments</u>
FISH	AQ	ESTUA				LC0 LC50	LC0 = 555mg/l (reported as 555ppm) for 24, 48, 72 and 96 hours, LC50 = > 1000mg/l (reported as > 1000ppm (w/v)).

References

Primary Reference : **#UREAF***
 Unpublished Report on Toxicity to Fish Test conducted by Environmental Agency, Japan

Secondary Reference : **!SIDSP***
 OECD/SIDS. Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, (1993)

Study

End Point : **AQUATIC ACUTE TOXICITY**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl**
CAS Number : **126-30-7**

Species/strain/system : Orange-red Killifish (*Oryzias latipes*)
Exposure Period : **48 h**

Test Substance

Impurities : **Water 0.03%, neopentyl hydroxy pivalate 0.44%, formic acid 0.002%**

Test Method and Conditions

Test method description : JIS K0102. Static test.

Test Results

<u>Organism</u>	<u>Medium</u>	<u>Spec.</u>	<u>Route</u>	<u>Lifestage</u>	<u>Sex</u>	<u>Effect</u>	<u>Effect Comments</u>
FISH	ESTUA					LC50	> 1000mg/l (reported > 1000 ppm)

References

Primary Reference : **#UREAF***
Unpublished Report on Toxicity to Fish Test conducted by Environmental Agency, Japan

Secondary Reference : **!SIDSP***
OECD/SIDS. Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, (1993)

Study

End Point : **AQUATIC TOXICITY**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl-**
CAS Number : **126-30-7**

Test Subject

Organism Medium Specification Route Lifestage Sex Number exposed Number controls

ALGAE

Species/strain/system : Algae (Selenastrum capricornutum)

Test Substance

Description of the test substance : Neopentyl glycol
Purity Grade : **>99%**

Test Method and Conditions

Test method description : OECD Test Guideline. GLP: no

Exposure

Exposure Period : **72 h**
Dose / Concentration : **>1000 mg/l w/v**

Test Results

<i>Organ</i>	<i>Effect</i>	<i>Rev.</i>	<i>OnSet</i>	<i>Sex</i>	<i>Affected in Exposed - Controls</i>
-----	-----	-----	-----	-----	-----
	EC50				

Effective concentration (reported as EBC50 > 1000ppm (w/v) for 42h)

References

Primary Reference : **#UREAA***
 Unpublished Report on Toxicity to Algae Test conducted by Environmental Agency, Japan, (1993)

Secondary Reference : **!SIDSP***
 Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, (1993)

Study

End Point : **AQUATIC TOXICITY**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl**
CAS Number : **126-30-7**

Test Subject

Organism Medium Specification Route Lifestage Sex Number exposed Number controls

CRUS

AQ

Species/strain/system : Water flea (Daphnia magna)

Test Substance

Purity Grade : **98%**

Test Method and Conditions

Test method description : Static test. Method used to calculate EC values: Probit method.

Exposure

Exposure Period : **21 d**
24-48 h

Test Results

<i>Organ</i>	<i>Effect</i>	<i>Rev.</i>	<i>OnSet</i>	<i>Sex</i>	<i>Affected in Exposed - Controls</i>
-----	-----	-----	-----	-----	-----
	NOEC				

Maximun concentration at which no effect was observed > 1000ppm (w/v)) for 21days.

EC0

For 24h lowest dose without effect: > 1000ppm (w/v))

EC50

For 24h lowest dose without effect: > 1000ppm (w/v))

References

Primary Reference : **#URTEA***
Unpublished Toxicity Test conducted by the Environmental Agency, (EA), Japan

Secondary Reference : **!SIDSP***
OECD/SIDS. Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, (1993)

Study

End Point : **AQUATIC TOXICITY**
Chemical Name : **1,3-Propanediol, 2,2-dimethyl-**
CAS Number : **126-30-7**

Test Subject

Organism Medium Specification Route Lifestage Sex Number exposed Number controls

CRUS

Species/strain/system : Water flea (Daphnia magna)

Test Substance

Description of the test substance : Neopentyl glycol
Purity Grade : **>98%**

Test Method and Conditions

Test method description : GLP: no. Probit method used to calculate these values.

Exposure

Exposure Type : **ACUTE**
Exposure Period : **24-48 h**
Dose / Concentration : **>1000 ppm w/v**

Test Results

<i>Organ</i>	<i>Effect</i>	<i>Rev.</i>	<i>OnSet</i>	<i>Sex</i>	<i>Affected in Exposed - Controls</i>
-----	-----	-----	-----	-----	-----
	EC0				

EC50

The 24h EC0 and EC50 are higher than 1000ppm w/v.

References

Primary Reference : **#URTEA***
 Unpublished Toxicity Test conducted by the Environmental Agency, (EA), Japan, (1993)

Secondary Reference : **!SIDSP***
 Screening Information Data Set (SIDS) of OECD High Production Volume Chemicals Programme, (1993)

SIDS INITIAL ASSESSMENT PROFILE

CAS No.	97-85-8	RECEIVED DPPT/CBIC 2006 DEC 28 AM 7:39
Chemical Name	Isobutyl isobutyrate	
Structural Formula	CH ₃ -CH(CH ₃)-CH ₂ -O-C(=O)-CH(CH ₃)-CH ₃	

SUMMARY CONCLUSIONS OF THE SIAR

Data from isobutanol toxicity studies have been included in the human health section. Data from isobutanol are useful when assessing the hazards associated with the systemic toxicity of isobutyl isobutyrate exposure due to the rapid and complete metabolism of isobutyl isobutyrate to isobutanol and isobutyric acid *in vivo*. Isobutanol is then further metabolized to isobutyric acid. Therefore, exposure to isobutyl isobutyrate via dermal, inhalation, and water or dietary administration results in the rapid appearance of isobutanol and isobutyric acid in the systemic circulation. Since exposure to either isobutyl isobutyrate or isobutanol results in systemic exposure to isobutanol and isobutyric acid, systemic toxicity data from studies that administer isobutanol directly are useful in identifying hazards associated with isobutyl isobutyrate exposure. Data from studies conducted with isobutyric acid were not included, since there were none available. The toxicokinetics of the metabolic reaction is documented and explained below.

The acute aquatic toxicity database of isobutyl isobutyrate was supported using data from a structural analog, compound, 2-ethylhexyl acetate (CAS# 103-09-3), alleviating the need for additional testing on isobutyl isobutyrate. Data from structurally similar compounds may be used to address the aquatic toxicity of isobutyl isobutyrate.

Toxicokinetics

Metabolism/toxicokinetic studies have been conducted with isobutyl isobutyrate using intravenous injections (Deisinger, 2003). Isobutyl isobutyrate levels peaked immediately after injection and rapidly decreased thereafter. The calculated T_{1/2} by one-compartment modeling was 11.1 seconds. Isobutyl isobutyrate is metabolized extremely rapidly *in vivo* to isobutanol and isobutyric acid.

Human Health

The oral LD₅₀ in rats is >6400 mg/kg bw. Dermal LD₅₀ in male rabbits was >10 ml/kg bw. Inhalation LC₆₆ values for vapor exposures were 5423 ppm (31,940 mg/m³) in rats (6 hours of exposure). Exposures to 658 ppm caused no deaths in 6 hours. Isobutyl isobutyrate is a slight skin irritant.

The NOAEL from an 18-week oral gavage study in rats was 1000 mg/kg bw/day for isobutyl isobutyrate. Studies with isobutanol generally corroborate this value although acute signs of toxicity were noted immediately after oral dosing with isobutanol. The use of different vehicles (corn oil with isobutyl isobutyrate and distilled water with isobutanol) affects the rate of absorption of these related materials and explains the presence or absence of clinical signs immediately after oral exposures. An inhalation two-generation reproductive toxicity study conducted with isobutanol (up to 2500 ppm; 7.58 mg/L) did not cause any parental systemic, reproductive, or neonatal toxicity when administered for two generations via whole-body exposure. No adverse developmental effects were noted in rats or rabbits exposed up to 10 mg/L isobutanol during gestation.

An *in vitro* mutagenicity study in bacteria indicates that isobutyl isobutyrate is not a genotoxicant. In addition,

isobutanol was negative in an *in vivo* mouse micronucleus study.

Environment

The available physicochemical data are adequate to describe the properties of isobutyl isobutyrate. Isobutyl isobutyrate has a melting point of -80°C , boiling point of 148.6°C and vapor pressure of 5.8 hPa at 25°C , a water solubility of 520 mg/L at 20°C and a log K_{ow} of 2.68. The photochemical removal of isobutyl isobutyrate as mediated by hydroxyl radicals occurs with a calculated half-life of 1.947 days. Isobutyl isobutyrate is readily biodegradable under aerobic conditions, based on data for isopropyl- and isobutyl-acetate. Isobutyl isobutyrate volatilizes easily from moving rivers, but volatilizes only moderately from quiescent lakes and other surface water bodies (calculated volatilization half-lives of 1.67 hours from a river and 4.955 days from a lake). Isobutyl isobutyrate is not persistent in the environment and is not likely to bioaccumulate in food webs. Using a log K_{ow} of 2.68, the BCF is 23.1. Based on Level III distribution modeling it is estimated that the majority of isobutyl isobutyrate released to the environment will partition into water (34.4%) and soil (52.7%), with a smaller amount in air (12.6%). The stability of isobutyl isobutyrate in water is pH dependent, at neutral pHs (7) the $T_{1/2} = 9.2$ years at 25°C and at higher pHs (8) the $T_{1/2}$ is shortened to 337 days.

Except for a study with the aquatic invertebrate, *Daphnia magna*, aquatic toxicity data are not available for isobutyl isobutyrate. Data for the structurally similar 2-ethylhexyl acetate (CAS# 103-09-3) were used to supplement the data for isobutyl isobutyrate. For fish, two studies with rainbow trout (*Oncorhynchus mykiss*) and 2-ethylhexyl acetate are available. Acute 96-h LC_{50} s of 8.27 and >4.2 mg/L were reported for 2-ethylhexyl acetate, respectively. Data are available with isobutyl isobutyrate and the invertebrate *Daphnia magna* with 48-h EC_{50} values of 55.8 to 59.3 mg/L reported. In addition, a daphnid study with 2-ethylhexyl acetate reported a 48-h EC_{50} of 22.9 mg/L. Data are available with 2-ethylhexyl acetate and the green alga *Selenastrum capricornutum* with a 72-h EC_{50} value of >21.9 mg/L and a 72-h NOEC of 10.3 mg/L reported. ECOSAR values for isobutyl isobutyrate were calculated to be 9.455 mg/L for fish, 27.556 mg/L for daphnids, and 0.771 mg/L for green algae. ECOSAR values for 2-ethylhexyl acetate were calculated to be 3.057 mg/L for fish, 3.571 mg/L for daphnids, and 0.260 mg/L for green algae. Collectively, the measured data with analog compounds and the ECOSAR values for isobutyl isobutyrate and its analog compound, allow the estimation of the acute aquatic toxicity of isobutyl isobutyrate. ECOSAR estimations of algae toxicity for esters are generally not very predictive of measured values.

Exposure

Workplace exposure may occur via inhalation or dermal contact. Exposure during manufacture is limited by the enclosed nature of the process and by bulk handling and good manufacturing practices. Industrial and occasional consumer exposure can occur both dermally and via inhalation during application of lacquer and thinner formulations containing isobutyl isobutyrate. General population exposure can occur through inhalation of ambient air that may contain low concentrations resulting from industrial or commercial releases. General population exposure also occur through ingestion of foods containing isobutyl isobutyrate either naturally or as a synthetic flavorant (21 CFR § 121.1164).

RECOMMENDATIONS AND_RATIONALE FOR THE RECOMMENDATION AND NATURE OF FURTHER WORK RECOMMENDED

The chemical is currently of low priority for further work. The chemical possesses properties indicating a hazard for the environment (toxicity to aquatic species). Based on data presented by the Sponsor country, and its ready biodegradability, under normal manufacturing, formulation, industrial and consumer use, this chemical is a low priority for further work for human health and the environment. Countries may wish to investigate any exposure scenarios that were not presented by the sponsor country.

Note: This document may only be reproduced integrally. The conclusions and recommendations (and their rationale) in this document are intended to be mutually supportive, and should be understood and interpreted together.

**ROBUST SUMMARIES and
SIDS DOSSIER for:
ISOBUTYL ISOBUTYRATE**

.....

CAS No. 97-85-8

2005 DEC 28 AM 7:39

RECEIVED
OPPT/CRIC

Sponsor Country: U.S.A.

DATE: December 2004

CONTENTS

	<u>Page</u>
SIDS PROFILE	1
SIDS SUMMARY	2
1.0 GENERAL INFORMATION	4
1.01 SUBSTANCE INFORMATION	4
A. CAS-Number	4
B. Name (IUPAC name)	4
C. Name (OECD name).....	4
D. CAS Descriptor.....	4
E. EINECS-Number	4
F. Molecular Formula	4
G. Structural Formula.....	4
H. Substance Group.....	4
I. Substance Remark	4
J. Molecular Weight.....	4
1.02 OECD INFORMATION	4
A. Sponsor Country:.....	4
B. Lead Organisation:.....	4
1.1 GENERAL SUBSTANCE INFORMATION	4
1.2 SYNONYMS	5
1.3 IMPURITIES	5
1.4 ADDITIVES	5
1.5 QUANTITY	5
1.6 LABELLING AND CLASSIFICATION	5
1.7 USE PATTERN	5
A. General.....	5
B. Uses in Consumer Products	5
1.8 OCCUPATIONAL EXPOSURE LIMIT VALUE	5
1.9 SOURCES OF EXPOSURE	6
1.10 ADDITIONAL REMARKS	6
2.1 MELTING POINT	6
2.2 BOILING POINT	6
2.3 DENSITY	6
2.4 VAPOUR PRESSURE	7
2.5 PARTITION COEFFICIENT log₁₀Pow	7
2.6 WATER SOLUBILITY	8
2.7 FLASH POINT (liquids)	9
2.8 AUTO FLAMMABILITY (solid/gases)	10
2.9 FLAMMABILITY	10
2.10 EXPLOSIVE PROPERTIES	10
2.11 OXIDIZING PROPERTIES	10
2.12 ADDITIONAL REMARKS	10
2.13 ADDITIONAL DATA	10
3.0 ENVIRONMENTAL FATE AND PATHWAYS	10
3.1 STABILITY	10
3.1.1 PHOTODEGRADATION	10
3.1.2 STABILITY IN WATER	11
3.1.3 STABILITY IN SOIL	11
3.2 MONITORING DATA (ENVIRONMENT)	11

3.3	TRANSPORT AND DISTRIBUTION BETWEEN ENVIRONMENTAL COMPARTMENTS INCLUDING ESTIMATED ENVIRONMENTAL CONCENTRATIONS AND DISTRIBUTION PATHWAYS	11
3.3.1	TRANSPORT BETWEEN ENVIRONMENTAL COMPARTMENTS	11
3.3.2	THEORETICAL DISTRIBUTION (FUGACITY CALCULATION)	12
3.4	IDENTIFICATION OF MAIN MODE OF DEGRADABILITY IN ACTUAL USE	12
3.5	BIODEGRADATION	12
3.6	BOD₅, COD OR RATIO BOD₅/COD	14
3.7	BIOACCUMULATION	14
3.8	ADDITIONAL REMARKS	14
	A. Sewage Treatment	14
	B. Other	14
4.0	ECOTOXICOLOGICAL DATA	15
4.1	ACUTE/PROLONGED TOXICITY TO FISH	15
4.3	TOXICITY TO AQUATIC PLANTS e.g. Algae	25
4.4	TOXICITY TO BACTERIA	28
4.5	CHRONIC TOXICITY TO AQUATIC ORGANISMS	28
4.5.1	CHRONIC TOXICITY TO FISH	28
4.5.2	CHRONIC TOXICITY TO AQUATIC INVERTEBRATES	28
4.6	TOXICITY TO TERRESTRIAL ORGANISMS	28
4.6.1	TOXICITY TO SOIL DWELLING ORGANISMS	28
4.6.2	TOXICITY TO TERRESTRIAL PLANTS	28
4.6.3	TOXICITY TO OTHER NON MAMMALIAN TERRESTRIAL SPECIES (INCLUDING AVIAN)	28
4.7	BIOLOGICAL EFFECTS MONITORING (INCLUDING BIOMAGNIFICATION)	28
4.8	BIOTRANSFORMATION AND KINETICS	29
4.9	ADDITIONAL REMARKS	29
5.0	TOXICITY	29
5.1	ACUTE TOXICITY	29
5.1.1	ACUTE ORAL TOXICITY	29
5.1.2	ACUTE INHALATION TOXICITY	32
5.1.3	ACUTE DERMAL TOXICITY	33
5.2.1	SKIN IRRITATION/CORROSION	33
5.2.2	EYE IRRITATION/CORROSION	33
5.4	REPEATED DOSE TOXICITY	33
5.5	GENETIC TOXICITY IN VITRO	36
A.	BACTERIAL IN VITRO TEST	36
B.	NON-BACTERIAL IN VITRO TEST	37
5.6	GENETIC TOXICITY IN VIVO	37
5.7	CARCINOGENICITY	38
5.8	TOXICITY TO REPRODUCTION	38
5.9	DEVELOPMENTAL TOXICITY/TERATOGENICITY	39
5.10	OTHER RELEVANT INFORMATION	41
A.	Specific toxicities	41
B.	Toxicodynamics, toxicokinetics	41
6.0	REFERENCES	43

SIDS PROFILE

DATE: December 2003

1.01 A.	CAS No.	97-85-8
1.01 C.	CHEMICAL NAME (OECD Name)	Isobutyl Isobutyrate
1.01 D.	CAS DESCRIPTOR	N/A
1.01 G.	STRUCTURAL FORMULA	CH ₃ -CH(CH ₃)-CH ₂ -o-C(=O)-CH(CH ₃)-CH ₃
	OTHER CHEMICAL IDENTITY INFORMATION	
1.5	QUANTITY	10M-50M lbs. 4.5-22.7 thousand metric tons in 2002 in the U.S. only
1.7	USE PATTERN	Solvent, especially for nitrocellulose lacquers and thinners; food flavorant and in perfumes, possible ingredient in insect repellent
1.9	SOURCES AND LEVELS OF EXPOSURE	From industrial waste streams, evaporation during use as a solvent, natural occurrence in fruits and essential oils.
ISSUES FOR DISCUSSION (IDENTIFY, IF ANY)		

SIDS SUMMARY

DATE: November, 2003

CAS No: 97-85-8		Info Available	OECD Study	GLP	Other Study	Estimation Method	Acceptable	Testing Required
Study		Y/N	Y/N	Y/N	Y/N	Y/N	Y/N	Y/N
PHYSICAL CHEMICAL DATA								
2.1	Melting Point	Y			Y		Y	N
2.2	Boiling Point	Y			Y		Y	N
2.3	Density	Y			Y		Y	N
2.4	Vapour Pressure	Y			Y		Y	N
2.5	Partition Coefficient	Y			Y		Y	N
2.6.	Water Solubility	Y			Y		Y	N
	pH and PkA values	N			Y		Y	N
2.12	Oxidation Reduction Potential	N			Y		Y	N
ENVIRONMENTAL FATE and PATHWAYS								
3.1.1	Photodegradation	Y				Y	Y	N
3.1.2	Solubility in water	Y				Y	Y	N
3.2	Monitoring data	Y				Y	Y	N
3.3	Transport and Distribution	Y				Y	Y	N
3.5	Biodegradation	Y			Y		Y	N
OTHER ENVIR FATE STUDIES RECEIVED								
ECOTOXICITY								
4.1	Acute Toxicity to Fish	Y						N
4.2	Acute Toxicity to Daphnia	Y	Y	Y	Y		Y	N
4.3	Toxicity to Algae	Y	Y	Y	Y		Y	N
4.5.2	Chronic Toxicity to Daphnia	N						N
4.6.1	Toxicity to Soil Dwelling Organisms	N						N
4.6.2	Toxicity to Terrestrial Plants	N						N
4.6.3	Toxicity to Birds	N						N
OTHER ECOTOXICITY STUDIES RECEIVED								

SIDS SUMMARY (Continued)

CAS No: 97-85-8		Info Available	OECD Study	GLP	Other Study	Estimation Method	Acceptable	Testing Required
Study		Y/N	Y/N	Y/N	Y/N	Y/N	Y/N	Y/N
TOXICITY								
5.1.1	Acute Oral	Y			Y		Y	N
5.1.2	Acute Inhalation	Y			Y		Y	N
5.1.3	Acute Dermal	Y			Y		Y	N
5.4	Repeated Dose	Y ¹	Y				Y	N
5.5	Genetic Toxicity <i>in vitro</i>	Y			Y		Y	N
	-Gene Mutation						Y	N
5.6	Genetic Toxicity <i>in vivo</i>	Y ¹	Y				Y	N
	-Chromosome Aberration						Y	N
5.8	Reproduction Toxicity	Y ¹					Y	N
5.9	Development/Teratogenicity	Y ¹	Y				Y	N
5.11	Human Experience							
OTHER TOXICITY STUDIES RECEIVED								

Y¹ = Data from isobutanol studies.

1.0 GENERAL INFORMATION

1.01 SUBSTANCE INFORMATION

A. CAS-Number

97-85-8

B. Name (IUPAC name)

Isobutyl Isobutyrate

C. Name (OECD name)

Isobutyl Isobutyrate

D. CAS Descriptor

Not applicable in this case

E. EINECS-Number

F. Molecular Formula

C₈ H₁₆ O₂

G. Structural Formula

CH₃-CH(CH₃)-CH₂-O-C(=O)-CH(CH₃)-CH₃

H. Substance Group

I. Substance Remark

J. Molecular Weight

144.21 g/mol

1.02 OECD INFORMATION

A. Sponsor Country:

United States of America

B. Lead Organisation:

Name of Lead Organisation: American Chemistry Council

Contact person: Barbara Francis

Address: 1300 Wilson Blvd.

Arlington, VA 22209

U.S.A.

Tel: 703-741-5609

Fax: 703-741-6091

1.1 GENERAL SUBSTANCE INFORMATION

A. Type of Substance

Natural organic compound

B. Physical State (at 20°C and 1.013 hPa)
Liquid

C. Purity (indicate the percentage by weight/weight)

1.2 SYNONYMS

Isobutyl 2-methylpropanoate
2-Methylpropyl 2-methyl propanoate
IBIB

1.3 IMPURITIES

None

1.4 ADDITIVES

None

1.5 QUANTITY

3.6 – 5.9 thousand metric tons in 2002 in the U.S. only
8M – 13 Mlbs

1.6 LABELLING AND CLASSIFICATION

No data available

1.7 USE PATTERN

A. General

Type of Use: **Category: Closed process**
Industrial, commercial solvent, automotive coatings, industrial coatings, graphic arts, wood furniture

Type of Use: **Category: Wide dispersive**

B. Uses in Consumer Products

Food flavorant, insect repellent, found in perfumes

1.8 OCCUPATIONAL EXPOSURE LIMIT VALUE

Remark: Thirty-five personal samples were gathered on personnel working in the isobutyl isobutyrate manufacturing area. All samples were determined to be below 1 ppm.
Reference: Eastman Chemical Company, Industrial Hygiene Department. Unpublished communications. February-March, 1984.

1.9 SOURCES OF EXPOSURE

From industrial waste streams, evaporation during use as a solvent, natural occurrence in fruits and essential oils.

1.10 ADDITIONAL REMARKS

No additional remarks

2.0 PHYSICAL-CHEMICAL DATA

2.1 MELTING POINT

- (a) Preferred value
Value: -80.7 °C
Method: other: not reported
GLP: no data
Test substance: Isobutyl Isobutyrate [CAS #97-85-8]; colorless liquid (Hawley's Condensed Chemical Dictionary. 1987.) 25.10.2000
Reliability: Score=2 valid with restrictions, lack of method details
Reference: CRC Handbook of Chemistry and Physics. 1991-1992. D.R. Lide (ed.) 72nd ed. CRC Press, Inc. Boca Raton, FL. P.3-155.

Hawley's Condensed Chemical Dictionary. 1987. N.I. Sax and R.J. Lewis, Sr. (eds.). 11th ed. Van Nostrand Reinhold Co. New York, NY. P.654.

2.2 BOILING POINT

- (a) Preferred value
Value: 148.6 °C
Method: other: not reported
GLP: no data
Test Substance: Isobutyl isobutyrate [CAS#97-85-8]; colorless liquid (Hawley's Condensed Chemical Dictionary. 1987.) 25.10.2000
Reliability: Score=2 valid with restrictions, lack of method details
Reference: CRC Handbook of Chemistry and Physics. 1991-1992. D.R. Lide (ed.) 72nd ed. CRC Press, Inc. Boca Raton, FL. P.3-155.

- (b) Value: 147-148 °C at 1013.25 hPa
Method: other: not reported
GLP: no data
Test Substance: Isobutyl isobutyrate [CAS#97-85-8]; colorless liquid (Hawley's Condensed Chemical Dictionary. 1987.) 25.10.2000
Remark: Reported as 147-148 deg. C @ 760 mm Hg
Reference: Sigma-Aldrich Product Information.
<http://www.sigma-aldrich.com>

2.3 DENSITY

- (a) Preferred result
Type: density

Isobutyl Isobutyrate SIDS Dossier

- Value: 0.855 g/cm³ at 20⁰C
Method: other: not reported
Test Substance: Isobutyl isobutyrate [CAS#97-85-8]; colorless liquid, .97% pure (confirmed by GC)
Reliability: Score=2 valid with restrictions, lack of method details
Reference: Merck-Schuchardt. Hohenbrunn, Germany. Safety Data Sheet. Isobutyl isobutyrate for synthesis. 04.2000. the Merck ChemicalDatabase-online. <http://www.merck.de>
- (b) Type: Relative density
Value: 0.855
Method: other: not reported
GLP: no data
Test substance: Isobutyl isobutyrate [CAS #97-85-8]; liquid, 99% pure
Reference: Sigma-Aldrich Product Information
<http://www.sigma-aldrich.com>
- (c) Type: relative density
Value: 0.875 at 4⁰C
Method: other: not reported
GLP: no data
Test substance: Isobutyl isobutyrate [CAS #97-85-8]; colorless liquid (Hawley's Condensed Chemical Dictionary. 1987.)
Remark: Value also reported for test conducted @ 0 deg. C
Reference: CRC Handbook of Chemistry and Physics. 1991-1992. D.R. Lide (ed.) 72nd ed. CRC Press, Inc. Boca Raton, FL. P.3-155.

2.4 VAPOUR PRESSURE

- (a) Preferred value
Value: 5.76 hPa at 25⁰C
Method: other (measured): not reported
GLP: no data
Remark: Reported as 4.33 mm Hg at 25° C
Test substance: Isobutyl Isobutyrate [CAS# 97-85-8]
Reliability: Score=2 valid with restrictions, lack of method details
Reference: Daubert, T.E. and R.P. Danner. 1989. Physical and Thermodynamic Properties of Pure Chemicals. Data Compilation. Hemisphere Publishing Corp. New York, NY.

2.5 PARTITION COEFFICIENT log₁₀Pow

- (a) Preferred value
Log pow: 2.68 at 25⁰C
Method: other (calculated): Atom/fragment contribution method; KOWWIN
GLP: no
Test substance: Isobutyl isobutyrate [CAS # 97-85-8]
Reliability: Score=2 valid with restrictions, calculated value

Reference: KOWWIN, v1.67. Log Octanol-Water Partition Coefficient. EPISUITE Version 3.10. Syracuse Research Corporation. Syracuse, NY.

2.6 WATER SOLUBILITY

- (a) Preferred value
 Value: 520 mg/L at 27±0.5° C
 Qualitative: Moderately soluble (100-1000 mg/L)
 Method: The slow-stir method described by Ellington (1999) was used. Duplicate glass tanks (20-L) were filled with 20 L deionised water, covered with glass, and sealed with parafilm. Test substance was placed in the tank by slow introduction via pipette. The tank was placed on a stir plate and stirred continuously with a Teflon stir bar at a rate that no vortex was observed. Sample aliquots were collected on days 0, 3, 4, 7, 8, 9, 10, and 14, extracted in methanol and analysed using GC-FID.
- GLP: yes
 Results: pH values of samples were 5.2±0.7. Daily results are shown in the table as follows:

Day	Slow-stirring Tanks	
	(Mean concentration, mg/L)	
Blank tanks	<2.0	<2.0
3	<i>423.4</i>	<i>455.2</i>
4	<i>485.4</i>	<i>518.8</i>
7	<i>546.7</i>	<i>525.8</i>
8	<i>526.9</i>	<i>511.0</i>
9	<i>506.2</i>	<i>533.8</i>
10	<i>495.7</i>	<i>496.6</i>
14	<i>517.5</i>	<i>537.4</i>
Mean	520.0	

Data in italics font not used to calculate the mean water solubility.

Test substance: other TS, Isobutyl isobutyrate [CAS #97-85-8]
 Reliability: (1) valid without restrictions
 Reference: Ellington JJ. (2003) J Chem Eng Data. 44: 1414-1418
 Hoffman CM (2003) Isobutyl Isobutyrate – Water Solubility. Final Report # 1727-WS, Eastman Kodak Company, Rochester, NY.

- (b) Value: 1000 mg/L at 20°C
 Qualitative: moderately soluble (100-1000 mg/L)
 Pka: at 25 °C
 Method: other: stated to be measured, details not given
 GLP: no data
 Test substance: Isobutyl isobutyrate [CAS #97-85-8]
 Reliability: Score=2 valid with restrictions, lack of method details

- Reference: Flick, E.W. 1991. Industrial Solvents Handbook. 4th ed. Noyes Data Corp. Park Ridge, NJ. P.815.
- (c) Value: 5000 mg/L
 Qualitative: soluble (1000-10000 mg/L)
 Pka:
 PH:
 Method: other: not reported
 GLP: no data
 Test substance: Isobutyl isobutyrate
 Remark: Reported as 0.5 g/100 mL
 Reference: Industrial Hygiene and Toxicology. 1989. F. Patty (ed.). 2nd ed. Volume II: Toxicology. Interscience Publishers. New York, NY. P. 1866.
- (d) Value: 670.5 mg/l at 25^oC
 Qualitative: moderately soluble (100-1000 mg/L)
 Pka:
 PH:
 Method: other, calculated, WSKOW
 GLP: no
 Test substance: Isobutyl isobutyrate
 Remark: Log Kow used for calculation = 2.68 (estimated); melting point used for calculation = -80.7 deg. C
 Reliability: Score=2, valid with restrictions, calculation procedure
 Reference: WSKOW, v1.41. Water Solubility Estimate from Log Kow. EPISUITE Version 3.11. Syracuse Research Corporation. Syracuse, NY.
- (e) Value: 1273 mg/L at 25^oC
 Qualitative: moderately soluble (100-1000 mg/L)
 Pka:
 PH:
 Method: Other, calculated, WATERNT
 GLP: No
 Test Substance: other, TS, Isobutyl Isobutyrate
 Remark: Structural fragment method
 Reliability: Score = 2, valid with restrictions, calculation procedure
 Reference: WATERNT, 1.01, EPI Suite v.3.11, United States Environmental Protection Agency, 2003.

2.7 FLASH POINT (*liquids*)

- (a) Preferred value
 Value: 37.2^oC
 Method: other: not reported
 Test substance: Isobutyl isobutyrate; liquid, 99% pure
 Reliability: Score=2, valid with restrictions, lack of method details
 Reference: Sigma-Aldrich Product Information
<http://www.sigma-aldrich.com>

(b) Value: 38°C
Method: other: not reported
GLP: no data
Test substance: other TS
Isobutyl isobutyrate; colorless liquid, .97% pure (confirmed by GC)
Reliability: Score = 2, valid with restrictions, lack of method details
Reference: Merck-Schuchardt. Hohenbrunn, Germany. Safety Data Sheet. Isobutyl isobutyrate for synthesis. 04.2000. The Merck Chemical Database-online. <http://www.merck.de>

2.8 AUTO FLAMMABILITY (solid/gases)

No data available

2.9 FLAMMABILITY

No data available

2.10 EXPLOSIVE PROPERTIES

No data available

2.11 OXIDIZING PROPERTIES

No data available

2.12 ADDITIONAL REMARKS

No additional remarks

2.13 ADDITIONAL DATA

No additional data

3.0 ENVIRONMENTAL FATE AND PATHWAYS

3.1 STABILITY

3.1.1 PHOTODEGRADATION

(a) Type: other: see remarks
Light source:
Light spect.: nm
Rel. intensity: based on Intensity of Sunlight
Deg. Product: N/A
Method: EPIWIN Program
Year: N/A
GLP: No
Test substance: Isobutyl isobutyrate
Remark: If released to the atmosphere, isobutyl isobutyrate is expected to degrade by reaction with photochemically-produced hydroxyl (OH) radicals. The 2nd order rate constant was calculated as 5.49E-12 cm³/(molecules-sec) at 25°C. Based on 1.5E6 OH molecules/cm³ and assuming 12 hours of sunlight per day, the estimated half-life is 1.947 days.
Reference: Atkinson, R. 1987. A Structure-activity relationship for the
Isobutyl Isobutyrate SIDS Dossier

estimation of rate constants for the gas-phase reaction of OH radicals with organic compounds. J. Inter. Chem. Kinet. 19:799-828.

AOPWIN, v1.91. Atmospheric Oxidation. EPI Suite Version 3.11. U.S. Environmental Protection Agency, 2003.

3.1.2 STABILITY IN WATER

- (a) Preferred result
Type: abiotic
T1/2 pH4:
T1/2 pH7: 9.217 year at 25 degree C
T1/2 pH9:
T1/2 pH8: 336.6 day at 25 degree C
Deg. Product:
Method; other (calculated): HYDROWIN
Year:
GLP: no
Test substance: other TS
Isobutyl isobutyrate
Remark: Rate constant for pH.8 @ 25 deg. C = 2.383e-2 L/mol*sec.
Reliability: Score=2 valid with restrictions
Reference: HYDROWIN, v1.67. Aqueous Base/Acid-Catalyzed Hydrolysis. EPI Suite, v. 3.11, U.S. Environmental Protection Agency, 2003.

3.1.3 STABILITY IN SOIL

No data available

3.2 MONITORING DATA (ENVIRONMENT)

3.3 TRANSPORT AND DISTRIBUTION BETWEEN ENVIRONMENTAL COMPARTMENTS INCLUDING ESTIMATED ENVIRONMENTAL CONCENTRATIONS AND DISTRIBUTION PATHWAYS

3.3.1 TRANSPORT BETWEEN ENVIRONMENTAL COMPARTMENTS

- (a) Type: Volatilization from surface waters
Test substance: Isobutyl isobutyrate
Method: Calculated using EPISUITE v3.11 (USEPA, 2003)
Result: Half-life from model river: 1.67 hours
Half-life from model lake: 4.955 days
Remark: Based on Henry's law constant of 1.58E-3 atm-m³/mol, vapor pressure of 4.33 mm Hg, water solubility of 520 mg/L, and a molecular weight of 144.21 g/mole, and model defaults (for model river: river 1m deep, water flow at 1 m/sec, wind speed of 5 m/sec; for model lake: 1 m deep, water flow 0.05 m/sec., wind speed 0.5 m/sec).
GLP: not applicable
Reference: EPISUITE v3.11, U.S. Environmental Protection Agency (2003).

Isobutyl Isobutyrate SIDS Dossier

- (b) Type: soil or sediment partition coefficient (Koc)
 Test substance: Isobutyl isobutyrate
 Method: Calculated using EPISUITE v.3.11 and PCKOCWIN v.1.66 using structural features of the molecule.
 Result: 53.3 L/kg
 GLP: not applicable
 Reference: PCKOWIN. Version 1.66. Soil Adsorption Coefficient. EPI Suite, v. 3.11, U.S. Environmental Protection Agency, 2003.
- (c) Type: Henry's Law Constant
 Test substance: Isobutyl isobutyrate
 Method: Calculated using water solubility 520 mg/L, vapor pressure 4.33 mm Hg, and molecular weight 144.21 g/mol.
 Result: 1.58E-4 atm-m³/mol
 GLP: not applicable
 Reference: EPI Suite, v. 3.11, U.S. Environmental Protection Agency, 2003.

3.3.2 THEORETICAL DISTRIBUTION (FUGACITY CALCULATION)

- (a) Preferred result
 Type: fugacity model level III
 Media: other: air, water, soil, and sediment
 Air (level III): 12.6%
 Water (level III) 34.4%
 Soil (level III): 52.7%
 Sediment (level III) 0.233%
 Method: other
 Year:
 Remark: Properties used were: melting point -80.7° C, boiling point 148.6°C, vapor pressure 4.33 mm Hg at 25°C, log Kow 2.68, water solubility 520 mg/L, Henry's Law Constant 1.58E-3 atm m³/mol. All other model parameters were default values.
 Remark: Air: half-life = 46.7 hr, emissions = 1000 kg/hr
 Water: half-life = 360 hr, emissions = 1000 kg/hr
 Soil: half-life = 360 hr, emissions = 1000 kg/hr
 Sediment: half-life = 1440 hr, emissions = 0 kg/hr
 Persistence Time: 194 hr
 Reliability: Score = 2, valid with restrictions, calculated
 Reference: Level III Fugacity Model. EPI Suite, v. 3.11, U.S. Environmental Protection Agency, 2003.

3.4 IDENTIFICATION OF MAIN MODE OF DEGRADABILITY IN ACTUAL USE

3.5 BIODEGRADATION

No data were available for isobutyl isobutyrate, so data for isopropyl acetate, isobutyl acetate and 2-ethylhexyl acetate are presented as analog compounds. The alcohol portions of the molecules contain C3, C4, C4, and C8 for the isopropyl acetate, isobutyl isobutyrate, isobutyl acetate, and 2-ethylhexyl acetate, respectively. The biodegradability of isobutyl isobutyrate is expected to be similar to the isobutyl acetate and within the range of acetates ranging from C3 to C8.

- (a) Preferred value
- Type: aerobic
- Inoculum: domestic sewage, non adapted
- Concentration: 3, 7, and 10 mg/L (at least two of these were tested in duplicate)
- Contact time: 20 days
- Degradation: 76% after 20 days
- Result: readily biodegradable
- Kinetic of test subst. 5 day = 61%, 10 day = 72%, 15 day = 74%, 20 days = 76%
- Method: BOD (Standard Methods for the Examination of Water and Wastewater. 1971. 13th Edi. American Public health Association, New York, NY)
- Method: Settled domestic wastewater was filtered through glass wool and added (3 mL/bottle) to clean 300 mL BOD bottles. Aerated dilution water containing minerals specified in the method were added to the bottles along with buffer. Test chemical was added to the bottles. Potential oxygen demand was 3 to 30 mg/L over 20 days. Dissolved oxygen was measured on days 0, 5, 10, 20 using a dissolved O₂ meter. When oxygen decreased to <4 mg/L in any bottle, it was reaerated.
- Year: 1971
- GLP: no data
- Test substance: isopropyl acetate (CAS #108-21-4)
- Remark: Typical unacclimated biodegradation curves for acetates were provided.
The biodegradation curve for isopropyl acetate showed steadily increasing oxidation from test initiation to Day 10, followed by a plateauing through day 20. isopropyl acetate is readily biodegradable. Measured COD was reported as 1.67 mg/mg; the theoretical oxygen demand was reported as 2.04 mg/mg.
- Reliability: Score = 2, valid with restrictions, not all procedures mentioned
- Reference: Price, K.S., G.T. Waggy, and R.A. Conway. 1974. Brine Shrimp Bioassay and Seawater BOD of Petrochemicals. J. Water Pollut. Contr. Fed. 46:63-77.
- (b) Type: aerobic
- Inoculum: domestic sewage, non-adapted
- Concentration: 3, 7, and 10 mg/L (at least two of these were tested in duplicate)
- Contact time: 20 days
- Degradation: 81% after 20 days
- Results: 5 day = 60%, 10 day = 74%, 15 day = 79%, 20 days = 81%
- Method: BOD (Standard Methods for the Examination of Water and Wastewater. 1971. 13th Edi. American Public Health Association, New York, NY)
- Year: 1971
- GLP: no data
- Test substance: Isobutyl acetate (reported as pure chemical)
- Remark: Typical unacclimated biodegradation curves for acetate esters were provided. The biodegradation curve for isobutyl acetate showed steadily increasing oxidation from test initiation to Day 20. Greater than 50% was biodegraded in 20 days; therefore, isobutyl acetate is expected to biodegrade in the environment

Isobutyl Isobutyrate SIDS Dossier

based on a standard BOD study. Measured COD was reported as 1.43 mg/mg; the theoretical oxygen demand was reported as 2.20 mg/mg.

Reliability: Score = 2, Valid with restrictions, not all procedures mentioned
Reference: American Public Health Association. 1971. Standard Methods for the Examination of Water and Wastewater. 1971. 13th Edi, New York, NY.

Price, K.S., G.T. Waggy, and R.A. Conway. 1974. Brine Shrimp Bioassay and Seawater BOD of Petrochemicals. J. Water Pollut. Contr. Fed. 46:63-77.

3.6 BOD₅,COD OR RATIO BOD₅/COD

No data available

3.7 BIOACCUMULATION

(a) BCF: 12.2
Elimination:
Method: other: calculated
Year:
GLP: no
Test substance: isobutyl isobutyrate
Remark: The BCF was calculated using a water solubility of 1000 mg/L at 20°C (Flick, 1991) and a recommended linear regression-derived equation (Lyman, et al. 1990). Based on the estimated BCF, isobutyl isobutyrate is not expected to bioaccumulate in fish or other aquatic organisms (Syracuse Research Corporation).
Reliability: Score = 2, Valid with restrictions, calculation method
Reference: Lyman, W.J., W.F. Reehl, and D.H. Rosenblatt. 1990. Handbook of Chemical Property Estimation Methods. American Chemical Society. Washington, D.C.

(b) BCF: 23.1
Elimination:
Method: other: calculated, BCFWIN
Year:
GLP: no
Test substance: Isobutyl isobutyrate
Remark: Log Kow used in calculation = 2.68 (estimated)
Reliability: Score = 2, Valid with restrictions, calculated procedure
Reference: BCFWIN v.2.15, EPISUITE v. 3.11 (U.S. Environmental Protection Agency, 2003).

3.8 ADDITIONAL REMARKS

A. Sewage Treatment

No data available

B. Other

No data available

4.0 ECOTOXICOLOGICAL DATA

4.1 ACUTE/PROLONGED TOXICITY TO FISH

No ecotoxicity data are available with isobutyl isobutyrate for fish or algae, so data for isopropyl and propyl acetate, n-butyl acetate and 2-ethylhexyl acetate are presented as analog compounds. The alcohol portions of the molecules contain C3, C3, C4, C4, and C8 for the isopropyl and propyl acetate, isobutyl isobutyrate, n-butyl acetate, and 2-ethylhexyl acetate, respectively. Since it is not possible to assign any specific analog compound to isobutyl isobutyrate, the “read-across” approach is used. Thus, the ecotoxicity of isobutyl isobutyrate is expected to be within the range of acetates ranging from C3 to C8.

(a) Remark:	Due to the absence of acute fish toxicity data for isobutyl isobutyrate, data for a structural analog are used (2-ethylhexyl acetate, CAS# 103-09-3).
Preferred Value	
Type:	static renewal
Species:	<i>Oncorhynchus mykiss</i> (Fish, fresh water)
Exposure period:	96 hour(s)
Unit:	mg/L
LC50:	8.27
Limit test:	no
Analyt. monitoring:	yes
Method:	OECD guideline 203 (specific protocol titled, "Acute Toxicity of the Water Accomodated Fraction (WAF) for 2-Ethylhexyl Acetate to the Rainbow Trout, <i>Oncorhynchus mykiss</i> , Determined Under Static Test Conditions, and Amendment"
Year:	2002
GLP:	yes
Test substance:	2-ethylhexyl acetate (CAS RN 103-09-3), analog for isobutyl acetate, 99.7% purity, source: Aldrich Chemical Co., Milwaukee, WI, USA.
Method:	Rainbow trout were obtained as eyed embryos from a commercial supplier. Fish were maintained in blended laboratory freshwater. After hatch and swim-up and absorption of the yolk sac, fish were fed brine shrimp and commercial fish food twice daily. A subset of fish were removed from the main population and acclimated to test dilutio water temperature (~15 deg. C) for 72 hours before testing Fish used in the toxicity tests were about 130 days old, had a mean length of 39 mm and a mean weight of 0.368 g. Fish were not fed during the test. The toxicity test was conducted using 4-L vessels. Test volume was 3 L. During

Isobutyl Isobutyrate SIDS Dossier

testing, vessels were placed in a 15±1 deg. C water bath. Light was maintained on a 16h daylight cycle. Fish were not fed during the test. No aeration was employed during the test.

The control/dilution water was a combination of naturally hard well water and well water that was de-mineralized by reverse osmosis yielding the desired range of hardness (total hardness ranging from 130 to 160 mg/L as CaCO₃).

The test employed a water accommodated fraction approach (WAF) in which an 9.0 gram of test material was placed on the surface of water in a 9.5 liter carboy filled with 9 liter dilution water. The solution was stirred for 20 hours with a stir bar creating a vortex <25% solution depth. After stirring the solution was allowed to separate for one hour. The aqueous solution was siphoned from the bottom of the carboy for use. Fresh solutions were prepared daily.

Test concentrations were analyzed daily by gas chromatography (GLC) equipped with a flame ionization detector (GC-FID).

The test was conducted in 4 liter glass jars. Two replicates of 5 fish each were exposed to the control/dilution water and to each of six dilutions of the WAF solution (1.6, 3.3, 6.5, 13, 25, 50% WAF). Renewals were made daily. Mortality and abnormal signs of behavior were recorded daily.

Results:

Water quality parameters measured during the test included: total hardness, 164 mg/l (as CaCO₃); alkalinity, 162 mg/l (as CaCO₃); dissolved oxygen ranged from ~100% saturation at renewal of test solution to 45% saturation mg/l; temperature ranged from 14 to 16 deg. C; and pH ranged from 7.61 to 8.26.

Test solutions were observed to be clear and colorless with no visible particulates, surface film, undissolved test substance, or precipitate.

Measured concentrations in the new solutions ranged from 0.41 to 16.7 mg/L across treatments. Measured concentrations in old solutions (24-h) ranged from <0.347 to 13.8 mg/L. Mean measured concentrations (using ½ the MDL for non-detects) were 0.284, 0.570, 1.34, 2.51, 6.42, and 15.3 mg/L across all treatments.

Results of the 96-h acute toxicity test, by concentration level tested:

Control: No mortality

0.284 mg/L: No mortality or sublethal effects

0.57 mg/L: No mortality or sublethal effects

1.34 mg/L: No mortality or sublethal effects

2.51 mg/L: No mortality or sublethal effects

Isobutyl Isobutyrate SIDS Dossier

6.42 mg/L: Partial mortality (20%) at 6.42 mg/L at 96 hours, all remaining fish (8/10) showed sublethal effects;
15.3 mg/L: 100% mortality in both replicates at 24 hours.

The 96-h LC50 was reported as 8.27 mg/l using the Trimmed Spearman-Kärber method. Confidence limits were 6.58 to 10.4 mg/L.

Reliability: Score = 1, valid without restrictions
Reference: Hahne R. 2002a. Acute Toxicity of the Water Accommodated Fraction (WAF) for 2-Ethylhexyl Acetate to the Rainbow Trout, *Oncorhynchus mykiss*, Determined under Static Test Conditions. Report #46955 prepared by ABC Laboratories Inc., Columbia Missouri. May 22, 2002.

(b) Type: flow through
Species: *Oncorhynchus mykiss* (Fish, fresh water)
Exposure period: 96 hour(s)
Unit: mg/l
Analytical monitoring: yes
NOEC: > 4.2
LC0: > 4.2
LC50: > 4.2
LC100: > 4.2
Limit Test: yes

Method: OECD Guide-line 203 "Fish, Acute Toxicity Test"
Year: 1992
GLP: yes
Test substance: other TS: 2-ethylhexyl acetate (BASF), purity : 99.3 %

Result: In conclusion the 96 hour LC50 values for 2-Ethylhexylacetate in the Rainbow trout (*Oncorhynchus mykiss*) were >4.5 mg/L based on the nominal concentration of the test substance and >4.2 mg/L based on the mean of analytically determined concentration.

No lethality was observed up to the solubility limit of the test substance in water (3.9 mg/L).

Test conditions: 0, 4.5 mg/L based on technical test substance. The test was carried out slightly above the solubility limit of the test substance in water (3.9 mg/L).

Test organisms
Test species: Rainbow trout (*Oncorhynchus mykiss*)
Animal supplier: Forellenzucht Trostadt GbR, Dorfstrasse 7,98646 Trostadt, Germany
Body weight: (wet weight): 2.08 g (1.21 - 3.46 g)

Body length (from the top of the snout to the end of the caudal fin): 6.1 cm (5.1 - 7.0 cm)
Age of the animals: approx. 5 months
Hatching date: Sep 08, 2003
Arrival at the testing facility: Dec 04, 2003

Remark: All animals used for the study were derived from the same batch of fish. All test animals were observed for their health state during insertion into the test vessels. No signs for sickness, injuries or abnormalities were observed.

EXPERIMENTAL PROCEDURE

1. Acclimatization

Duration of acclimatization to testing conditions including light regime: 14 days, acclimatization to test water and temperature: 8 days

Photoperiod: 16 hours light, 8 hours dark

Water quality: Non chlorinated charcoal filtered tap water

(Frankenthal, Germany) mixed with deionized water

Total hardness: Approx. 1.0 mmol/L = 100 mg/L CaCO₃

Acid capacity: Approx. 2.2 mmol/L

Oxygen content: > 80% saturation

pH-value: Generally 7.5 - 8.5

Temperature: Approx. 15°C (8 days)

Diet: Ecostart 17 (Bio Mar), supplier Kofu Tiernahrung, Betriebsstätte Wesel, Hafenstr. 11 - 13, 46483 Wesel, Germany, ad libitum, additionally generally on workdays frozen brine shrimp (artemia).

Withdrawal of feed: During the last day before start of exposure

Medical treatment: None during acclimatization

Mortality during the last week before start of exposure: 0 %

Test water

Water quality: Non chlorinated charcoal filtered tap water (Frankenthal, Germany, aerated) mixed with deionized water

Total hardness: Approx. 1.0 mmol/L = 100 mg/L CaCO₃

Conductivity: Approx. 250 µS/cm (at 25°C)

Ca content: Approx. 40 mg/L

Mg content: Approx. 5 mg/L

Acid capacity: Approx. 2.2 mmol/L

pH-value: Generally 7.7 - 8.1

Temperature: Approx. 15°C

The tap water is regularly assayed for chemical contaminants by the municipal authorities of Frankenthal and the Technical Services of BASF Aktiengesellschaft as well as for presence of microbes by a contract laboratory. On the basis of the analytical findings, the drinking water was found to be suitable for toxicity tests. German Drinking Water Regulation (Trinkwasserverordnung, Bundesgesetzblatt, December 05, 1990) served as a guideline for maximum tolerable contaminants. The individual results are to be found in the archives of the testing facility.

Preparation of the test concentrations:

The test was conducted in a flow through system. The flow-through-system started 3 days before the fishes were placed into the test vessels to ensure that the test system was saturated with the test substance. The stock solution was prepared in the following way: The test substance was weighed into a volumetric flask. The volumetric flask was made up with acetone. The stock solution was completely dissolved and clear. The stock solution was transferred into a 50 mL glass injector which was attached to a perfusion pump. The stock solution was continuously injected into the mixing vessel, where it was admixed to a continuous flow of dilution water. The mixture of dilution water and the stock solution of the test substance was flowing continuously into the test vessels. The flow rate of the dilution water was calibrated (maximum deviation less than 10%) before the system was started. The test vessels were tempered by the dilution water. The apparatus was checked daily for proper function. The flow meter (rotameter) levels were checked daily and corrected if necessary. The control group was performed as solvent control with the same concentration of acetone as the concentration group. A stock solution of 62.5 mL acetone in 2 L dilution water was mixed at a flow rate of 60 mL / hour with 18.69 L dilution water / hour. The concentration of acetone in all test vessels was 0.1 mL acetone / L test solution.

The nominal test concentration was 4.5 mg/L.

Test conditions

Isobutyl Isobutyrate SIDS Dossier

The exposure was carried out in a flow-through system.

Test vessels: glass aquaria (59 cm long, 28.5 cm wide, 30 cm high) with an overflow 26.5 cm above the bottom,

overflow covered with plastic gauze

Volume of water: 45 L

Flow rate: 18.75 L / hour

Volume of exchange/day: Approx. 10

Animals / test vessel: 10

Loading (g fish/L water): 0.5

Loading (g fish/L water/day): 0.05

Test vessel / concentration: 1

Photoperiod: 16 hours light, 8 hours darkness

Light intensity: Approx. 82 - 280 Lux (the light intensity is determined in regular intervals at the surface of the aquaria)

Temperature: 15°C

Aeration: None

Feeding: None

Insertion of test organisms

The test organisms were introduced into the test vessels according to a randomization plan prepared by the testing laboratory using a program of the laboratory data evaluation group of the testing facility.

Observations

Symptoms and lethality:

The fish were observed for survival and toxic signs changes in appearance, swimming behavior and behavior in comparison to the control group) within 1 hour after start of exposure and 4, 24, 48, 72 and 96 hours after start of exposure. Fish were considered dead if there was no visible movement and no reaction after touching. Dead fish were removed from the test vessels.

Water parameter:

Temperature, oxygen content and pH-value were measured in each test vessel within 1 hour after start of exposure and after 24, 48, 72 and 96 hours.

Analytical determinations

An analytical method for the determination of the test substance in the test water was developed by the Analytical Department of BASF Aktiengesellschaft. The analyses were carried out at the Analytical Chemistry Laboratory of the Experimental Toxicology and Ecology of BASF

Aktiengesellschaft under the responsibility of Dr. E. Leibold / Lan Ma in compliance with the Principles of Good Laboratory Practice. For methods and details see Appendix.

Stability in water: See analytical concentration control

Samples for analytical determination of the concentration were taken within the first hour after the start of exposure, after 24 hours and at the end of the exposure after approx. 96 hours. Water samples were taken from the middle of the test vessel using a beaker and were transported to the analytical laboratory in glass ampoules, which were rinsed with test solution before they were filled. The transport to the Analytical Laboratory was done on the day of sampling.

Statistics:

No statistical analysis was carried out since no lethality was observed at the highest tested concentration. The median lethal concentration (LC50) was above this value.

Test substance: 2-Ethylhexylacetate (BASF), purity: 99.3 %
Reliability: (1) valid without restrictions
Reference: BASF AG, Product Safety Department, unpublished study, project no. 12F0674/025053 (to be finalized).

(c) Value: 9.455 mg/L
Test substance: isobutyl isobutyrate (CAS # 97-85-8)
Remark: An acute fish 96-h LC50 was calculated using ECOSAR, from the USEPA. The SAR for esters was used. The structure was determined from the CAS RN as stored in the accompanying database of SMILES notations within ECOSAR.
Reliability: Score = 2, valid with restrictions, calculated value
Reference: EPA's EcoSAR model (v. 0.99g). EPWIN (Estimation Program Interface for Windows). Version 3.10. U.S. Environmental Protection Agency (2000).

(d) Value: 3.057 mg/L
Test substance: 2-Ethylhexyl acetate (CAS # 103-09-3)
Remark: An acute fish 96-h LC50 was calculated using ECOSAR, from the USEPA. The SAR for esters was used. The structure was determined from the CAS RN as stored in the accompanying database of SMILES notations within ECOSAR.
Reliability: Score = 2, valid with restrictions, calculated value

Isobutyl Isobutyrate SIDS Dossier

Reference: EPA's EcoSAR model (v. 0.99g). EPI Suite v3.10 (USEPA, 2000)

4.2 ACUTE TOXICITY TO AQUATIC INVERTEBRATES

Daphnia

(a) Preferred result
Type: static renewal
Test substance: Isobutyl isobutyrate (Eastman IBIB) [CAS No.97-85-8]; 99.6% purity
Species: *Daphnia magna* (Crustacea)
Exposure period: 48 hour(s)
Unit: mg/L
NOEC: 53.7
EC50: 55.8-59.3
Analytical monitoring: yes
Method: OECD 202 and EEC/Annex V C.2
Year: 1995
GLP: yes
Test substance: isobutyl isobutyrate (CAS# 97-85-8)
Method: First-instar *Daphnia magna* neonates (<24 hours old) were obtained from an in-house culture. Daphnids were cultured in a temperature controlled water bath at 20 degrees C and maintained on a 16 hour light cycle. During holding, daphnids were fed algae and occasionally artificial food. Dilution water was an treated, filtered and aerated water from Lake Ontario. Total hardness and alkalinity were 120 and 91 mg/L as CaCO₃, respectively. It was biologically aged by including aquatic organisms in the stored water prior to testing.

The toxicity test was conducted using 250 mL beakers filled to 200 mL with test solution. During testing, vessels were place in a 20±1 deg. C water bath. Light was maintained on a 16h daylight cycle. Daphnids were not fed during the test. No aeration was employed during the test.

Exposure solutions were prepared daily by adding the appropriate amounts of test material to 500-mL volumetric flasks. Aliquots of 200 mL were added to each vessel.

Test concentrations were analyzed daily by gas chromatography (GLC) equipped with a flame ionization detector (GC-FID).

Two replicates of 10 daphnids each were exposed to the control/dilution water and to each of five concentrations of the TS (95.2, 171.5, 308.6, 555.5, and 1000 mg/L). Renewals were

Isobutyl Isobutyrate SIDS Dossier

Results: made daily. Numbers of immobilized daphnids and abnormal signs of behavior were recorded at 0, 6, 24, and 48 hours. Water quality parameters measured during the test included: dissolved oxygen, which ranged from 8.4 to 9.2 mg/L during the test; temperature, which ranged from 20 to 21 deg. C; and pH, which ranged from 6.9 to 9.2. Measured concentrations in the new and 24-h solutions were 53.72-55.82, 70.3-82.8, 109.1-131.35, 227.08-285.2, and 178.15 mg/L (range of two replicates).

Results of the 48-h acute toxicity test, by concentration level tested (for each replicate):

Control: 0, 1 mortality
53.72-55.82 mg/L: 3, 5 mortalities
70.3-82.8 mg/L: 10 (at 24 hours), 10 mortalities
109.31-131.35 mg/L: 10 (at 24 hours), 10 mortalities
227.08-285.2 mg/L: 10 (at 24 hours), 10 mortalities
178.15 mg/L: 10, 10 mortalities (all by 24 hours)

The 48-h LC50s were calculated using the nonlinear interpolation method. Confidence limits were 53.7-82.8 for replicate series A and inestimable to 70.3 mg/L for replicate B.

Reliability: No sublethal effects were reported.
Score = 1, valid without restrictions
Reference: Patterson KM, Hirsch MP. 1995. An Acute Aquatic Effects Test with the Daphnid *Daphnia magna*. Study #EN-431-900945-1, Eastman Kodak Company, Health and Environment Laboratories, Rochester, NY. August 3, 1995.

- (b) Data presented because no data for fish and algae are available for isobutyl isobutyrate. Data for the structurally similar 2-ethylhexyl acetate (CAS# 103-09-3) are presented to support the endpoints for isobutyl isobutyrate.

Type: static
Species: *Daphnia magna* (Crustacea)
Exposure period: 48 hour(s)
Unit: mg/l
NOEC: = 15.7
EC50: = 22.9
Analytical monitoring: yes
Method: other: OECD 202 (Part 1)
Year: 2002
GLP: yes
Test substance: 2-ethylhexyl acetate (CAS# 103-09-3)
Method: First-instar *Daphnia magna* neonates (<24 hours old) were obtained from an in-house culture. Daphnids were cultured in a temperature controlled water bath at 20 degrees C and maintained on a 16 hour light cycle. During holding, daphnids were fed algae and occasionally artificial food. Dilution water was an aged laboratory freshwater. The control / dilution water was a combination of naturally hard well water and well water

Isobutyl Isobutyrate SIDS Dossier

that was de-mineralized by reverse osmosis yielding the desired range of hardness (total hardness ranging from 130 to 160 mg/L as CaCO₃). It was biologically aged by including aquatic organisms in the tank.

The toxicity test was conducted using 250 mL beakers filled to 200 mL with test solution. During testing, vessels were placed in a 20±2 deg. C water bath. Light was maintained on a 16h daylight cycle. Daphnids were not fed during the test. No aeration was employed during the test.

The test employed a water accommodated fraction approach (WAF) in which an 9.0 gram of test material was placed on the surface of water in a 9.5 liter carboy filled with 9 liter dilution water. The solution was stirred for 20 hours with a stir bar creating a vortex <25% solution depth. After stirring the solution was allowed to separate for one hour. The aqueous solution was siphoned from the bottom of the carboy for use. Fresh solutions were prepared daily.

Test concentrations were analyzed daily by gas chromatography (GLC) equipped with a flame ionization detector (GC-FID).

Two replicates of 10 daphnids each were exposed to the control/dilution water and to each of six dilutions of the WAF solution (3.3, 6.5, 13, 25, 50, and 100% WAF). Renewals were made daily. Numbers of immobilized daphnids and abnormal signs of behavior were recorded daily.

Result: Water quality parameters measured during the test included: total hardness, 148 mg/l (as CaCO₃); alkalinity, 164 mg/l (as CaCO₃); dissolved oxygen ranged from 101 to 117% of saturation during the test; temperature ranged from 19.7 to 22.0 deg. C; and pH ranged from 7.99 to 8.35.

Test solutions were observed to be clear and colorless with no visible particulates, surface film, undissolved test substance, or precipitate.

Measured concentrations in the new solutions ranged from 0.951 to 45.8 mg/L across treatments. Measured concentrations in old solutions (24-h) ranged from <0.688 to 28.1 mg/L. Mean measured concentrations (using ½ the MDL for non-detects) were 0.828, 2.06, 4.12, 7.99, 15.7, and 33.4 mg/L across all treatments.

Results of the 48-h acute toxicity test, by concentration level tested:

Control: No mortality
0.828 mg/L: No mortality
2.06 mg/L: No mortality
4.12 mg/L: No mortality

Isobutyl Isobutyrate SIDS Dossier

7.99 mg/L: No mortality
15.7 mg/L: No mortality;
33.4 mg/L: 50% mortality in both replicates at 24 hours, 100% mortality at 48 hours.

No sublethal effects were reported.

The 48-h LC50 was reported as 22.9 mg/l using the Trimmed Spearman-Kärber method. Confidence limits were 15.7 to 33.4 mg/L.

- Test substance: 2-Ethylhexyl acetate [CAS No.103-09-3]; 99.77% purity, source: Aldrich Chemical Co., Milwaukee, WI, USA.
- Reference: Hahne R. 2002b. Acute Toxicity of the Water Accomodated Fraction (WAF) for 2-Ethylhexyl Acetate to the Water Flea *Daphnia magna*, Determined under Static Test Conditions. Report #46956 prepared by ABC Laboratories Inc., Columbia Missouri. May 02, 2002.
- (c) Value: 27.566 mg/L
Test substance: Isobutyl isobutyrate (CAS # 97-85-8)
Remark: An acute daphnid 48-h LC50 was calculated using ECOSAR, from the USEPA. The SAR for esters was used. The structure was determined from the CAS RN as stored in the accompanying database of SMILES notations within ECOSAR.
Reliability: (2) valid with restrictions, calculated value
Reference: EPA's EcoSAR model (v. 0.99g). EPWIN (Estimation Program Interface for Windows). Version 3.10. U.S. Environmental Protection Agency (2000).
- (d) Value: 3.571 g/L
Test substance: 2-ethylhexyl acetate (CAS# 103-09-3)
Remark: An acute daphnid 48-h LC50 was calculated using ECOSAR, from the US EPA. The SAR for esters was used. The structure was determined using the CAS RN as stored in the accompanying database of SMILES notations within ECOSAR.
Reliability: Score = 2, valid with restrictions, calculated value.
Reference: EPA's ECOSAR model (v. 0.99g). EPISuite v. 3.10 (US EPA, 2000).

4.3 TOXICITY TO AQUATIC PLANTS e.g. Algae

- (a) Data presented because no data for fish and algae are available for isobutyl isobutyrate. Data for the structurally similar 2-ethylhexyl acetate (CAS# 103-09-3) are presented to support the endpoints for isobutyl isobutyrate.
- Preferred result
- Species: *Selenastrum capricornutum* (Freshwater green algae)
- Test substance: 2-Ethylhexyl acetate [CAS No.103-09-3]; 99.77% purity, source: Aldrich Chemical Co., Milwaukee, WI, USA.
- Endpoint: growth inhibition (based on area under growth curve), cell counts
- Exposure period: 72 hour(s)
- Units: mg/L

Isobutyl Isobutyrate SIDS Dossier

Limit test: no
EC50: >21.9
NOEC: 10.3
LOEC: 21.9
Analytical monitoring: yes
Method: OECD Guideline 201
GLP: yes
Test substance: 2-Ethylhexyl acetate (CAS# 103-09-3)
Method: The green alga *Selenastrum capricornutum* used was maintained in the laboratory and originated from the Univ. Texas Culture Collection (UTEX #1648), Austin, TX. Cultures were maintained in synthetic algal nutrient medium in flasks under continuous cool-white fluorescent illumination of ~8000 lumens/m² at 24±2°C and continuously shaken at 100 oscillations/minute.

Test medium was prepared by adding requisite amounts of each of the macro- and micro-nutrients into laboratory-grade water. After pH adjustment to 7.5±0.1 S.U., the media was filtered using a 0.45 micrometer porosity filter and refrigerated.

The test employed a water accommodated fraction approach (WAF) in which an 3.0 gram of test material was placed on the surface of water in a 4 liter carboy filled with 3 liter dilution water. The solution was stirred for 20 hours with a stir bar creating a vortex <25% solution depth. After stirring the solution was allowed to separate for one hour. The aqueous solution was siphoned from the bottom of the carboy for use and was as the 100% WAF.

Test vessels were sterilized 250-mL flasks fitted with foam stoppers. Besides six control replicates, three replicates were prepared for each test concentration. Replicates contained 100 mL test solution. Nominal test concentrations of 6.5, 13, 25, 50, and 100% WAF were chosen based on the results of two range finding tests.

Each flask (except for a fourth flask of the lowest concentration used for 72-h chemical analysis) was inoculated with 1 mL of the algae containing approximately 1.0 E+6 cells/mL, resulting in an initial cell density of approximately 1.0 E+4 cells/mL. Flasks were kept under the same conditions as the stock cultures. Cell counts were performed using a light microscope and a hemacytometer every 24 hours in all flasks. Results were expressed as cells/mL.

Temperature and pH was measured in each test vessel and control at test initiation and in one replicate of the control and each treatment at 72 hours.

Samples of the initial test solutions were analyzed for TS concentration. A fourth replicate at the lowest concentration was analyzed at 72 hours. Chemical analysis was performed using gas chromatography (GC) equipped with a Flame Ionization Detector (FID).

Final samples were prepared by extracting a 10-mL sample twice with toluene and shaken for two minutes. Samples were vialled and immediately analyzed.

EC and NOEC values were calculated based on area under the growth curve and growth rate versus mean measured concentration. Shapiro-Wilk's test and Levene's test were conducted to test for normality and homogeneity of variances. A one-way ANOVA and a Dunnett's comparison to the control was conducted for each time point to determine the NOECs.

Results:

Temperature ranged from 23.5 to 25.0°C at 0 and 72 hours. The pH values ranged from 7.40 to 7.57 S.U. at test initiation across all test vessels and pH ranged from 8.97 to 9.26 S.U. at 72 hours across all vessels. Although the pH increased by more than 1 pH unit in some flasks, the integrity of the test was deemed unaffected, since the control performance was unaffected (control pH at 72 hours was 9.03 S.U.)

Measured concentrations on day 0 ranged from 2.49 to 43.4 mg/L across treatments (6.5 to 100% WAF). Measured concentrations in day 3 solutions were all <0.688. Mean measured concentrations (using ½ the MDL for non-detects) were 1.42, 2.70, 5.27, 10.3, and 21.9 mg/L across all treatments.

At 72 hours, the control had 116 E+4 cells/mL, an increase of 116 times that at test initiation indicating an acceptable test. The percent change in cell density ranged from -3% in the 25% and 100% WAF treatments to +14% in the 13% WAF treatment. At 72 hours, no significant reduction in algal growth was found in treatments #50% WAF as measured by area under the growth curve and #100% WAF as measured by growth rate.

72-h EC50 = >21.9 mg/L (based on area under the growth curve and growth rate)

72-h NOEC = 10.3 mg/L, 72-h LOEC = 21.9 mg/L

Reliability:

Score = 1, valid without restrictions

Reference:

Hicks SL. 2002. Toxicity of 2-Ethylhexyl Acetate to the Unicellular Green Alga, *Selenastrum capricornutum*. Report #46957 prepared by ABC Laboratories Inc., Columbia Missouri. May 01, 2002.

(b)

Value:

0.771 mg/L

Test substance:

Isobutyl isobutyrate (CAS# 97-85-8)

Isobutyl Isobutyrate SIDS Dossier

- Remark: An acute algal 96-h EC50 was calculated using ECOSAR, from the USEPA. The SAR for esters was used. The structure was determined from the CAS RN as stored in the accompanying database of SMILES notations within ECOSAR.
- Reliability: (2) valid with restrictions, calculated value
- Reference: EPA's EcoSAR model (v. 0.99g). EPI Suite v3.10 (USEPA, 2000)
- (c) Value: 0.26 mg/L
- Remark: An acute algal 96-h EC50 was calculated using ECOSAR, from the USEPA. The SAR for esters was used. The structure was determined from the CAS RN as stored in the accompanying database of SMILES notations within ECOSAR.
- Test substance: 2-Ethylhexyl acetate [CAS No.103-09-3]
- Reliability: (2) valid with restrictions, calculated value
- Reference: EPA's EcoSAR model (v. 0.99g). EPI Suite v3.10 (USEPA, 2000)

4.4 TOXICITY TO BACTERIA

No data available

4.5 CHRONIC TOXICITY TO AQUATIC ORGANISMS

No data available

4.5.1 CHRONIC TOXICITY TO FISH

No data available

4.5.2 CHRONIC TOXICITY TO AQUATIC INVERTEBRATES

No data available

4.6 TOXICITY TO TERRESTRIAL ORGANISMS

No data available

4.6.1 TOXICITY TO SOIL DWELLING ORGANISMS

No data available

4.6.2 TOXICITY TO TERRESTRIAL PLANTS

No data available

4.6.3 TOXICITY TO OTHER NON MAMMALIAN TERRESTRIAL SPECIES (INCLUDING AVIAN)

No data available

4.7 BIOLOGICAL EFFECTS MONITORING (INCLUDING BIOMAGNIFICATION)

No data available

4.8 BIOTRANSFORMATION AND KINETICS

No data available

4.9 ADDITIONAL REMARKS

No additional remarks

5.0 TOXICITY

5.1 ACUTE TOXICITY

5.1.1 ACUTE ORAL TOXICITY

Isobutyl isobutyrate (IBIB) is rapidly broken down to form equal amounts of isobutyric acid and isobutanol. The resulting isobutanol is further broken down to ultimately form isobutyric acid. In order to support the limited acute oral toxicity information on IBIB, acute oral toxicity data for the two metabolites (isobutanol and isobutyric acid) have been added to this section.

- (a) Preferred result
- | | |
|-----------------|---|
| Type: | LD50 |
| Species: | rat |
| Value: | >6400 mg/kg |
| Method: | The test material was administered undiluted to seven animals at dose levels ranging from 200 to 12800 mg/kg. One rat was used at each dose level. The animals were observed for 14 days following dosing and no necropsies were performed. |
| Year: | 1956 |
| GLP: | no |
| Test substance: | isobutyl isobutyrate |
| Remark: | Weakness, and ataxia were observed in the animal receiving 12800 mg/kg. The time of death was the 2 nd day following dosing. |
| Reliability: | Score = 2, acceptable w/restrictions |
| Reference: | Laboratory of Industrial Medicine, Eastman Kodak Co., 1956 |
- (b) Type: LD50
- | | |
|-----------------|--|
| Species: | mouse |
| Value: | >6400 mg/kg |
| Method: | The test material was administered undiluted to seven animals at dose levels ranging from 1600 to 25600 mg/kg. One mouse was used at each dose level. The animals were observed for 14 days following dosing and no necropsies were performed. |
| Year: | 1956 |
| GLP: | no |
| Test substance: | isobutyl isobutyrate |
| Remark: | Weakness, and ataxia were observed in the animal receiving 12800 mg/kg. The time of death was within 3 hours following dosing. |
| Reliability: | Score = 2, acceptable w/restrictions |
| Reference: | Laboratory of Industrial Medicine, Eastman Kodak Co., 1956 |

- (c) Type: LD50
Species: rat
Value: > 2830 mg/kg bw (males)
3350 (2860 to 3920) mg/kg bw (females)
- Method: EPA (TSCA) Health Effects Testing Guidelines 40 CFR Part 798 (Subpart B, Section 798.1175:acute oral toxicity; and 1987 OECD Guidelines for Testing of Chemicals (Section 4: Health Effects; 401:acute oral toxicity). Rat (Harlan Sprague Dawley) body weights were within +/- 20% of the group mean for each sex. The body weight range on the day of dosing was 281 to 292 g for males and 210 to 259 g for females (including those used for preliminary testing). A total of 3 male and 20 female rats were used for the definitive peroral test. An additional 2 female rats were used for preliminary testing. The animals were acclimated for at least 5 days before dosing. Detailed clinical observations were conducted twice, at the time of receipt and during animal identification and/or dosing. Cage-side observations and mortality checks were conducted at least once daily. Animals considered unacceptable for the study, based on the clinical signs were rejected for use on this study. Each dosing mixture was prepared just prior to administration by diluting the appropriate amount of isobutanol with 0.25% w/v aqueous methyl cellulose solution. All resulting emulsions were mixed for approximately 15 to 30 minutes on a magnetic stirrer. Doses were administered by stomach intubation through a commercial 16-gauge (3-inch) ball-end stainless steel needle attached to a disposable syringe. The exact amounts of test substance and emulsion given to each rat were recorded on the raw data form. The rats were fasted overnight before dosing. Five female rats were included on each of several dose levels in order to determine an LD₅₀. Three male rats were included on an intermediate dose level for comparison. An additional 2 female rats were used for preliminary peroral toxicity testing. For individual animals, the dosing volume was adjusted according to body weight. Dosed rats were observed frequently for signs of toxicity on the first day of the test and twice a day thereafter (except on weekends or holidays when they were examined for death alone). Weights were recorded on the day of dosing and at 7 and 14 days after dosing or at death. After 14 days, all survivors were sacrificed by CO₂ overdose. Necropsies were performed on all animals that died or were sacrificed. Unless tissues were judged to be excessively autolyzed, the following tissues were collected from selected animals and retained in 10% neutral buffered formalin: kidneys, urinary bladder, liver, sciatic nerve, stomach, intestines and spleen. Lungs were also saved because of possible lung damage, based on clinical signs. An LD₅₀ was calculated for female rats, based on the 14-day observation period. It was calculated by the moving average method. An estimate of the slope was made by the formula developed by Weil. During the acute peroral toxicity test, several animals (including survivors) had varying amounts of

Isobutyl Isobutyrate SIDS Dossier

blood present in the urine. Therefore, histology evaluations were performed on all saved kidney and urinary bladder tissues. One female rat appeared to be pregnant at necropsy and the uterus was saved in order to verify this condition (since the animals are ordered to be non-pregnant).

Year: 1993

GLP: Yes

Test

substance: Isobutanol (99.9% purity by capillary GC, GC/MS and NMR used to confirm identity)

Remark: In preliminary testing, 1 female rat was dosed with 2000 mg/kg of isobutanol and 1 female rat was dosed with 8000 mg/kg (20% w/v emulsions in 0.25% aqueous methyl cellulose solution). The rat receiving 8000 mg/kg died. In the definitive test, the peroral LD₅₀ for female rats dosed with the test substance (emulsions in 0.25% aqueous methyl cellulose solution) was 3350 mg/kg. None of 3 male rats died after receiving peroral doses of 2830 mg/kg of isobutanol (a comparison dose that produced 0 of 5 female deaths), although signs were apparent. Signs of toxicity included sluggishness, unsteady gait, lacrimation, piloerection, slow breathing, prostration and a trace to large amount of blood in urine (positive by HEMASTIX Reagent Strips). Several females exhibited a slight weight loss within 7 to 14 days. Deaths occurred within 2 hours to 1 day. Survivors recovered within 0.5 hour to 6 days. Necropsy of animals that died revealed discolored and/or mottled lungs (bright to dark red), tan to dark maroon and/or mottled livers (in 2), discolored stomachs (gray and/or yellow), 1 liquid-filled stomach, dark red and/or gray areas on the intestines, red to brown kidneys (in 1) and a large amount of blood in the urine of 1 (positive by HEMASTIX Reagent Strips). There were no gross lesions apparent in any survivor at necropsy. One female survivor dosed with 2830 mg/kg of isobutanol appeared pregnant at necropsy (determined to be a pseudopregnancy during microscopic valuation). The kidneys and urinary bladders from 1 or 2 rats from each dose group (except 1000 mg/kg) were saved and examined microscopically (see Appendix 2). The only kidney lesions evident were single instances of tubular proteinosis, tubular basophilia, mineralization and congestion, which were not considered to be attributable to the test substance. There were no lesions observed in the urinary bladders. In the uterus of the 1 female rat (2830 mg/kg) that appeared pregnant at necropsy, deciduoma of pseudopregnancy were apparent. This condition is somewhat unusual for animals of this age group. Subsequent investigations revealed that the female rats ordered for this study had undergone vaginal swabbing on the day of shipment at the animal supplier. This female animal (and one other from the acute inhalation study) had pseudopregnancy due to cervical stimulation from the vaginal swabbing procedure. The male rat oral (fasted) LD₅₀ was > 2830 mg/kg bw; 0 of 3 died. The female rat oral (fasted) LD₅₀ was 3350 (2860 to 3920) mg/kg bw.

Isobutyl Isobutyrate SIDS Dossier

Microscopic kidney lesions were evident but probably not related to treatment.

Reliability: Score = 1, GLP guideline study for Isobutanol
Reference: Christopher, S.M. November 30, 1993. "Isobutanol: Acute toxicity and irritancy testing using the rat (peroral and inhalation toxicity) and the rabbit (cutaneous and ocular tests)". Bushy Run Research Center, Union Carbide Corp. Lab. Proj. ID 92U1166

(d) Type: LD50
Species: rat
Value: >500 mg/kg
Method: Ten male albino rats (Charles River CD strain) were fasted 18 hours prior to receiving 500 mg/kg of isobutyric acid as a 2% suspension in corn oil. Clinical signs were observed on the day of dosing and daily during the 14-day observation period. The initial and terminal body weights were determined on the test animals. A gross autopsy was not performed.

Year: 1979
GLP: no data
Test substance: isobutyric acid
Remark: No signs of intoxication were observed and all animals gained weight normally during the 14 day observation period. This study was conducted in support of the use of isobutyric acid as a fumigant in dry corn to prevent growth of mold and fungus.

Reliability: Score = 2, accepted scientific standard methods – For Isobutyric acid
Reference: Report No. A79-146. May 25, 1979. Toxicity Data Report. Animal Industry R/D, Agricultural Division, American Cyanamid Co. P.O. Box 400, Princeton, NJ 08540

5.1.2 ACUTE INHALATION TOXICITY

(a) Preferred result

Type: LC66 and LC0
Species: rat
Value: LC66 = 5423 ppm, LC0 = 658 ppm
Method: Groups of three rats were exposed to an atmospheres containing aerosolised isobutyl acetate at a concentration of either 5423 or 658 ppm for 6 hours. Exposure values were based on nominal concentrations.

Year: 1956
GLP: no
Test substance: isobutyl isobutyrate
Remark: No effects were noted in the rats exposed to 658 ppm for 6 hours. Clinical signs noted in the 5423 ppm group included prostration and narcosis. Deaths occurred during exposure. All animals that

survived the exposure gained weight at the end of the 14 day observation period.
Reliability: Score = 2, accepted scientific methodology
Reference: Laboratory of Industrial Medicine, Eastman Kodak Co., 1956

5.1.3 ACUTE DERMAL TOXICITY

(a) Preferred result
Type: LD50
Species: guinea pig
Value: >10,000 ml/kg
Method: According to the 24 hour cuff method (Draize, et al.)
Year: 1956
GLP: no
Test substance: isobutyl isobutyrate
Remark: Contact period = 24 hours, observation period = 14 days. Based on a lack of weight gain in the animal receiving 10 ml/kg for 24 hours under an occlusive cuff, isobutyl isobutyrate can be absorbed thru the skin and cause toxicity.
Reliability: Score = 2, accepted scientific methodology
Reference: Laboratory of Industrial Medicine, Eastman Kodak Co., 1956

5.2.1 SKIN IRRITATION/CORROSION

(a) Preferred result
Species: guinea pig
Result: slight irritant
Classification:
Method: A gauze pad soaked with undiluted isobutyl isobutyrate was held in contact with the shaved skin of a guinea pig under an occlusive wrap for 24 hours. The animals were observed for 14 days after exposure.
Year: 1978
GLP: no
Test substance: isobutyl isobutyrate
Remark:
Reliability: Score = 2, accepted scientific methodology
Reference: Laboratory of Industrial Medicine, Eastman Kodak Co., 1956

5.2.2 EYE IRRITATION/CORROSION

No data available

5.3 SKIN SENSITISATION

No data available

5.4 REPEATED DOSE TOXICITY

Isobutyl isobutyrate (IBIB) is rapidly broken down to form equal amounts of isobutyric acid and isobutanol. In order to compare the toxicity information on IBIB with isobutanol, a repeated-dose oral toxicity study (also conducted by gavage) for isobutanol has been added to this section. There are no repeated-dose toxicity studies for isobutyric acid.

Isobutyl Isobutyrate SIDS Dossier

(a) Preferred result

Species: rat

Strain: Wistar

Sex: male and female

Route of Admin: oral gavage as a solution in corn oil

Exposure Period: 18 weeks

Freq. of Treatment: 7 days/week for 18 weeks

Post Exposure Observation Period: none

Doses: 0, 10, 100, and 1000 mg/kg

Control Group: yes

NOAEL: 1000 mg/kg

LOAEL: NA

Method: The study consisted of male and female animals (15/sex/group) receiving 0, 10, 100, or 1000 mg/kg isobutyl isobutyrate by oral gavage for 18 weeks. The dose volume was 5 ml/kg with the control animals receiving corn oil alone. In addition, groups of five rats of each sex were given the same doses for either 2 or 6 weeks. The animals were weighed on Day 1, 2, 6 and weekly thereafter. Feed and water consumption were determined on a cage basis (5/cage by sex). After the final dose, the animals were fasted for 24 hours and killed under barbiturate anesthesia. Hematology parameters were collected at 3, 6, and 18 weeks inot the study. A gross necropsy was performed and brain, heart, liver, spleen, kidney, adrenals, stomach, small intestine, caecum, gonads, pituitary, and thyroid(s) were weighed. Typical organs were collected, fixed in formalin, and processed for histological exam. Urine was collected during week 2, 6, and 18 of treatment and examined. Kidney function tests were also performed during week 6 and 18.

Year: 1977

GLP: no

Test substance: isobutyl isobutyrate (>98% pure).

Remark: This study was performed due to the food additive use.

Results: No deaths or abnormal behaviour was noted during the study. No differences in body weight, feed or water consumption, or histological findings were noted. The only change in haematology parameters was a decrease in hemoglobin concentration in the male 100 and 1000 mg/kg bw/day groups after two weeks of exposure. These changes were no accompanied by changes in other red blood cell parameters in these groups, and the female animals were not affected. In addition, similar findings were not observed at the 6 and 18-week haematology determinations. Kidney function tests and urine examination was normal. The only change in organ weights was an increase (14%) in relative spleen weight (when

Isobutyl Isobutyrate SIDS Dossier

corrected for body weights) in the 1000 mg/kg bw/day male group. The mean terminal body weights of animals in this group were decreased by 7% compared to the control group (not statistically significant but may explain the increased relative spleen weight). Due to the fact that there was no histological changes in this or any other organ, this finding was considered spurious.

Reliability: Score = 2, accepted scientific methodology
Reference: Drake, J. J-P., K.R. Butterworth, I.F. Gaunt, and P. Grasso. 1978. Short-Term Toxicity Study of Isobutyl Isobutyrate in Rats. *Food Cosmetics Toxicology*, 16:337-342.

(b) Species: rat
Strain: CD
Sex: male and female
Route of Admin: gavage
Exposure Period: 90 days
Freq. of Treatment: daily
Post Exposure Observation Period: N/A
Doses: 0, 100, 316, or 1000 mg/kg/day
Control Group: yes
NOAEL: 316 mg/kg
LOAEL: 1000 mg/kg
Method: Four groups of male and female rats (30/sex/group) were dosed daily by gavage with 0, 100, 316 or 1000 mg/kg/day of isobutanol for either 4 weeks (interim sacrifice; 10/sex/group) or 13 weeks (remaining animals). Dosing solutions of isobutanol in deionized water were used and 10 mL/kg was the constant dosing volume. Body weights and feed consumption were recorded weekly. Clinical signs were recorded daily. Blood and urine were collected for clinical pathology at pre-dose (10 sentinel animals), and at the 4 and 13-week necropsies. Organ weights and results of gross pathology exams were recorded at both the 4 and 13-week necropsies. Histopathological examinations of tissues from the control and 1000 mg/kg groups were conducted as well as examination of hearts, livers, and kidneys from the 100 and 316 mg/kg dose groups.

Year: 1987
GLP: yes
Test substance: isobutanol (purity 99.9%)
Remark: Analysis of dosing solutions confirmed concentrations and stability. The difference between the presence or absence of acute clinical signs of toxicity (ataxia, hypoactivity) following oral administration between the isobutanol and isobutyl isobutyrate studies is most probably due to the vehicle used. The isobutanol study

Isobutyl Isobutyrate SIDS Dossier

uses distilled water, a vehicle that would allow rapid absorption of the alcohol while the limited solubility of isobutyl isobutyrate in water necessitated the use of a corn oil vehicle. The corn oil vehicle would allow for a slower absorption of the test article and hence, the lack of acute signs of intoxication.

- Results: Treatment-related clinical signs noted in the 1000 mg/kg dose group included hypoactivity, ataxia, salivation, labored respiration, rales, prostration, hypothermia, and emaciation. Hypoactivity and ataxia were the most common clinical signs and these resolved primarily after week 4. There were no compound related clinical signs in the 100 or 316 mg/kg dose groups. The mortality rate was 1/60, 1/60, 2/60, and 11/60 for the control, 100, 316, and 1000 mg/kg groups, respectively. The only difference in body weights, body weight gain, or feed consumption was during weeks 1 and 2 of the study and were restricted to the 1000 mg/kg/day dose group. In addition, there were no dose-related differences observed in organ weights, gross pathology or histopathological examination. The mortality observed in the different dose groups was due to gavage errors, and was not due to compound administration.
- Reliability: Score = 2, standard method with restrictions; conducted with isobutanol
- Reference: "Rat Oral Subchronic Toxicity Study Final Report. Compound: Isobutyl Alcohol." Toxicity Research Laboratories, Ltd. Muskegon, MI. TRL Study #032-002 dated 1987.

5.5 GENETIC TOXICITY IN VITRO

A. BACTERIAL IN VITRO TEST

- (a) Preferred value
- Type: Ames test with additional E. coli strain
- System of Testing: Salmonella typhimurium strains TA98, TA100, TA1535, TA1537, Escherichia coli strain WP2 $uvrA^-$
- Concentration: ug/plate
- Metabolic
- Activation: with and without
- Result: Isobutyl isobutyrate was non-mutagenic when tested to a maximum concentration of 5000 micrograms/plate, using the plate incorporation protocol. At least five doses were tested in triplicate, without metabolic activation, and with 10% liver S-9 from rat. Replicate testes were performed after the initial trial to confirm these results.
- Method: Isobutyl isobutyrate was tested as a coded chemical using the plate incorporation method using S. typhimurium tester strains, TA98, TA100, TA1535, and TA1537 and E. coli strain WP2 $uvrA^-$ in the presence and absence of phenobarbitone/ β -naphthoflavone-induced liver S9 from male Sprague Dawley rats. Test concentrations of isobutyl isobutyrate (up to 5000 ug/plate) were
- Isobutyl Isobutyrate SIDS Dossier*

prepared using dimethyl sulphoxide as the solvent; a maximum of 0.1 ml solvent was added to each plate. Each dose was tested in triplicate without activation, and with 10% rat liver S-9. Concurrent positive and solvent controls were run with each trial. Repeat experiments were performed following the initial trial. A material was considered mutagenic if it produced a reproducible, dose-related increase in revertants over the solvent control, under a single metabolic activation condition, in replicate trials. A material was considered questionable if the positive response was elicited at only one concentration, or if the response could not be reproduced. A chemical was designated as non-mutagenic only after it was tested without metabolic activation, and with 10% rat S-9.

Year: 2003
GLP: Yes
Test substance: isobutyl isobutyrate
Purity: >99%
Reliability: Score = 1, guideline study

Reference: SafePharm communication to Karen Ruble, Eastman Chemical Company, December 4, 2003.

B. NON-BACTERIAL IN VITRO TEST

5.6 GENETIC TOXICITY IN VIVO

Data from isobutanol in vivo genetic toxicity studies have been included in this section. Data from isobutanol is useful when assessing the hazard associated with isobutyl isobutyrate exposure due to the rapid and complete hydrolysis of isobutyl isobutyrate to isobutanol in vivo. There are no in vivo genetic toxicity studies conducted with isobutyric acid. The toxicokinetics of the metabolism is documented and explained further in the Toxicokinetics Section (5.10.B).

(a) Preferred result

Test substance: isobutanol
Test species: Mouse/NMRI (male and female)
Test method: OECD No. 474 (Proposal for updating, ENV/EPOC (96)4)
EPA/TSCA 789.5395 (August 1997)
EEC Directive 92/69, B 12 (December 1992)

GLP: yes
Test results: Oral gavage dose of 500, 1,000 or 2,000 mg/kg of isobutanol did not have any chromosome-damaging (clastogenic) effect, and there were no indications of any impairment of chromosome distribution in the course of mitosis.

Lowest dose producing toxicity: 1000 mg/kg
Effect on Mitotic Index or P/N Ratio: None
Genotoxic effects: negative
Comments: Both of the positive control chemicals, i.e. cyclophosphamide for

clastogenicity and vincristine for spindle poison effects, led to the expected increase in the rate of polychromatic erythrocytes containing small or large micronuclei.

Reliability: Score = 1, GLP guideline study

Reference: Engelhardt, D., and Hoffmann, H.D. Cytogenetic Study In Vivo with Isobutanol in the Mouse Micronucleus Test - Single Oral Administration. (2000) Project No. 26M0243/994085, Department of Toxicology, BASF Aktiengesellschaft, D-67056 Ludwigshafen/Rhein, FRG.

5.7 CARCINOGENICITY

No data available

5.8 TOXICITY TO REPRODUCTION

Data from a two-generation reproductive toxicity study with isobutanol has been included in this section. Data from isobutanol is useful when assessing the hazard associated with isobutyl isobutyrate exposure due to the rapid and complete hydrolysis of isobutyl isobutyrate to isobutanol in vivo. There are no reproductive toxicity studies conducted with isobutyric acid. The toxicokinetics of the metabolism is documented and explained further in the Toxicokinetics Section (5.10.B).

- (a) Preferred value
- | | |
|-----------------|---|
| Type: | Two generation study |
| Species/strain: | Rat/Sprague-Dawley |
| Sex: | Male and Female |
| Route of Adm.: | inhalation |
| Method: | Conducted according to US EPA Health Effects Test Guidelines OPPTS 870.3800, Reproduction and Fertility Effects, August 1998. |

Briefly, groups of male and female rats (30/sex/group) were exposed to 0, 500, 1000, or 2500 ppm isobutanol for six hours/day, seven days/week for ten weeks prior to mating. Exposures continued in the male animals until sacrifice. The female animals were exposed thru gestation day 20, with exposure reinitiated on lactation day 5 and continued thru lactation day 28. The F1 pups were weaned on postnatal day 29 and those chosen to represent the next generation started direct inhalation exposures on postnatal day 29. These F1 male and female animals (30/sex/group) were exposed for ten weeks prior to mating. The F1 males continued exposure until sacrifice. The F1 female animals were exposed thru gestation day 20, with exposure reinitiated on lactation day 5 and continued thru lactation day 21. Body weight, feed consumption, exposure parameters, necropsy endpoints, and reproductive and developmental endpoints were collected according to the test guideline.

- | | |
|---------------------|--|
| Exposure period: | 6 hours/day |
| Freq. of treatment: | 7 days/week prior to mating, during mating and gestation; treatment was suspended during lactation days 0-4 and re-initiated on lactation day 5. |

Isobutyl Isobutyrate SIDS Dossier

Premating exposure period:	10 weeks
Exposure conc.:	0, 500, 1000 and 2500 ppm
Control group:	Concurrent
NOAEL Parental:	2500 ppm
NOAEL F1 Offspring:	2500 ppm
NOAEL F2 Offspring:	2500 ppm
Results:	Exposure to isobutanol concentrations up to 2500 ppm did not cause any parental systemic, reproductive, or neonatal toxicity when administered for two generations via whole body exposure.
GLP:	yes
Test substance:	isobutanol (>99.9% purity)
Remarks:	The highest exposure concentration was chosen based upon decreases in reaction to an external stimuli reported in a previous neurotoxicity study (Le, et al., 2001). However, the animals exposed to 2500 ppm in this study did not demonstrate decreases in response to external stimuli as was previously reported.
Reliability:	Score =1 GLP guideline study
Reference:	“An inhalation two-generation reproductive toxicity study of isobutanol in rats.” WIL Research Laboratory Study Number WIL-186013, WIL Research Laboratories, Inc., 1407 George Rd, Ashland, OH 44805-9281, sponsored by the Oxo-Process Panel of the American Chemistry Council, 1300 Wilson Boulevard, Arlington VA 22209.2003

5.9 DEVELOPMENTAL TOXICITY/TERATOGENICITY

Data from isobutanol developmental toxicity studies have been included in this section. Data from isobutanol is useful when assessing the hazard associated with isobutyl isobutyrate exposure due to the rapid and complete hydrolysis of isobutyl isobutyrate to isobutanol in vivo. There are no developmental toxicity studies conducted with isobutyric acid. The toxicokinetics of the metabolism is documented and explained further in the Toxicokinetics Section (5.10.B).

- (a) Preferred value
- | | |
|----------------------------|--|
| Species: | rat |
| Strain: | Wistar |
| Sex: | female |
| Route of Admin: | inhalation |
| Exposure Period: | Day 6 thru 15 of gestation |
| Frequency of Treatment: | Daily for 6 hours/day |
| Duration of Test: | 10 treatment days/animal |
| Exposure Conc.: | 0, 0.5, 2.5, or 10.0 mg/l |
| Control Group: | yes, concurrent no treatment |
| LOAEL (Maternal Toxicity): | 10.0 mg/l |
| NOAEL (Teratogenicity): | 10.0 mg/l |
| Method: | Pregnant rats were exposed to isobutanol by whole body inhalation from gestation day 6 thru 15. Body weights, feed consumption, and clinical sign data were collected throughout the |
- Isobutyl Isobutyrate SIDS Dossier*

	study. Chamber concentrations (actual and nominal), temperature, and absolute and relative humidity values were collected.
Year:	1990
GLP:	yes
Test substance:	isobutanol (purity >99.8%)
Result:	No treatment related effects on either the dams or the offspring were observed. Therefore, under the conditions of this study, 10 mg/l was considered a No-Observed-Effect Level for both maternal and fetal outcomes. This study corroborates the findings of Nelson, et al., <i>Tox. Ind. Health</i> , 6(314):373-387, 1990.
Reliability:	Score = 1, GLP guideline study
Reference	Klimisch, H.-J. 1990. Prenatal Toxicity of 2-Methyl-1-Propanol in Rats After Inhalation. Project No. 37R0057/88047. BASF Department of Toxicology, BASF Corp. 6700 Ludwigshafen, West Germany.
(b)	Species: rabbit
	Strain: Himalayan
	Sex: Female
	Route of Admin.: Inhalation
	Exposure
	Period: Day 7-19 of gestation
	Freq. of
	Treatment: 6 hours/day
	Duration of
	Test: Up to Day 29 post-implantation
	Exposure
	Concentrations: 0.5; 2.51; 10 mg/l
	Control Group: Yes
	NOAEL Maternal
	Toxicity: 2.51 mg/l
	NOAEL Developmental
	Toxicity: 10 mg/l
	Method: OECD Guideline 414 "Teratogenicity"
	Year: 1990
	GLP: Yes
	Test substance: Isobutanol purity >99.8%
Result:	Each control and study group contained 15 pregnant females. A slight (non-significant) retardation in body weight was observed in rabbits of the high-dose group throughout the exposure period. Otherwise, no compound-related effects indicative of maternal toxicity were found. Significantly increased incidences of intraventricular foramen/septum membranaceum (cardiac septal defects) were found for the high-dose group; this finding was not considered to be of biological significance, because by comparison with historical control data, the incidences were found to lie fully within the range of biological variation. Substance related effects on the offspring, indicative of embryo-/fetotoxicity or teratogenicity, were not observed.
Reliability:	Score = 1, GLP guideline study

Reference: BASF AG, Department of Toxicology: "Prenatal Toxicity of 2-Methyl-1-propanol in Rabbits After Inhalation", BG No.96, Project No. 90R0057/88048, 12.14.1990, conducted under the auspices of the BG Chemie, Heidelberg, (1990); Klimisch H.-J. and Hellwig J.: Fund. Appl. Toxicol., 27, 77-89, (1995).

5.10 OTHER RELEVANT INFORMATION

A. Specific toxicities

No data available

B. Toxicodynamics, toxicokinetics

(a) Preferred value

Species: rat
Strain: Sprague-Dawley
Sex: male
Route of Admin: intravenous
Exposure Period: Bolus injection into indwelling catheter
Freq. of Treatment: Single
Duration of Test: Intravenous blood sampling occurred from immediately post dosing until 240 seconds (4 minutes) after bolus injection
Exposure Conc.: 0.125 mmol/Kg
Control Group: no, used pre-injection samples as background controls
Method: Preliminary studies were conducted to select dose levels, dose formulations, and sampling times for the definitive studies. Isobutyl isobutyrate in saline with 1% Tween 20 was administered individually to seven animals via an indwelling femoral vein catheter. Serial blood samples were collected from an indwelling jugular vein catheter and immediately deproteinized to halt enzymatic activity. All catheters were made of Teflon as the test material adhered to the other types of plastic catheter materials. Extensive methods development was conducted to insure the test material was delivered into the systemic circulation and that the sampling times would provide useful metabolism/toxicokinetic data. The entire sampling period lasted only 240 seconds. Concentrations of isobutyl isobutyrate as well as down stream metabolites (isobutanol and isobutyric acid) were assayed by an internal standard GC-MS selected ion monitoring method.
Year: 2003
GLP: yes
Test substance: isobutyl isobutyrate purity > 99%
Result: Following intravenous administration of isobutyl isobutyrate, blood collection were done as fast as possible due to the rapid metabolism of the test article. Analysis of these blood samples demonstrated a extremely rapid hydrolysis of isobutyl isobutyrate to form isobutanol and isobutyric acid. The estimated $T_{1/2}$ (from a simple one-compartment model with bolus input and first-order output) was 11.1 seconds. Peak isobutyl isobutyrate levels were found in the less than 15 seconds

Isobutyl Isobutyrate SIDS Dossier

sampling time period with mean values of 1045 micromolar. The isobutyl isobutyrate levels decreased very quickly (by 46 seconds, they were at 43 micromolar) and could not be detected 166 seconds after dose administration. Isobutanol levels increased to the 78-220 micromolar range within the 15 seconds required for the first sample and stayed in this range until the end of sampling at 240 seconds. Isobutyric acid levels increased up to 304 micromolar within the 31-45 seconds time point and remained above 100 micromolar until 196 seconds. Isobutyric acids were consistently higher than the isobutanol levels suggesting formation of isobutyric acid from the further metabolism of isobutanol. This study demonstrates a extremely rapid hydrolysis of isobutyl isobutyrate, with a half-life measured in seconds. It also demonstrates the rapid appearance of the down stream metabolites, isobutanol and isobutyric acid.

Isobutyl isobutyrate, isobutanol and isobutyric acid blood levels Found following isobutyl isobutyrate intravenous injection.

Sampling Time (seconds)	Isobutyl isobutyrate*	Isobutanol	Isobutyric acid*
0	Nd**	Nd	Nd
<15	1045	103	129
16-30	405	209	268
31-45	115	218	304
46-60	43	196	295
61-75	18	163	252
76-90	12	153	225
91-105	8	135	188
106-120	6	115	161
121-135	6	128	162
136-150	3	97	126
151-165	6	103	126
166-180	Nd	98	122
181-195	Nd	87	107
196-210	Nd	78	95
211-225	Nd	82	98
226-240	Nd	96	107

*mean μM whole blood, **Nd = not detected

Reliability:
Reference:

score = 2, accepted scientific methodology
Deisinger, P.J. (2003) Unpublished data. Health and Environment Laboratories, Eastman Kodak Company, Rochester, NY. For Eastman Chemical Company, Kingsport, TN.

6.0 REFERENCES

American Public Health Association. 1971. Standard Methods for the Examination of Water and Wastewater. 1971. 13th Edi, New York, NY.

“An inhalation two-generation reproductive toxicity study of isobutanol in rats.” WIL Research Laboratory Study Number WIL-186013, WIL Research Laboratories, Inc., 1407 George Road, Ashland, OH 44805-9281, sponsored by the Oxo-Process Panel of the American Chemistry Council, 1300 Wilson Boulevard, Arlington, VA 22209.

AOPWIN. Version 1.90. Atmospheric Oxidation. EPIWIN (Estimation Program Interface for Windows). Version 3.05. Syracuse Research Corporation. Syracuse, NY.

Atkinson, R. 1987. A Structure-activity relationship for the estimation of rate constants for the gas-phase reaction of OH radicals with organic compounds. *J. Inter. Chem. Kinet.* 19:799-828.

BASF AG, Department of Toxicology: “Prenatal Toxicity of 2-Methyl-1-propanol in Rabbits After Inhalation,” BG No. 96, Project No. 90R0057/88048, 12.14.1990, conducted under the auspices of the BG Chemie, Heidelberg, (1990); Klimisch H.-J. and Hellwig J: *Fund. Appl. Toxicol.*, 27, 77-89, (1995).

BCFWIN. Version 2.14. BCF Estimate from Log Kow. EPIWIN (Estimation Program Interface for Windows). Version 3.05. Syracuse Research Corporation. Syracuse, NY.

Brook, L.T. et al., 1984. Acute Toxicities of Organic Chemicals to Fathead Minnows (*Pimephales Promelas*). Center for Lake Superior Environmental Studies. University of Wisconsin.

Christopher, S.M. November 30, 1993. “Isobutanol: Acute toxicity and irritancy testing using the rat (peroral and inhalation toxicity) and the rabbit (cutaneous and ocular tests)”. Bushy Run Research Center, Union Carbide Corp. Lab. Proj. ID 92U1166.

CRC Handbook of Chemistry and Physics. 1991-1992. D.R. Lide (ed.) 72nd ed. CRC Press, Inc. Boca Raton, FL. P.3-155.

Daubert, T.E. and R.P. Danner. 1989. Physical and Thermodynamic Properties of Pure Chemicals. Data Compilation. Hemisphere Publishing Corp. New York, NY.

Deisinger, P.J. (2003) Unpublished data. Health and Environment Laboratories, Eastman Chemical Company, Rochester, NY. For Eastman Chemical Company, Kingsport, TN.

Drake, J. J-P., K.R. Butterworth, I.F. Gaunt, and P. Grasso. 1978. Short-Term Toxicity Study of Isobutyl Isobutyrate in Rats. *Food Cosmetics Toxicology*, 16:337-342.

Eastman Chemical Company Unpublished Information, 2003.

Eastman Chemical Company, Industrial Hygiene Department. Unpublished information. February-March, 1984.

Engelhardt, D., and Hoffmann, H.D. Cytogenetic Study In Vivo with Isobutanol in the Mouse Micronucleus Test- Single Oral Administration. (2000) Project No. 26M0243/994085, Department of Toxicology, BASF Aktiengesellschaft, D-67056 Ludwigshafen/Rhein, FRG.

Isobutyl Isobutyrate SIDS Dossier

EPA's EcoSAR model (v. 0.99g). EPWIN (Estimation Program Interface for Windows). Version 3.10. U.S. Environmental Protection Agency (2000).

EPISUITE, v.3.10. United States Environmental Protection Agency, 2000.

Flick, E.W. 1991. Industrial Solvents Handbook. 4th ed. Noyes Data Corp. Park Ridge, NJ. P. 815.

Hahne R. 2002a. Acute Toxicity of the Water Accommodated Fraction (WAF) for 2 Ethylhexyl Acetate to the Rainbow Trout, *Oncorhynchus mykiss*, Determined under Static Test Conditions. Report #46955 prepared by ABC Laboratories Inc., Columbia Missouri. May 22, 2002.

Hawley's Condensed Chemical Dictionary. 1987. N.I. Sax and R.J. Lewis, Sr. (eds.). 11th ed. Van Nostrand Reinhold Co. New York, NY. P.654.

Hicks SL. 2002. Toxicity of 2-Ethylhexyl Acetate to the Unicellular Green Alga, *Selenastrum capricornutum*. Report #46957 prepared by ABC Laboratories Inc., Columbia Missouri. May 01, 2002.

HYDROWIN. Version 1.67. Aqueous Base/Acid-Catalyzed Hydrolysis. EPIWIN (Estimation Program Interface for Windows). Version 3.05. Syracuse Research Corporation. Syracuse, NY.

Industrial Hygiene and Toxicology. 1989. F.Patty (ed). 2nd ed. Volume II: Toxicology. Interscience Publishers. New York, NY. P. 1866.

Klimisch, H.-J. 1990. Prenatal Toxicity of 2-Methyl-1-propanol in Rats after Inhalation. Project No. 37R0057/88047. BASF Department of Toxicology, BASF Corp. 6700 Ludwigshafen, West Germany.

KOWIN. Version 1.66 Log Octanol-Water Partition Coefficient. EPIWIN (Estimation Program Interface for Windows). Version 3.05. Syracuse Research Corporation. Syracuse, NY.

Laboratory of Industrial Medicine, Eastman Kodak Co., 1956.

Level III Fugacity Model. EPIWIN (Estimation Program Interface for Windows). Version 3.05. Syracuse Research Corporation. Syracuse, NY.

Lyman, W.J, W.F. Reehl, and D.H. Rosenblatt. 1990. Handbook of Chemical Property Estimation Methods. American Chemical Society. Washington, D.C.

Lyman, W.J., et al. 1982. Handbook of Chemical Property Estimation Methods. Environmental Behavior of Organic Compounds. McGraw-Hill, NY.

Merck-Schuchardt. Hohenbrunn, Germany. Safety Data Sheet. Isobutyl isobutyrate for synthesis. 04.2000. The Merck Chemical Database-online. <http://www.merck.de> (accessed October 17, 2000).

Patterson KM, Hirsch MP. 1995. An Acute Aquatic Effects Test with the Daphnid *Daphnia magna*. Study #EN-431-900945-1, Eastman Kodak Company, Health and

Isobutyl Isobutyrate SIDS Dossier

Environment Laboratories, Trochester, NY. August 3, 1995.

PCKOCWIN. Version 1.66. Soil Adsorption Coefficient. EPIWIN (Estimation Program Interface for Windows). Version 3.05. Syracuse Research Corporation. Syracuse, NY.

Price, K.S., G.T. Waggy, and R.A. Conway. 1974. Brine Shrimp Bioassay and Seawater BOD of Petrochemicals. *J. Water Pollut. Contr. Fed.* 46:63-77.

“Rat Oral Subchronic Toxicity Study Final Report. Compound: Isobutyl Alcohol.” Toxicity Research Laboratories, Ltd. Muskegon, MI. TRL Study #032-002 dated 1987.

Report No. A79-146. May 25, 1979. Toxicity Data Report. Animal Industry R/D, Agricultural Division, American Cyanamid Co. P.O. Box 400, Princeton, NJ 08540.

Safe-Pharm communication to Karen Ruble, Eastman Chemical Company, December 2003.

Sigma Aldrich Product Information. <http://www.sigma-aldrich.com/saws.nsf/ProductSearch?OpenFrameet> (accessed September 25, 2000).

WSKOW. Version 1.37. Water Solubility Estimate from Log Kow. EPIWIN (Estimation Program Interface for Windows). Version 3.10. Syracuse Research Corporation. Syracuse, NY.

SIDS Initial Assessment Report**For****SIAM 20**

April 19-22, 2005, Paris, France

RECEIVED
OPPT/PAID
2006 DEC 28 AM 7:39

- 1. Chemical Name:** Isobutyl isobutyrate
- 2. CAS Number:** 97-85-8
- 3. Sponsor Country:** United States of America
National SIDS Contact Point in Sponsor Country:
Mr. Oscar Hernandez, Director
U.S. Environmental Protection Agency
Risk Assessment Division (7403 M)
1200 Pennsylvania Avenue, NW
Washington DC 20460
Phone: (202) 564-7461
E-mail: hernandez.oscar@epa.gov
- 4. Shared Partnership with:** American Chemistry Council, Oxo Process Panel
- 5. Roles/Responsibilities of the Partners:** Not applicable
- Name of industry sponsor /consortium: American Chemistry Council
Barbara Francis, Oxo Process Panel
1300 Wilson Blvd
Arlington, VA 22209
Phone: (703) 741-5609
 - Process used: Robust Summaries/dossiers, the SIAR, and the SIAP were drafted by the Oxo Process Panel's toxicologists. Documents were reviewed by the Oxo Process Panel and the United States Environmental Protection Agency.
- 6. Sponsorship History**
- How was the chemical or category brought into the SIDS Program? The American Chemistry Council's Oxo Process Panel submitted a test plan and robust summaries for this chemical to the U.S. Environmental Protection Agency in December 2001, under the International Council of Chemical Associations (ICCA) Global Initiative on High Production Volume (HPV) Chemicals Program.
- 7. Review Process Prior to** Members of the Oxo Process Panel conducted a comprehensive

- the SIAM:** literature search. Documents were prepared by the Panel and reviewed by industry toxicologists prior to submission to the United States Environmental Protection Agency (U.S. EPA). The EPA conducted reviews of submitted data and offered comments to industry. The EPA submitted documents to OECD for consideration at SIAM 18.
- 8. Quality check process:** The quality of existing data was determined using guidance provided in the Manual for Investigation of HPV Chemicals, Chapter 3: Data Evaluation (OECD, 2002).
- 9. Date of Submission:** December 22, 2003
- 10. Date of last Update:** December 20, 2004
- 11. Comments:**

CONTENTS

1	IDENTITY	5
1.1	Identification of the Substance	5
1.2	Purity/Impurities/Additives	5
1.3	Physico-Chemical properties	5
2	GENERAL INFORMATION ON EXPOSURE.....	5
2.1	Production Volumes and Use Pattern	5
2.2	Environmental Exposure and Fate.....	6
2.2.1	Sources of Environmental Exposure	6
2.2.2	Photodegradation.....	6
2.2.3	Stability in Water	6
2.2.4	Transport between Environmental Compartments	6
2.2.5	Biodegradation	7
2.2.6	Bioaccumulation.....	7
2.2.7	Other Information on Environmental Fate	8
2.3	Human Exposure	8
2.3.1	Occupational Exposure.....	8
2.3.2	Consumer Exposure	8
3	HUMAN HEALTH HAZARDS.....	8
3.1	Effects on Human Health	8
3.1.1	Toxicokinetics, Metabolism and Distribution	9
3.1.2	Acute Toxicity.....	9
3.1.3	Irritation.....	10
3.1.4	Sensitisation	10
3.1.5	Repeated Dose Toxicity	10
3.1.6	Mutagenicity.....	11
3.1.7	Carcinogenicity	11
3.1.8	Toxicity for Reproduction.....	11
3.2	Initial Assessment for Human Health.....	12
4	HAZARDS TO THE ENVIRONMENT.....	12
4.1	Aquatic Effects	12
4.2	Terrestrial Effects	14
4.3	Other Environmental Effects	14
4.4	Initial Assessment for the Environment.....	14
5	RECOMMENDATIONS	14
6	REFERENCES	16

Tables

Table 1	Summary of physico-chemical properties	5
Table 2	Aquatic toxicity data for isobutyl isobutyrate and its analogs.....	13

SIDS Initial Assessment Report

1 IDENTITY

1.1 Identification of the Substance

CAS Number: 97-85-8
 IUPAC Name: Isobutyl isobutyrate
 Molecular Formula: C₈ H₁₆ O₂
 Structural Formula: CH₃-CH(CH₃)-CH₂-O-C(=O)-CH(CH₃)-CH₃
 Molecular Weight: 144.21 g/mol
 Synonyms: Isobutyl 2-methylpropanoate
 2-Methylpropyl 2-methyl propanoate
 IBIB

1.2 Purity/Impurities/Additives

No data available.

1.3 Physico-Chemical properties

Table 1 Summary of physico-chemical properties

Property	Value
Physical state	Liquid at ambient temperature
Melting point	-80.7°C
Boiling point	148.6°C
Relative density	0.855
Vapour pressure	5.76 hPa @ 20°C
Water solubility	520 mg/L @ 27°C
Partition coefficient n-octanol/water (log value)	2.68 @ 25°C
Henry's law constant	1.58x10 ⁻³ atm·m ³ /mol @ 25°C

The references for the values found in Table 1 are in the Dossier.

2 GENERAL INFORMATION ON EXPOSURE

2.1 Production Volumes and Use Pattern

Manufacture

Isobutyl isobutyrate is produced by a single manufacturer in the United States by the catalyzed esterification of isobutyric acid with isobutyl alcohol, using a continuous enclosed process. It is

also produced as a co-product in the manufacture of Texanol® Ester Alcohol. In both cases the product is purified by distillation and stored in tanks. It is typically shipped via tank cars and tank trucks (Eastman Chemical Company unpublished information (2003)). Current U.S. annual estimated production volume is about 4.1 thousand metric tons (< 9.0 million pounds). The non-confidential U.S. production range reported for the EPA TSCA Inventory Update Rule is 3.6-5.9 thousand metric tons (8-13 million pounds) in 2002. It is sold primarily in the United States except for about 300 metric tons, which are exported to Western Europe, Japan and Latin America. The sponsors are not aware of any isobutyl isobutyrate manufacture outside of the United States.

Use

Isobutyl isobutyrate is used principally as a solvent, especially for nitrocellulose lacquers and thinners as well as coatings for plastic substrates, and high-solids coatings. The majority of end uses are in coatings (automotive, industrial, wood furniture, and graphic arts). The concentration of isobutyl isobutyrate in these formulations typically ranges from 5-15% (Eastman Chemical Company unpublished information (2003)). It is also permitted by the Federal Food, Drug and Cosmetic Act for use as a flavorant in food (21 CFR 172.515 (4/1/90)), and is used as an insect repellent. (Sax, N.I. and R.J. Lewis, Hawley's Condensed Chemical Dictionary (1987)). It has a reported use in formulations of perfumes (Kirk-Othmer Encyclopedia of Chemical Technology (1990)).

2.2 Environmental Exposure and Fate

2.2.1 Sources of Environmental Exposure

Isobutyl isobutyrate occurs naturally in fruits and essential oils. Environmental concentrations of isobutyl isobutyrate may also occur from waste streams during manufacture or through solvent use in lacquers and thinners.

2.2.2 Photodegradation

The photochemical removal of isobutyl isobutyrate from the troposphere occurs by reaction with hydroxyl radicals. This reaction is the rate-limiting step governing the overall residence time of isobutyl isobutyrate in air. Other processes, such as photolysis, wet deposition (rain-out), and dry deposition (aerosol formation) are not expected to play an important role in the atmospheric removal of isobutyl isobutyrate. Using a global average tropospheric hydroxyl radical concentration of 1.5×10^6 molecules/cm³, a second order photo-oxidation rate constant of 5.4945×10^{-12} cm³/molecule-sec, and a 12-h daylight period, the total tropospheric lifetime of isobutyl isobutyrate is expected to be about 1.947 days (Atkinson et al., 1989, EPISUITE v.3.11).

2.2.3 Stability in Water

The stability of isobutyl isobutyrate in water is pH dependent. At neutral pHs (pH 7) the half-life is about 9.217 years at 25°C. At pH 8, the hydrolysis half-life decreases to 337 days. Results were estimated using HYDROWIN v.1.67 (EPISUITE v.3.11).

2.2.4 Transport between Environmental Compartments

The values for the properties melting point, boiling point, vapor pressure, and aqueous solubility for isobutyl isobutyrate were taken from standard references (e.g., CRC Handbook, Merck Index) that contain no experimental details. Results of the use of the properties for the estimation of chemical transport between environmental compartments should be used with caution. The vapor pressure of

isobutyl isobutyrate is 5.8 hPa at 25°C and the water solubility is 520 mg/L at 27°C. A Henry's law constant was calculated to be 1.58×10^{-3} atm-m³/mol, using a molecular mass of 144.21 g/mol and the preferred vapor pressure and water solubility. For chemicals with a Henry's Law constant $< 1.0 \times 10^{-3}$ atm-m³/mole, volatilization from water is expected to be moderate. Isobutyl isobutyrate is on the borderline of being moderately volatile.

The potential for isobutyl isobutyrate to volatilize from model rivers and lakes was calculated by EPISUITE (v.3.11) using a water solubility of 520 mg/L, a vapor pressure of 5.8 hPa, and a Henry's law constant of 1.58×10^{-3} atm-m³/mol and default model assumptions. Volatilization half-lives from a model river and lake were 1.67 hours and 4.955 days, respectively. Thus, volatilization from rapidly flowing surface waters is an important removal process for isobutyl isobutyrate, while volatilization is only moderate from more quiescent lakes and other surface waters.

The preferred log K_{ow} value is 2.68. This octanol/water partition coefficient suggests that isobutyl isobutyrate would partition moderately from water to soil, sediment, or biota. Using EPISUITE (v.3.11) and PCKOCWIN (v.1.66), the soil or sediment K_{oc} for isobutyl isobutyrate was calculated to be 53.3 L/kg based on the structural features of the molecule. This soil/sediment partitioning values indicate that isobutyl isobutyrate moves moderately through soil to groundwater, with some sorption to soil expected.

Fugacity modeling (Level III) was conducted for isobutyl isobutyrate using EPISUITE (v.3.11). Input parameters included molecular weight 144.21 g/mol, melting point -80.7°C, boiling point 148.6°C, water solubility 520 mg/L, log Kow 2.68, and Henry's law constant 1.58×10^{-3} atm-m³/mol. Equal releases to air, water and soil were assumed. Media-specific half-lives were selected or calculated by the model. The model selected 5.495×10^{-12} cm³/molecule-sec as the second order rate constant for atmospheric photo-oxidation (Atkinson, 1989). Biodegradation half-lives in water, soil and sediment (360 h, 360 h, and 1440 h, respectively) were selected by the model based on the biodegradation submodels within EPISUITE (v.3.11). All other parameters used were the model default values. The results support the above conclusions regarding the movement of isobutyl isobutyrate in the environment with 12.6% distributing to air, 34.4% to water, 52.7% to soil and 0.233% to sediment.

2.2.5 Biodegradation

Biodegradation data are not available for isobutyl isobutyrate. Data from analog compounds can be used to estimate the biodegradability of isobutyl isobutyrate. Biodegradation studies with isopropyl acetate (CAS# 108-21-4) and isobutyl acetate (CAS# 110-19-0) are available. Using a standard method from the American Public Health Association (APHA, 1971), Price et al. (1974) reported 76% biodegradation of isopropyl acetate and 81% biodegradation of isobutyl acetate after 20 days, indicating that these analog compounds are readily biodegradable. These data with the analog compounds support the determination that isobutyl isobutyrate would also likely to be readily biodegradable.

2.2.6 Bioaccumulation

The bioaccumulation potential of isobutyl isobutyrate is low to moderate. The log K_{ow} value for isobutyl isobutyrate is 2.68. This octanol:water partition coefficient suggests that isobutyl isobutyrate would only accumulate in biological tissue to a moderate degree, but would not biomagnify in food chains. An estimated bioconcentration factor of 23.1 L/kg was calculated using the log Kow value of 2.68, which further suggests a low to moderate bioaccumulation potential (BCFWIN v.2.15, EPISUITE v.3.11).

2.2.7 Other Information on Environmental Fate

2.3 Human Exposure

Human exposure may occur during manufacture, formulation into products or during the use of product formulations containing isobutyl isobutyrate. Exposure also occurs through ingestion of foods that contain isobutyl isobutyrate, either naturally or as an intended food additive.

2.3.1 Occupational Exposure

No workplace exposure limits for isobutyl isobutyrate have been established. Limited exposure opportunity exists during production, which is by a single manufacturer using a closed continuous production process. The reactor, distillation column and storage tank are closed and connected via direct piping. The product is shipped primarily via tank car or tank truck, although smaller quantities may be shipped in drums. Exposure or release could occur during transport through an infrequent spill or accident. Occupational exposure during formulation of isobutyl isobutyrate solvent into products, such as lacquers and thinners is also likely to be limited, because the reactors used to mix up formulations are also enclosed. Exposure is more likely, by inhalation or through dermal contact, when products, such as lacquers or thinners, which contain isobutyl isobutyrate are used. Thirty-five personal samples were gathered on personnel working in the isobutyl isobutyrate manufacturing area. All samples were determined to be below 1 ppm (Eastman Chemical, 1984).

2.3.2 Consumer Exposure

Consumer exposure may occur during the use of product formulations that contain isobutyl isobutyrate. Such products include lacquers and thinners (Sax, N.I. and R.J. Lewis, Hawley's Condensed Chemical Dictionary (1981))(Kuney JH; Chemyclopedia(1990)). End use data shows a 55% use in automotive coatings, a 43% use in general industrial coatings, and a 2% use in graphic arts and wood furniture (Eastman Chemical Company, 2003). Isobutyl isobutyrate's use as a solvent can result in release into the atmosphere through evaporation (SRC). General population exposure can occur through inhalation of ambient air containing low concentrations of this substance. Consumers also ingest foods that contain isobutyl isobutyrate naturally or that is added as a flavorant. Isobutyl isobutyrate has been identified as a volatile constituent of strawberries, cantaloupes and muskmelons (Yamashita I et al (1976); Dirinck P et al (1984))(Yabumoto K et al (1977)). It is reported to be found in the essential oil of hops and in *vitis vinifera* (Fenaroli, (1971)). Of the 22 constituents isolated from *Eriocephalus punctulatus* oil, 50% were esters. Isobutyl isobutyrate was one of the most prevalent at 6.5% concentration (Roard D et al (1997)).

Isobutyl isobutyrate is a food additive permitted for direct addition to food for human consumption (Federal Food, Drug and Cosmetic Act and Fenaroli's Handbook of Flavor Ingredients (1971)). Synthetic flavoring substances and adjuvants, including isobutyl isobutyrate may be used in food under the Federal Food, Drug, and Cosmetic Act (Furia, T.E. Handbook of Food Additives. 2nd ed. (1968)) (21 CFR §121.1164).

3 HUMAN HEALTH HAZARDS

3.1 Effects on Human Health

Data from isobutanol toxicity studies have been included in the human health section. Data from isobutanol are useful when assessing the hazards associated with the systemic toxicity of isobutyl

isobutyrate exposure due to the rapid and complete metabolism of isobutyl isobutyrate to isobutanol and isobutyric acid *in vivo*. Isobutanol is then further metabolized to isobutyric acid. Therefore, exposure to isobutyl isobutyrate via dermal, inhalation, and water or dietary administration results in the rapid appearance of isobutanol and isobutyric acid in the systemic circulation. Since exposure to either isobutyl isobutyrate or isobutanol results in systemic exposure to isobutanol and isobutyric acid, systemic toxicity data from studies that administer isobutanol directly are useful in identifying hazards associated with isobutyl isobutyrate exposure. Data from studies conducted with isobutyric acid were not included, since there were none available. The toxicokinetics of the metabolic reaction is documented and explained below.

3.1.1 Toxicokinetics, Metabolism and Distribution

Isobutyl isobutyrate metabolism has been studied in rats.

Studies in Animals

In vivo Studies

Metabolism/toxicokinetic studies have been conducted with isobutyl isobutyrate using intravenous injections (femoral vein) with simultaneous intravenous (jugular vein) sampling (Deisinger, 2003). Isobutyl isobutyrate levels within the first seconds had mean values of 1045 μM and rapidly decreased thereafter. The calculated $T_{1/2}$ by one-compartment modeling was 11.1 seconds. Isobutanol and isobutyric acid levels increased rapidly up to peak levels of 218 and 304 μM , respectively. Isobutyric acid levels were consistently higher than isobutanol levels, suggesting further metabolism of the isobutanol metabolite to isobutyric acid. Both isobutanol and isobutyric acid levels remained increased throughout the 240 second sampling period.

Conclusion

Isobutyl isobutyrate is rapidly metabolised to isobutanol and isobutyric acid in rats.

3.1.2 Acute Toxicity

The acute toxicity values from the robust studies for all three routes of administration (oral, dermal, inhalation) are those conducted in 1956 at the Laboratory of Industrial Medicine of the Eastman Kodak Company. Although these studies were conducted prior to the promulgation of test guidelines, the design and conduct of these studies were judged acceptable by current standards.

Studies in Animals

Inhalation

Inhalation exposure to 5423 ppm (31.94 mg/L) of vapors of isobutyl isobutyrate for six hours killed 2 of 3 rats while 658 ppm (3.88 mg/L) killed 0 of 3 rats (Eastman Kodak Co., 1956). Prostration and narcosis was noted during exposure to the 5423 ppm. All of the animals that survived the 6-hour exposures gained weight at the end of the 14-day observation period. The toxicity data and clinical signs are in agreement with what has been observed with isobutanol.

Dermal

The dermal LD₅₀ values for isobutyl isobutyrate in guinea pigs was >10,000 ml/kg bw (Eastman Kodak Co., 1956).

Oral

The acute oral LD₅₀ value in rats was >6,400 mg/kg bw (Eastman Kodak Co., 1956). Weakness, ataxia, and death were noted at 12,800 mg/kg bw. The data from isobutanol acute oral toxicity studies are in agreement with the isobutyl isobutyrate data.

Conclusion

The data from the acute toxicity studies with isobutyl isobutyrate suggest that it is only slightly acutely toxic to experimental animals via the oral, dermal, and inhalation routes of exposure.

3.1.3 Irritation

Skin Irritation

Studies in Animals

Isobutyl isobutyrate was reported to cause slight skin irritation in guinea pigs (Eastman Kodak Co., 1956).

Eye Irritation

No data available

Respiratory Tract Irritation

No data available

Conclusion

Isobutyl isobutyrate is a slight skin irritant. There is no data for eye irritation.

3.1.4 Sensitisation

No data available

3.1.5 Repeated Dose Toxicity

Data from isobutanol toxicity studies have been included in this section. Data from isobutanol studies are useful for corroborating the isobutyl isobutyrate data due to the rapid and complete metabolism of isobutyl isobutyrate to isobutanol *in vivo*. Therefore, exposure to isobutyl isobutyrate via dermal, inhalation, and water or dietary administration results in the rapid appearance of isobutanol and isobutyric acid in the systemic circulation. Since exposure to either isobutyl isobutyrate or isobutanol results in systemic exposure to isobutanol and isobutyric acid, systemic toxicity data from studies that administer isobutanol directly are useful in corroborating studies conducted with isobutyl isobutyrate. The toxicokinetics of the metabolic reaction is documented and explained above. There are no repeated-dose toxicity studies conducted with isobutyric acid.

Studies in Animals

Oral

An 18-week subchronic oral gavage toxicity study has been done in rats using dose levels of 0, 10, 100, or 1,000 mg/kg bw/day (Drake, et al., 1978). The NOAEL was 1000 mg/kg bw/day in rats. Oral gavage studies with isobutanol support these findings. Acute signs of toxicity (ataxia,

hypoactivity) were noted immediately after oral exposure to 1,000 mg/kg day with isobutanol, while no clinical signs of exposure were noted following oral exposure to isobutyl isobutyrate. The difference in acute responses is related to the rate of absorption due to the vehicles used in each study. The isobutanol study used distilled water while the isobutyl isobutyrate study used a corn oil vehicle.

Conclusion

The NOAEL for oral exposure to isobutyl isobutyrate, in rats, is 1000 mg/kg bw/day.

3.1.6 Mutagenicity

Studies in Animals

In vitro Studies

Isobutyl isobutyrate was not mutagenic in a standard plate incorporation. Ames assay or an Escherichia coli WP2 uvrA gene mutation assay (SafePharm, 2003).

In vivo Studies

Data from an isobutanol toxicity study has been included in this section. Data from isobutanol is useful when assessing the hazards associated with the *in vivo* genetic toxicity of isobutyl isobutyrate exposure due to the rapid and complete hydrolysis of isobutyl isobutyrate to isobutanol and isobutyric acid *in vivo*. There are no *in vivo* genetic toxicity studies conducted with isobutyric acid.

An oral *in vivo* mouse micronucleus test conducted with isobutanol is available from BASF Corporation (Engelhardt and Hoffman, 2000). Isobutanol was administered once orally to male and female NMRI mice at doses up to 2,000 mg/kg/day body weight. Positive and negative controls all produced appropriate responses. Isobutanol did not produce any chromosome-damaging (clastogenic) effect, and there were no indications of any impairment of chromosome distribution in the course of mitosis (spindle poison effect).

Conclusion

Isobutyl isobutyrate was not genotoxic in *in vitro* experiments using bacterial cells and isobutanol was not genotoxic in *in vivo* experiments in mice.

3.1.7 Carcinogenicity

No data for carcinogenicity are available for isobutyl isobutyrate or its metabolites.

3.1.8 Toxicity for Reproduction

Studies in Animals

Data from isobutanol reproductive and developmental toxicity studies have been included in this section. Data from isobutanol are useful when assessing the hazards associated with the reproductive and developmental toxicity of isobutyl isobutyrate exposure due to the rapid and complete hydrolysis of isobutyl isobutyrate to isobutanol and isobutyric acid *in vivo*. There are no reproductive or developmental toxicity studies conducted with isobutyric acid.

Effects on Fertility

A two-generation reproductive toxicity study has been conducted by inhalation with isobutanol (WIL Research Labs, 2003). Groups of male and female rats were exposed by inhalation (6 hours/day, seven days/week) to 0, 1.52, 3.03, or 7.58 mg/L (500, 1000, or 2500-ppm) isobutanol for two generations. Daily treatments were continuous with the exception of the period between gestation day 21 through postnatal day 4 (removal of the dams from the pups during this period typically causes pup mortality). Exposure to 2500 ppm or 7.58 mg/L isobutanol did not cause any parental systemic, reproductive, or neonatal toxicity when administered for two generations via whole-body exposure.

Developmental Toxicity

In two definitive developmental toxicity studies (BASF, 1990 and Klimisch, 1990), groups of pregnant female rats (25/group) or rabbits (15/group) were exposed via inhalation to 0, 0.5, 2.5 or 10 mg/L isobutanol for 6 hours/day during gestation (rats - days 6-15; rabbits – days 7-19). Rabbit dams exposed to 10 mg/L had slight decreases in body weight gain during gestation while exposures in rats had no treatment-related effects. No evidence of developmental or fetotoxicity was reported in either the rats or the rabbits fetuses.

Conclusion

Isobutyl isobutyrate is not expected to cause reproductive or developmental toxicity based upon the available data for isobutanol.

3.2 Initial Assessment for Human Health

Isobutyl isobutyrate was only slightly toxic to experimental animals following acute oral, dermal, or inhalation exposure. Isobutyl isobutyrate is a slight skin irritant. Isobutyl isobutyrate is rapidly hydrolysed *in vivo* to isobutanol and isobutyric acid. Data from isobutanol acute oral toxicity and repeated-dose toxicity studies have been used to corroborate conclusions from studies conducted with isobutyl isobutyrate. In addition, *in vivo* mutagenicity, and reproductive and developmental toxicity studies has been used to identify hazards associated with isobutyl isobutyrate exposure. Repeated dose toxicity studies for isobutyl isobutyrate using the oral route of exposure had NOAELs of 1000 mg/kg bw/day. The NOAEL for reproductive and developmental toxicity from a two-generation reproductive toxicity study and developmental toxicity studies with isobutanol in rats was approximately 2500 ppm 7.58 mg/L for the two-generation study and 10 mg/L for the developmental toxicity studies. An *in vitro* mutagenicity test with isobutyl isobutyrate was negative and an *in vivo* micronucleus test with isobutanol was negative for genotoxicity.

4 HAZARDS TO THE ENVIRONMENT

4.1 Aquatic Effects

Analog Justification

The only acute toxicity data available for isobutyl isobutyrate is with the daphnid, *Daphnia magna*. Based on general structural similarities as acetates, data for 2-ethylhexyl acetate (CAS# 103-09-3) are used to address the aquatic toxicity for isobutyl isobutyrate.

Acute Toxicity Test Results

Fish

No data on fish toxicity are available for isobutyl isobutyrate. Data are available for the analog compound 2-ethylhexyl acetate. Available data as well as ECOSAR predicted toxicity values are shown in Table 2 below. Two studies with rainbow trout (*Oncorhynchus mykiss*) are available for 2-ethylhexyl acetate. Acute 96-hour LC₅₀s of 8.27 and >4.2 mg/L were reported.

ECOSAR values for isobutyl isobutyrate and 2-ethylhexyl acetate are included for comparison to the available measured data. For isobutyl isobutyrate, an ECOSAR value of 9.455 mg/L for fish was calculated. An ECOSAR value of 3.057 mg/L was calculated for 2-ethylhexyl acetate.

Invertebrates

Data are available isobutyl isobutyrate and the invertebrate *Daphnia magna* with reported 48-hour EC₅₀ values of 55.8 to 59.3 mg/L reported. For comparison purposes, a 48-h EC₅₀ with *Daphnia magna* of 22.9 mg/L 2-ethylhexyl acetate is also presented. ECOSAR values for isobutyl isobutyrate and 2-ethylhexyl acetate were 27.566 and 3.571 mg/L, respectively.

Green Algae

No acute toxicity data are available for green algae with isobutyl isobutyrate. Data are available with 2-ethylhexyl acetate and the green alga *Selenastrum capricornutum* with a 72-hour EC₅₀ value of >21.9 mg/L reported and 72-h NOEC (LOEC) values of 10.3 (21.9) mg/L.

ECOSAR values for isobutyl isobutyrate and 2-ethylhexyl acetate were calculated to be 0.771 mg/L and 0.26 mg/L respectively. It should be noted that the ECOSAR value for algae for esters is generally not very predictive of measured values because ECOSAR is overly conservative in its predictions for green algae.

Aquatic Toxicity Summary

The data for isobutyl isobutyrate, the structural analog 2-ethylhexyl acetate and their ECOSAR values allow the estimation of the acute aquatic toxicity of isobutyl isobutyrate. All data are presented in Table 2.

Table 2 Aquatic toxicity data for isobutyl isobutyrate and its analogs.

	Isobutyl isobutyrate	2-Ethylhexyl acetate
	CAS# 97-85-8	CAS# 103-09-3
	C ₈ H ₁₆ O ₂	C ₁₀ H ₂₀ O ₂
	MW=144.22	MW=172.27
Fish	No data available	8.27 mg/L >4.2 mg/L
Fish (ECOSAR)	9.455 mg/L	3.057 mg/L
Daphnids	55.8 to 59.3 mg/L	22.9 mg/L
Daphnid (ECOSAR)	27.566 mg/L	3.571
Green algae	No data available	>21.9 mg/L NOEC (LOEC)= 10.3 (21.9) mg/L
Green algae (ECOSAR)	0.771 mg/L	0.260 mg/L

Chronic Toxicity Test Results

No data available.

4.2 Terrestrial Effects

No data available.

4.3 Other Environmental Effects

4.4 Initial Assessment for the Environment

The available physicochemical data are adequate to describe the properties of isobutyl isobutyrate. Isobutyl isobutyrate has a melting point of $-80\text{ }^{\circ}\text{C}$, boiling point of $148.6\text{ }^{\circ}\text{C}$ and has a vapor pressure of 5.8 hPa at 25°C , a water solubility of 520 mg/L at 27°C and a log K_{ow} of 2.68. The photochemical removal of isobutyl isobutyrate as mediated by hydroxyl radicals occurs with a calculated half-life of 1.947 days. Isobutyl isobutyrate is likely to be readily biodegradable under aerobic conditions, based on data for isopropyl- and isobutyl-acetate. Isobutyl isobutyrate volatilises easily from moving rivers, but volatilises only moderately from quiescent lakes and other surface water bodies (calculated volatilization half-lives of 1.67 hours from a river and 4.955 days from a lake). Isobutyl isobutyrate is not persistent in the environment and is not likely to bioaccumulate in food webs. Based on Level III distribution modelling it is estimated that the majority of isobutyl isobutyrate released to the environment will partition into water (34.4%) and soil (52.7%), with a smaller amount in air (12.6%). The stability of isobutyl isobutyrate in water is pH dependent, at neutral pHs (7) the $T_{1/2} = 9.2$ years at 25°C and at higher pHs (8) the $T_{1/2}$ is shortened to 337 days.

Except for a study with the aquatic invertebrate *Daphnia magna*, aquatic toxicity data are not available for isobutyl isobutyrate. Data for the structurally similar 2-ethylhexyl acetate (CAS# 103-09-3) were used to supplement the data for isobutyl isobutyrate. For fish, two studies with rainbow trout (*Oncorhynchus mykiss*) and 2-ethylhexyl acetate are available. Acute 96-hour LC_{50s} of 8.27 and >4.2 mg/L were reported for 2-ethylhexyl acetate. Data are available with isobutyl isobutyrate and the invertebrate *Daphnia magna* with 48-hour EC_{50} values of 55.8 to 59.3 mg/L reported. In addition, a daphnid study with 2 ethylhexyl acetate reported a 48-hour EC_{50} of 22.9 mg/L. Data are available with 2-ethylhexyl acetate and the green alga *Selenastrum capricornutum* with a 72-hour EC_{50} value of >21.9 mg/L and a 72-h EC_3 of 21.9 mg/L reported. ECOSAR values for 2-ethylhexyl acetate were calculated to be 3.057 mg/L for fish, 3.571 mg/L for daphnids, and 0.260 mg/L for green algae. The measured data with the analog compound and the ECOSAR values for isobutyl isobutyrate and its analog compound allow the estimation of the acute aquatic toxicity of isobutyl isobutyrate. Terrestrial data are not available.

5 RECOMMENDATIONS

The chemical is currently of low priority for further work.

This conclusion is supported by the fact that adequate SIDS level physicochemical and toxicological data are available to characterize isobutyl isobutyrate. Isobutyl isobutyrate is not persistent in the environment and is not expected to bioaccumulate. Although environmental monitoring data are not found for isobutyl isobutyrate, fugacity-dependent modeling indicates that the majority of isobutyl isobutyrate released to the environment will partition into water (34.4%) and soil (52.7%), with a smaller amount in air (12.6%). The amounts of isobutyl isobutyrate in

other compartments (e.g., sediment) are expected to be negligible. Since isobutyl isobutyrate has a fairly short half-life in the environment, these theoretical levels are not expected to persist or bioconcentrate. Terrestrial data are not available.

6 REFERENCES

American Public Health Association. 1971. Standard Methods for the Examination of Water and Wastewater. 1971. 13th Edi, New York, NY.

“An inhalation two-generation reproductive toxicity study of isobutanol in rats.” WIL Research Laboratory Study Number WIL-186013, WIL Research Laboratories, Inc., 1407 George Road, Ashland, OH 44805-9281, sponsored by the Oxo-Process Panel of the American Chemistry Council, 1300 Wilson Boulevard, Arlington, VA 22209.

Atkinson R. 1987. A Structure-activity relationship for the estimation of rate constants for the gas-phase reaction of OH radicals with organic compounds. *J. Inter. Chem. Kinet.* 19:799-828.

BASF AG, Department of Toxicology: “Prenatal Toxicity of 2-Methyl-1-Propanol in Rabbits After Inhalation,” BG No. 96, Project No. 90R0057/88048, 12.14.1990, conducted under the auspices of the BG Chemie, Heidelberg, 1990.

Budvari, S. editor. Merck Index-An Encyclopedia of Chemicals, Drugs, and Biologicals. 12th edition. Merck and Co., Inc., Rahway, NJ, 1996.

CRC Handbook of Chemistry and Physics. 1991-1992. D.R. Lide (ed.) 72nd ed. CRC Press, Inc. Boca Raton, FL. P.3-155.

Deisinger, P.J.(2003). Unpublished data. Health and Environment Laboratories, Eastman Chemical Company, Rochester, NY, For Eastman Chemical Company, Kingsport, TN.

Dirinck P et al; Anal Volatiles: Methods Appl, Proc – Workshop 1983: 381-400 (1984).

Drake, J. J-P., K.R. Butterworth, I.F. Gaunt, and P. Grasso. 1978. Short-Term Toxicity Study of Isobutyl Isobutyrate in Rats. *Food Cosmetics Toxicology*, 16:337-342.

Eastman Chemical Company unpublished information (2003).

Eastman Chemical Company, Industrial Hygiene Department. Unpublished information. February-March, 1984.

Engelhardt, D., and Hoffman, H.D. Cytogenetic Study In Vivo with Isobutanol in the Mouse Micronucleus Test- Single Oral Administration. (2000) Project No. 26M0243/994085, Department of Toxicology, BASF Aktiengesellschaft, D-67056 Ludwigshafen/Rhein, FRG.

EPA's EcoSAR model (v. 0.99g). EPWIN (Estimation Program Interface for Windows). Version 3.10. U.S. Environmental Protection Agency (2000).

EPIWIN (Estimation Program Interface for Windows) v.3.10. Syracuse Research Corporation. Syracuse, NY.

Federal Food, Drug and Cosmetic Act, 21 CFR 172.515 (4/1/90) and §121.1164.

Fenaroli's Handbook of Flavor Ingredients. Volume 2. Edited, translated, and reviewed by T.E. Furia and N. Bellance. 1st ed. Cleveland: The Chemical Rubber Co., (1971) 461.

Furia, T.E. (ed.). CRC Handbook of Food Additives. 2nd ed. Volume 2. Boca Raton, Florida: CRC Press, Inc., (1968) 729, 873

HYDROWIN v.1.67. Aqueous Base/Acid-Catalyzed Hydrolysis. EPIWIN (Estimation Program Interface for Windows). Version 3.05. Syracuse Research Corporation. Syracuse, NY.

Kirk-Othmer Encyclopedia of Chemical Technology. 3rd ed., Volumes 1-26. New York, NY: John Wiley and Sons, 1978-84, p. V9 334 (1990).

Klimisch, H.J. 1990. Prenatal Toxicity of 2-Methyl-1-propanol in Rats after Inhalation. Project No. 37R0057/88047. BASF Department of Toxicology, BASF Corp. 6700 Ludwigshafen, West Germany.

Kuney JH; Chemyclopedia 1991 Washington DC: Amer Chem Soc p. 83 (1990).

Laboratory of Industrial Medicine, Eastman Kodak Co., 1956.

Level III Fugacity Model. EPIWIN (Estimation Program Interface for Windows) Version 3.05. Syracuse Research Corporation, Syracuse, NY.

PCKOWCIN. V.1.66. Soil Adsorption Coefficient. EPIWIN (Estimation Program Interface for Windows). Version 3.05. Syracuse Research Corporation. Syracuse, NY.

Price, K.S., G.T. Waggy, and R.A. Conway. 1974. Brine Shrimp Bioassay and Seawater BOD of Petrochemicals. J. Water Pollut. Contr. Fed. 46:63-77.

Road D et al Parfums Cosmet Aromes 14: 57 (1997).

Safe-Pharm communication to Karen Ruble, Eastman Chemical Company, December 2003.

Sax, N.I. and R.J. Lewis Sr. (eds). Hawley's Condensed Chemical Dictionary, 11th ed.1987. Van Nostrand Reinhold Co. New York, NY. P654.

Sax, N.I. and R.J. Lewis, Sr. (eds.),Hawley's Condensed Chemical Dictionary 10th ed. NY: Van Nostrand Reinhold p. 576 (1981).

Yabumoto K et al; J Food Sci 42: 32-7 (1977).

Yamashita I et al; Phytochemistry 15: 1633-7 (1976).