

Part II

Reviews of Representative Stream Assessment and Mitigation Protocols

ABBREVIATIONS

| | |
|-------|--|
| Dbkf | Bankfull depth |
| DO | Dissolved oxygen |
| DOC | Dissolved organic carbon |
| EMAP | Environmental Monitoring and Assessment Program |
| est | Estimate |
| GIS | Geographic Information System |
| IBI | Index of Biotic Integrity |
| max | Maximum |
| min | Minimum |
| QA/QC | Quality Assurance / Quality Control |
| O/E | Observed:Expected ratio |
| opt | Optional |
| RBP | Rapid Bioassessment Protocols (Barbour et al., 1999) |
| REMAP | Regional Environmental Monitoring and Assessment Program |
| TSS | Total dissolved solids |
| USACE | U.S. Army Corps of Engineers |
| USDA | U.S. Department of Agriculture |
| USEPA | U.S. Environmental Protection Agency |
| USFS | U.S. Forest Service |
| USGS | U.S. Geologic Survey |
| Wbkf | Bankfull width |
| Wfpa | Flood prone width |

LIST OF PROTOCOL REVIEWS

| | | |
|-----|---|--------|
| 1. | Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers, U.S. Environmental Protection Agency | II - 1 |
| 2. | Revised Methods for Characterizing Stream Habitat in the National Water Quality Assessment Program, U.S. Geologic Survey | II - 3 |
| 3. | Field Operations Manual for Assessing the Hydrologic Permanence and Ecological Condition of Headwater Streams, U.S. Environmental Protection Agency | II - 5 |
| 4. | Environmental Monitoring and Assessment Program (EMAP), Physical Habitat Characterization, U.S. Environmental Protection Agency | II - 7 |
| 5. | Methods for Evaluating Stream, Riparian, and Biotic Conditions, U.S. Forest Service | II -10 |
| 6. | Wadeable Streams Assessment: Field Operations Manual, U.S. Environmental Protection Agency | II -12 |
| 7. | Watershed Assessment of River Stability & Sediment Supply (WARSSS), Rosgen/U.S. Environmental Protection Agency | II -16 |
| 8. | Stream Geomorphic Assessment Protocol Handbooks, Vermont Agency of Natural Resources | II -18 |
| 9. | A Physical Habitat Index for Freshwater Wadeable Streams in Maryland, Maryland Department of Natural Resources..... | II -21 |
| 10. | Physical Habitat and Water Chemistry Assessment Protocol for Wadeable Streams Monitoring, Minnesota Pollution Control Agency..... | II -24 |
| 11. | Field evaluation manual for Ohio's primary headwater habitat streams, Ohio Environmental Protection Agency | II -26 |
| 12. | The Qualitative Habitat Evaluation Index (QHEI), Ohio Environmental Protection Agency | II -29 |
| 13. | Guidelines for Evaluating Fish Habitat in Wisconsin Streams, U.S. Forest Service | II -31 |
| 14. | Physical Habitat of Aquatic Ecosystems, Texas Commission on Environmental Quality | II -33 |
| 15. | Subjective Evaluation of Aquatic Habitats, Kansas Department of Wildlife & Parks | II -35 |

| | | |
|-----|---|---------|
| 16. | Effectiveness monitoring for streams and riparian areas: sampling protocol for stream channel attributes, AREMP & PACFISH/INFISH (PIBO)..... | II - 37 |
| 17. | R1/R4 (Northern and Intermountain Regions) Fish and Fish Habitat Standard Inventory Procedures Handbook, U.S. Forest Service..... | II - 39 |
| 18. | Effectiveness monitoring for streams and riparian areas within the Pacific Northwest: stream channel methods for core attributes, AREMP & PACFISH/INFISH (PIBO) | II - 41 |
| 19. | A Manual of Procedures for Sampling Surface Waters, Arizona Department of Environmental Quality | II - 43 |
| 20. | Stream Condition Inventory (SCI) Technical Guide, U.S. Forest Service..... | II - 46 |
| 21. | Idaho Small Stream Ecological Assessment Framework, Idaho Department of Environmental Quality | II - 48 |
| 22. | Idaho River Ecological Assessment Framework, Idaho Department of Environmental Quality | II - 51 |
| 23. | Beneficial Use Reconnaissance Program Field Manual for Streams, Idaho Department of Environmental Quality | II - 54 |
| 24. | Methods for Stream Habitat Surveys, Oregon Department of Fish and Wildlife..... | II - 56 |
| 25. | Stream Inventory Handbook: Levels I & II, U.S. Forest Service | II - 58 |
| 26. | Functional Assessment Approach for High Gradient Streams: West Virginia, U.S. Army Corps of Engineers, Huntington District | II - 60 |
| 27. | West Virginia Stream and Wetland Valuation Metric, West Virginia Interagency Review Team | II - 62 |
| 28. | [Virginia] Unified Stream Methodology, U.S. Army Corps of Engineers, Norfolk District and Virginia Department of Environmental Quality..... | II - 65 |
| 29. | Standard Operating Procedure: Compensatory Mitigation, U.S. Army Corps of Engineers, Charleston District..... | II - 67 |
| 30. | [Kentucky] Draft Stream Relocation/Mitigation Guidelines, Kentucky Division of Water..... | II - 69 |
| 31. | Stream Assessment Protocol for Headwater Streams in the Eastern Kentucky Coalfield Region, U.S. Army Corps of Engineers, Louisville District..... | II - 71 |
| 32. | Stream Mitigation Guidelines [NC], U.S. Army Corps of Engineers, Wilmington District | II - 73 |

| | | |
|---------------------------------|--|----------------------|
| Name | Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers | Catalog No. 1 |
| Primary Author/ Agency | USEPA Barbour, M.T., J. Gerritsen, B.D. Snyder, and J.B. Stribling. 1999. Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition. EPA 841-B-99-002. USEPA Office of Water, Washington, D.C. | |
| Electronic Resource | http://www.epa.gov/owow/monitoring/rbp/ | |
| Intended Use/Purpose | Non-Regulatory Condition Assessment; Inventory; Ambient Monitoring. | |
| Target Resource Type | Wadeable Streams | |
| Scale/Unit of Assessment | Stream reach, 100 meters | |
| Geographic Applicability | Nationwide | |
| General Level of Effort | Varies based on the specific components of the protocol that are employed: Easy (rapid), Moderate, or Intensive (1 day± in the field by a trained or experienced crew of 2 or more persons). | |
| Assessment Parameters | <p><u>Habitat Assessment Index (based on visual observation)</u></p> <p>Channel/Valley Morphology: Channel alteration (H, L)¹; frequency of riffles or bends (H); sinuosity (L); pool substrate characterization (L); Velocity/depth combinations (H); pool variability (L); bank stability (H, L).</p> <p>Physical Habitat: Epifaunal substrate/available cover (H, L); embeddedness (H); sediment deposition (H, L); channel flow status (H, L); bank vegetative protection (H, L); riparian zone width (H, L).</p> <p>Water Quality: --</p> <p>Biology: --</p> <p>Other: --</p> <p>¹ H = applicable in high gradient streams; L = applicable in low gradient streams.</p> <p><u>Additional Assessment Parameters</u></p> <p>Channel/Valley Morphology: --</p> <p>Physical Habitat: Stream velocity; stream depth; canopy cover class; woody debris tally; substrate particle size classes (est.); predominant riparian vegetation type; dominant aquatic vegetation type and species.</p> <p>Water Quality: Temperature, specific conductivity; dissolved oxygen; pH; turbidity; water odors (classes); surface oils (classes); sediment odors (classes).</p> <p>Biology: Periphyton (quantitative protocols for single habitat and multi-habitat provided and field-based rapid periphyton survey protocol described); benthic macroinvertebrates (single habitat and multi-habitat protocols provided); fish.</p> <p>Other: Predominant surrounding land use.</p> | |
| Resolution | Qualitative (descriptive); Semi-Quantitative (ordinal scale, rank, etc.) ~ mostly applicable to physical habitat assessment; Quantitative (actual measurement or estimate) ~ mostly applicable to biological assessment(s). | |

| | | |
|---|---|----------------------|
| Name | Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers | Catalog No. 1 |
| Output | Condition Assessment ~ once data analyses and regional relationships have been developed. Index (e.g. numeric score) ~ physical habitat; Raw data ~ biological data. | |
| Reference | Barbour et al. (1999) stress that regional reference conditions should be used to scale the assessment to the 'best attainable conditions' for synoptic surveys or those for monitoring trends over time. However, the authors also state that site-specific reference conditions may be better suited to assess specific sources of stream impact. | |
| QA/QC | The RBP stresses that practitioners should be trained in the assessment procedure and work in teams in order to minimize observer bias. Specific QA/QC measures for both field sampling and laboratory analysis (if applicable) are provided for each main chapter in the RBP manual (e.g. benthic macroinvertebrates, fish, etc). | |
| Description/ Summary | <p>The primary purpose of the Rapid Bioassessment Protocols for Use in Wadeable Streams and Rivers (RBP) is "to describe a practical technical reference for conducting cost-effective biological assessments of lotic ecosystems," (Barbour et al., 1999). The author advocate integrated assessments of stream condition that incorporate physical habitat, water quality, and biological measures, such as periphyton, benthic macroinvertebrates, and fish.</p> <p>The RBP stream habitat assessment is a visual-based rapid assessment that relies upon visual characterizations of ten stream features in order to categorize the quality of those features as either poor, marginal, suboptimal, or optimal. The range of quality from poor to optimal is further defined on a point scale from 0 to 20 for each stream habitat parameter assessed. Thus, the maximum point score for the RBP habitat assessment is 200. Quality descriptions are outlined on the field data sheets and further described and illustrated in the text of the RBP manual itself. There are a few different or modified stream habitat parameters used in the assessment based on whether the stream has a high gradient and therefore dominated by riffle/run habitat types and coarse substrate, or a low gradient dominated by glide/pool habitats and typically finer substrates.</p> <p>Barbour et al. (1999) also outline biological data analysis techniques, discuss the integration of physical habitat data and biological data, and suggest methods of reporting and graphically summarizing RBP data. Numerous data forms are provided, and examples of concepts and ideas are illustrated with real data from around the country. Step by step field procedures are suggested and equipment lists provided.</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Barbour et al. (1999) describe the general RBP habitat assessment, as reviewed herein, as a Level I approach that takes approximately 15-20 minutes in the field. However, the authors also suggest that more quantitative and less ambiguous measures of stream habitat parameters, such as USEPA EMAP methods (Kaufmann and Robison, 1997), result in considerably greater precision. | |
| Seasonality | Periphyton: Late summer or early fall; Benthic Macroinvertebrates: Depends on program objectives. Fish: Mid to late summer. Physical Habitat: Not stated. | |
| Related Procedures/ References | Kaufmann, P.R., and E.G. Robison. 1998. Physical Habitat Characterization, Section 7 in J.M. Lazorchak et al. (eds). EMAP- Surface Waters: Field Operations and Methods for Measuring the Ecological Condition of Wadeable Streams. EPA/620/R-94/004F, USEPA, Washington, D.C. | |
| Other/Notes | <p>The RBP has become a defining framework for biological assessment programs in many U.S. States. The RBP Habitat Assessment Index in particular is an especially common component of other local or regional stream assessment protocols.</p> <p>Barbour et al (1999) stress that implementation of the RBP is enhanced by developing empirical relationships between habitat quality and biological conditions within specific geographic regions.</p> | |

| | | | |
|---------------------------------|--|---|----------|
| Name | Revised Methods for Characterizing Stream Habitat in the National Water Quality Assessment Program | Catalog No. | 2 |
| Primary Author/ Agency | U.S. Geologic Survey Fitzpatrick, F.A., I.R. Waite, P.J. D'Arconte, M.R. Meador, M.A. Maupin, and M.E. Gurtz. 1998. Revised Methods for Characterizing Stream Habitat in the National Water Quality Assessment Program. U.S. Geologic Survey, WRI Report 98-4052, Raleigh, NC. 67 pp. | | |
| Electronic Resource | http://pubs.usgs.gov/wri/wri984052/ | | |
| Intended Use/Purpose | Inventory; Ambient Monitoring. | | |
| Target Resource Type | Wadeable and non-wadeable streams. | | |
| Scale/Unit of Assessment | Stream reach, 20X the mean wetted channel width (Wadeable streams: minimum 150 meters, maximum 300 meters; Non-wadeable streams: minimum 500 meters, maximum 1,000 meters). Fitzpatrick et al. (1998) also present procedures for collecting and analyzing data at basin and channel segment scales via GIS, topographic mapping, and aerial photography. | | |
| Geographic Applicability | Nationwide. | | |
| General Level of Effort | Moderate to Intensive. | | |
| Assessment Parameters | Channel/Valley Morphology: (optional) | Stream discharge; water surface gradient; water depth; flow velocity; wetted channel width; channel habitat units [bed forms]; sinuosity; channel gradient; bankfull stage; bank angle; bank height; bank stability index (based on bank angle, bank vegetative cover, bank height, & dominant bank substrate); cross-sectional channel dimensions; substrate particle size analysis (est. required; pebble counts, optional). | |
| | Physical Habitat: | In-stream cover (type and percent-cover); bank vegetative cover; embeddedness; riparian vegetative cover (densiometer). | |
| | Water Quality: | -- | |
| | Biology: | Riparian vegetation stem density, basal area, & speciation (via point-centered quarter method, optional). | |
| | Other: | Stream order; watershed area; cumulative perennial stream length; drainage density; basin length; drainage shape (ratio of drainage area and the square of the basin length); basin relief; basin relief ratio (ratio of basin relief and basin length); entire stream gradient (ratio of difference between elevation at 85% and 10% of stream length and the stream length between these two points); dominant riparian land use. | |
| Resolution | Qualitative (descriptive); Semi-Quantitative (ordinal scale, rank, etc.); and Quantitative (actual measurement or estimate). | | |
| Output | Raw data | | |
| Reference | N/A (The objective of the method or procedure is not presented in the context of defining the condition of a resource. However, it may be used to identify or establish reference conditions.) | | |

| | | |
|---|--|-----------------------------|
| Name | Revised Methods for Characterizing Stream Habitat in the National Water Quality Assessment Program | Catalog No. 2 |
| QA/QC | Not stated. | |
| Description/ Summary | <p>The goal of the National Water Quality Assessment (NAWQA) Program is to assess status and trends in water quality nationwide and to develop an understanding of the major factors influencing observed conditions and trends. Stream habitat assessments are conducted as part of the NAWQA Program in order measure habitat characteristics essential in describing and interpreting water chemistry and biological conditions (Fitzpatrick et al., 1998). These procedures allow for appropriate habitat descriptions and standardization of measurement techniques to facilitate unbiased evaluations of habitat influences on stream conditions at local, regional, and national scales (Fitzpatrick et al., 1998).</p> <p>The Revised Methods for NAWQA stream habitat characterizations integrate data at four spatial scales: 1) basin (watershed); 2) segment; 3) reach; and 4) microhabitat. Basin and segment-scale assessments are undertaken using GIS, topographic maps, aerial photographs, etc. A stream segment is defined in the NAWQA program as "a length of stream that is relatively homogeneous with respect to physical, chemical, and biological properties," and may be over several kilometers long (Fitzpatrick et al., 1998). Watershed size, climate and potential runoff characteristics, and land use are determined at the basin-scale, while stream gradient, sinuosity, and water management features are measured at the segment-scale. A computer program called "Basinsoft" has been developed by USGS to quantify a number of basin characteristics using GIS (Harvey and Eash, 1996). The stream reach scale is most commonly at issue for restoration and mitigation projects, and the remainder of this summary will focus primarily on stream reach scale aspects of the NAWQA Revised Methods.</p> <p>Reach-scale data is collected in the field from 11 systematically placed, equally-spaced transects (channel cross-sections); the spacing of which is based on stream width. The Revised Methods includes quantitative, semi-quantitative, and qualitative metrics. Specific methods for measuring or estimating reach-scale data are provided, and numerous illustrative graphs and figures are used to clarify concepts and instructions. There are additional sampling procedures for optional parameters, as noted in the Assessment Parameters section above.</p> <p>Data forms are provided for recording basin, segment, and reach scale data, although it is acknowledged that some may need revision to meet local needs. The Revised Methods manual also includes a suggested data management hierarchy that is available on the internet, which can be imported into a variety of commercial spreadsheet and database software applications. Data analysis is described, and specific statistical procedures that can be utilized to identify relationships among habitat variables and/or relationships among habitat variables and biological components of the stream system are recommended.</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | Not stated. | |
| Related Procedures/ References | Harvey, C.A. and D.A. Eash. 1996. Description, instructions, and verification for Basinsoft, a computer program to quantify drainage-basin characteristics, U.S. Geologic Survey Water Resources Investigations Report 95-4287. 25 pp. | |
| Other/Notes | | |

| | | |
|---------------------------------|--|----------------------|
| Name | Field Operations Manual for Assessing the Hydrologic Permanence and Ecological Condition of Headwater Streams | Catalog No. 3 |
| Primary Author/ Agency | U.S. EPA Fritz, K.M., B.R. Johnson, and D.M. Walters. 2006. Field Operations Manual for Assessing the Hydrologic Permanence and Ecological Condition of Headwater Streams. EPA/600/R-06/126. U.S. Environmental Protection Agency, office of Research and Development, Washington, D.C. | |
| Electronic Resource | http://www.epa.gov/eerd/manual/headwater.htm | |
| Intended Use/Purpose | Inventory; Ambient Monitoring. | |
| Target Resource Type | Headwater streams (ephemeral, intermittent, and perennial) with a drainage area less than 1 square mile. | |
| Scale/Unit of Assessment | Stream reach, 40x the channel width (~30 meters), absent of any tributary confluence | |
| Geographic Applicability | Forested, temperate regions (study sites were located in Indiana, Illinois, Kentucky, Ohio, New Hampshire, New York, Vermont, Washington, and West Virginia). | |
| General Level of Effort | Intensive | |
| Assessment Parameters | <p>Channel/Valley Morphology: Stream discharge; water depth; flow velocity; wetted channel width; channel gradient; categorical channel habitat units (erosional habitats vs. depositional habitats); sinuosity (no. of complete meanders in sample reach); bankfull width; bankfull depth; floodprone area width; depth to bedrock; depth to groundwater; streambed sediment moisture content; substrate particle size classes.</p> <p>Physical Habitat: Riparian vegetative cover (densiometer).</p> <p>Water Quality: Temperature; conductivity; pH; dissolved oxygen.</p> <p>Biology: Bryophytes (qualitative or quantitative); algae (qualitative or quantitative); benthic invertebrates (quantitative); amphibians (semi-quantitative).</p> <p>Other: - -</p> | |
| Resolution | Qualitative (descriptive; categorical), Semi-Quantitative (ordinal scale, rank, etc.), and Quantitative (actual measurement or estimates). | |
| Output | Raw data | |
| Reference | N/A (The objective of the method or procedure is not presented in the context of defining the condition of a resource. However, it may be used to identify or establish reference conditions.) | |
| QA/QC | Not stated. | |

| | | |
|---|--|----------------------|
| Name | Field Operations Manual for Assessing the Hydrologic Permanence and Ecological Condition of Headwater Streams | Catalog No. 3 |
| Description/ Summary | <p>The “Field Operations Manual for Assessing the Hydrologic Permanence and Ecological Condition of Headwater Streams” provides a compilation of methods useful to characterize headwater streams. The Manual does not present information allowing the user to immediately assess the <i>condition</i> of any given headwater stream (i.e. there is no reference condition or index provided for any particular geographic region). Instead, the Manual provides an assemblage of recommended methods and/or tools potentially useful to undertake an exercise aimed at developing a regional reference database. It does however include a section outlining considerations for field sampling design, including minimum sample size, hypothesis testing, and even a brief introduction to BACI study designs (before/after control/impact). The Manual also provides conceptual backgrounds explaining the purpose and relevance of each suggested parameter.</p> <p>Study sites used for testing the methods included in the Manual were limited to basin areas consistent with the “Primary Headwater Habitat Streams” protocol of the Ohio Environmental Protection Agency (OEPA, 2002), and the methods for some parameters included in the Manual are adapted from OEPA (2002).</p> <p>Instructions for each step are well defined, including photographs and/or diagrams. Materials lists and literature references for each step of each method are included following each section of the report. Recommended field data sheets are provided.</p> <p>Alternative sampling methods are provided for documenting many stream parameters based on the type of equipment available (e.g. stream discharge; flow velocity; channel slope; etc.).</p> | |
| Expertise Required | Not stated. However, proposed sampling, sorting, data reduction, and analysis of biological community assemblages should only be undertaken by persons with appropriate levels of expertise and training. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | Time of year is critical when sampling headwater streams, because precipitation and evapotranspiration can have such profound influences on stream flow. Ideally, sampling would be conducted during both the wettest and driest times of the year to capture the extreme limits of variability in physical conditions. However, if only one field sampling visit is possible, sampling should be conducted during a Spring index period when stream flow is greatest and most aquatic organisms can be collected. | |
| Related Procedures/ References | OEPA. 2002. Field Evaluation Manual for Ohio's Primary Headwater Headwater Habitat Streams, Final Version 1.0. Ohio Environmental Protection Agency, Columbus, OH. http://www.epa.ohio.gov/dsw/wqs/headwaters/index.aspx | |
| Other/ Notes | Although the authors note that land use change within a stream's watershed and the habitat degradation that may result is considered by some authors to be the greatest threat to streams and their biological communities, there is no parameter included in the Manual to estimate or otherwise document land cover or land uses within a watershed of interest. | |

| | | |
|---------------------------------|---|----------------------------|
| Name | Environmental Monitoring and Assessment Program (EMAP), Physical Habitat Characterization | Catalog No 4 |
| Primary Author/ Agency | Kaufmann, P.R. and E.G. Robison. 1998. Physical Habitat Characterization, Section 7 in J.M. Lazorchak, D.J. Klemm, and D.V. Peck (eds), Environmental Monitoring and Assessment Program- Surface Waters: Field Operations and Methods for Measuring the Ecological Condition of Wadeable Streams. U.S. Environmental Protection Agency, EPA/620/R-94/004F, Washington, D.C. | |
| Electronic Resource | http://www.epa.gov/emap/html/pubs/docs/groupdocs/surfwatr/field/ws_abs.html | |
| Intended Use/Purpose | Non-Regulatory Condition Assessment; Ambient Monitoring | |
| Target Resource Type | Wadeable Streams | |
| Scale/Unit of Assessment | Stream reach, 40X low flow wetted width (minimum 150 meters) | |
| Geographic Applicability | Nationwide | |
| General Level of Effort | Moderate | |
| Assessment Parameters | <p>Channel / Valley Morphology Stream discharge; water depth; channel habitat units [bed forms]; pool formative features; wetted channel width; channel gradient; bankfull width; bankfull height; bank height; bank angle; substrate particle size classes (est.); embeddedness (est.); bank undercut distance.</p> <p>Physical Habitat Woody debris tally; areal cover class of fish concealment structures (est.); aerial cover class (est.) of aquatic macrophytes and filamentous algae; riparian vegetative cover (densiometer); relative aerial cover class (est.) and type (e.g. woody trees) of riparian vegetation in canopy, mid-layer, and ground cover.</p> <p>Water Quality Temperature; conductivity; acid neutralizing capacity; dissolved organic carbon; nutrients; turbidity; total suspended solids; color; major cations and anions.</p> <p>Biology - -</p> <p>Other Observation of human disturbance and proximity to stream channel.</p> | |
| Resolution | Qualitative (descriptive); Semi-Quantitative (ordinal scale, rank, etc.); Quantitative (actual measurement or estimate). | |
| Output | Raw data. However, Kaufmann et al. (1999) provide detailed procedures that can be used to calculate metrics related to stream reach and riparian habitat quality using EMAP PHC field data. | |
| Reference | N/A (The objective of the method or procedure is not presented in the context of defining the condition of a resource. However, it may be used to identify or establish reference conditions.) | |
| QA/QC | Kaufmann et al. (1999) discuss the precision associated with EMAP Physical Habitat Characterization measurements and metrics based on extensive field trials in Oregon and the Mid-Atlantic region. | |

| | | |
|---|--|---------------------|
| Name | Environmental Monitoring and Assessment Program (EMAP), Physical Habitat Characterization | Catalog No 4 |
| Description/ Summary | <p>The USEPA Environmental Monitoring and Assessment Program (EMAP) is a research program aimed at developing the tools necessary to monitor and assess the status and trends of national ecological resources. EMAP protocols have been used to assess the regional condition of wadeable streams in the Pacific Northwest, the Mid-Atlantic, the greater 12-State western U.S., and the central U.S. They also served as the basis for the Wadeable Streams Assessment (USEPA, 2006), which was a nationwide State and Federal agency collaborative effort to statistically summarize the condition of the Nation's streams. The EMAP Physical Habitat Characterization (PHC) is one component of the broader EMAP protocols, which also include: water chemistry, stream discharge, periphyton, sediment community metabolism, sediment toxicity, benthic macroinvertebrates, aquatic vertebrates, fish tissue contaminants, and rapid habitat and visual stream assessments (Lazorchak et al. 1998).</p> <p>There are four broad components of EMAP PHC: 1) stream discharge; 2) thalweg profile; 3) large woody debris tally; and 4) channel and riparian characterization. The target stream reach is divided into 10 equally spaced segments with cross-sections established at each union for a total of 11 cross-sections; the first being established at the downstream end of the reach. Stream discharge is measured at a single carefully selected cross-section following methods in Kaufmann (1998). The thalweg profile is a longitudinal survey of depth, channel habitat units, and presence of soft/small sediment at predetermined intervals based on channel width. The woody debris tally is recorded in each of the 10 reach segments between the cross-sections. Channel and riparian characterization includes measures and/or visual estimation of channel dimensions, substrate, fish cover, bank characteristics, riparian vegetation structure, and evidence of human disturbance. These measures are obtained at each of the 11 cross-sections.</p> <p>The EMAP PHC provides very detailed step-by-step instructions for laying out the sample reach and describes what to measure, how to measure, and in what sequence to measure all of the EMAP PHC components. Channel habitat unit classes are defined for the thalweg profile, large woody debris is defined and various "influence zones" are illustrated for the debris tally, and precise descriptions are provided for the whole suite of channel and riparian characterization variables. Comprehensive data forms are provided, and the EMAP PHC provides a list of equipment and supplies necessary to execute the characterization.</p> <p>Finally, the EMAP PHC recommends notation and data entry features and styles to facilitate quantitative statistical assessment and series analysis of the data following methods in Kaufmann et al. (1999).</p> <p>Kaufmann (draft 2001) revised the EMAP PHC as part of a Western Pilot Study Field Operations Manual for Wadeable Streams (Peck et al., Unpublished 2001 Draft). The Western Pilot PHC includes a number of procedural modifications for collecting data on substrate particle size, in-stream fish cover, human influence, and thalweg channel habitat classification. There are also three new PHC metrics in the Western Pilot: 1) size and proximity of large, old riparian trees and occurrence of invasive plant species in the riparian area; 2) degree of geomorphic channel constraint; and 3) evidence of major floods or debris torrents.</p> | |
| Expertise Required | None specified, but the authors stress that the EMAP PHC field methods are easily learned. | |
| Time Necessary to Conduct Assessment | 1.5 to 3.5 hours in the field for a two-person crew | |
| Seasonality | The EMAP PHC field procedures are most efficiently applied during low flow conditions during the vegetative growing season, but they may be applied during other seasons and higher stream flows. | |

| | | |
|--------------------------------------|---|----------------------------|
| Name | Environmental Monitoring and Assessment Program (EMAP), Physical Habitat Characterization | Catalog No 4 |
| Related Procedures/References | <p>Cuffney, T.F, M.E. Gurtz, and M.R. Meador. 1993. Methods for Collecting Benthic Invertebrate Samples as Part of the National Water-Quality Assessment Program. U.S. Geological Survey Open-File Report 93-406, Raleigh, North Carolina.</p> <p>Kaufmann, P.R. Unpublished 2001 Draft. Physical Habitat Characterization, Section 7 In D.V. Peck, J.M. Lazorchak, and D.J. Klemm (eds). Environmental Monitoring and Assessment Program-Surface Waters: Western Pilot Study Field Operations for Wadeable Streams. U.S. Environmental Protection Agency, EPA/xxx/x-xx/xxx, April 2001. Washington, D.C.</p> <p>Kaufmann, P.R., P. Levine, E.G. Robison, C. Seeliger, and D.V. Peck. 1999. Quantifying Physical Habitat in Streams. U.S. Environmental Protection Agency, EPA/620/R-99/003, Washington, D.C.</p> <p>Lazorchak, J.M., D.J. Klemm, and D.V. Peck (eds). 1998., Environmental Monitoring and Assessment Program- Surface Waters: Field Operations and Methods for Measuring the Ecological Condition of Wadeable Streams. U.S. Environmental Protection Agency, EPA/620/R-94/004F, Washington, D.C.</p> <p>USEPA. 2006. Draft Wadeable Stream Assessment: A Collaborative Survey of the Nation's Streams. U.S. Environmental Protection Agency, Office of Water, EPA-841-B-06-002, Washington, D.C.</p> | |
| Other/Notes | <p>EMAP procedures for sampling benthic macroinvertebrates are based on the USEPA Rapid Bioassessment Protocols, but sampling equipment has been modified to allow a single field investigator to conduct the sampling, as recommended by the U.S. Geological Survey National Water Quality Assessment Program (Cuffney et al., 1993).</p> <p>EMAP Aquatic Vertebrate sampling procedures for fish and amphibians utilize the same stream cross-sectional transects as other EMAP procedures. Aquatic vertebrate sampling in wadeable streams utilizes a backpack electro-shocker and block nets or seines. Collection time is based on transect width and should take place for not less than 45 minutes, but no longer than 3 hours.</p> | |

| | | |
|---------------------------------|--|----------------------|
| Name | Methods for Evaluating Stream, Riparian, and Biotic Conditions | Catalog No. 5 |
| Primary Author/ Agency | Platts, W.S., W.F. Megahan, and G.W. Minshall. 1983. Methods for Evaluating Stream, Riparian, and Biotic Conditions. USDA Forest Service Intermountain Forest and Range Experiment Station, General Technical Report INT-138, Ogden, UT. 70 pp. | |
| Electronic Resource | http://www.treesearch.fs.fed.us/pubs/29138 | |
| Intended Use/Purpose | Inventory; Ambient Monitoring | |
| Target Resource Type | Wadeable Streams | |
| Scale/Unit of Assessment | Stream reach of unspecified length. | |
| Geographic Applicability | Nationwide | |
| General Level of Effort | Moderate to Intensive. | |
| Assessment Parameters | <p>Channel / Valley Morphology: Stream discharge; water depth; channel habitat units [bed forms]; percent pool; percent riffle; pool formative features; channel gradient; channel elevation; sinuosity; bank angle; physical bank stability; channel cross-sectional dimensions; stream width; substrate particle size classes (est.); embeddedness (est.); bank undercut distance; vegetative bank stability.</p> <p>Physical Habitat: Woody debris tally; pool quality; in-stream vegetative cover; solar radiation on water surface; riparian vegetative cover type; vegetation overhanging water surface.</p> <p>Water Quality: - -</p> <p>Biology: Vegetation use by animals (est.); herbage production and utilization; fish (numerous sampling methods described); benthic macroinvertebrates (numerous sampling methods described).</p> <p>Other: Stream order.</p> | |
| Resolution | Primarily quantitative (actual measurement or estimate) with some semi-quantitative components. | |
| Output | Raw data | |
| Reference | N/A (The objective of the method or procedure is not presented in the context of defining the condition of a resource. However, it may be used to identify or establish reference conditions.) | |
| QA/QC | An analysis of the accuracy and precision of most of the assessment variables is provided based on time series graphical interpretation of habitat estimates over a 2 to 15 year period in Idaho, Utah, and Nevada relative to the true value of the respective variable. Precision was similarly rated based on confidence intervals obtained for each habitat measurement. | |

| | | |
|---|---|----------------------|
| Name | Methods for Evaluating Stream, Riparian, and Biotic Conditions | Catalog No. 5 |
| Description/ Summary | <p>Platts et al. (1983) set out to propose a "valid, objective, quantitative, repeatable procedure that will provide accurate evaluation of the stream and its biotic communities under any set of conditions." Methods for Evaluating Stream, Riparian, and Biotic Conditions presents standardized techniques for measuring aquatic, riparian, and biotic attributes of stream systems, including fish populations and macroinvertebrate assemblages.</p> <p>Platts et al. (1983) stress transect-based methods for physical stream characterization, whereby channel and riparian zone cross-sections (transects) are established from which one or more physical stream and riparian zone attributes are inventoried as they intersect each transect.</p> <p>The authors do not suggest any means of aggregating data collected using these methods into any specific evaluation of stream condition.</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | Not stated. | |
| Related Procedures/ References | Many of the recommended methods in Platts et al. (1983) have been modified and/or incorporated for use in other stream monitoring and assessment protocols in the two decades since the this manual was published. | |
| Other/Notes | | |

| | | | |
|---------------------------------|---|--|----------------------|
| Name | Wadeable Stream Assessment: Field Operations Manual | | Catalog No. 6 |
| Primary Author/ Agency | <p>U.S. EPA</p> <p>USEPA. 2004a. Wadeable Streams Assessment: Field Operations Manual. EPA-841-B-04-004. U.S. Environmental Protection Agency, Office of Water and Office of Research and Development, Washington, D.C.</p> <p>USEPA. 2006. Wadeable Stream Assessment: A Collaborative Survey of the Nation's Streams. EPA-841-B-06-002. U.S. Environmental Protection Agency, Office of Water, Washington, D.C.</p> | | |
| Electronic Resource | <p>http://www.epa.gov/owow/monitoring/wsa/wsa_fulldocument.pdf</p> | | |
| Intended Use/Purpose | <p>Ambient Monitoring</p> | | |
| Target Resource Type | <p>Wadeable streams; generally 1st thru 3rd order streams (excluding intermittent and ephemeral streams)</p> | | |
| Scale/Unit of Assessment | <p>Stream reach, 40X the channel width, absent of any tributary confluence or impoundment.</p> | | |
| Geographic Applicability | <p>Nationwide</p> | | |
| General Level of Effort | <p>Intense (1 day± in the field by a trained or experienced crew of 2 or more persons)</p> | | |
| Assessment Parameters | <p>Channel / Valley Morphology:</p> <p>Physical Habitat:</p> <p>Water Quality: - -</p> <p>Biology:</p> <p>Other:</p> | <p>Stream discharge; channel gradient; channel sinuosity; channel cross-sectional dimensions; bank height; bank angle; channel habitat units [aka bed forms]; wetted channel width; substrate particle size classes (est.); bank undercut distance;.</p> <p>Woody debris tally; areal cover class of fish concealment structures (est.); embeddedness (est.); riparian vegetative cover (densiometer) and type in canopy, mid-layer, and ground cover; rapid visual-based habitat assessment (RBP).</p> <p>Benthic macroinvertebrates.</p> <p>Presence of anthropogenic disturbance within 10 meters of streambanks.</p> | |
| Resolution | <p>Semi-Quantitative (ordinal scale, rank, etc.): Most of methods included are quantitative, except for the rapid habitat assessment and some other estimates of metrics in lieu of actual measurements.</p> <p>Quantitative (actual measurement or estimate): According to the authors, systematic spatial sampling design for physical habitat measurements collected from channel cross sections scales the sample reach and resolution in proportion to stream size and allows for statistical and series analyses of the data.</p> | | |

| | | |
|-----------------------------|--|----------------------|
| Name | Wadeable Stream Assessment: Field Operations Manual | Catalog No. 6 |
| Output | <p>Index (e.g. Rapid visual-based habitat assessment (RBP habitat assessment); macroinvertebrate numeric score): IBI; macroinvertebrate O/E index; relative bed stability index; riparian disturbance index.</p> <p>Qualitative Description:</p> <p>Raw data: Most of the Field Operation Manual's methods result in raw data and/or field data sheets. Appendix A of USEPA (2006) and Kaufmann et al. (1999) summarize data assessment and formulation of various indices.</p> | |
| Reference | <p>Internal [See Appendix A of USEPA (2006)].</p> <p>Reference conditions for the Wadeable Streams Assessment were defined using data for nine (9) chemical and physical parameters to identify <i>least disturbed conditions</i> per ecoregion. Those nine parameters included total nitrogen, total phosphorus, chloride, sulfate, acid-neutralizing capacity, turbidity, in-stream fish habitat complexity, percent fine substrate, and a riparian disturbance index. Benthic macroinvertebrate assemblages present at those reference sites were then used to develop the condition indices introduced below.</p> | |
| QA/QC | <p>A comprehensive training program that included practice field sampling was instituted prior to data collection activities for the Wadeable Streams Assessment. Each field crew was subsequently audited, and 10% of sample sites were revisited to assess data quality.</p> <p>Comprehensive step-by-step instructions are provided for every step of every field data method proposed. Data forms, recommended guidelines for documenting field data, and comprehensive materials and equipment lists are provided. Instructions for equipment calibration, maintenance, and storage are included. A flow chart illustrating a recommended general sequence of sampling activities per team member is provided, and text further describes logistics and work flow. The Field Operations Manual (USEPA, 2004a) does not itself include any information about data analysis, but recommended methods are outlined in related documents (Appendix A of USEPA (2006) and Kaufmann et al. (1999)).</p> | |
| Description/ Summary | <p>This document describes procedures for collecting data, samples, and information in the field about biotic assemblages and environmental attributes of stream ecosystems that have been used to assess stream conditions over large geographic areas as part of a collaborative State and Federal assessment of the condition of Wadeable streams nationwide. The procedures presented in this manual are based on standard USEPA methods used for the EMAP and REMAP studies. Methods of analysis are summarized in Appendix A of USEPA (2006), and more detailed information on many of the specific indicators used in the Wadeable Streams Assessment is located in Kaufmann et al. (1999). None of these documents by themselves provide a template from which the ecological condition of a given stream in the field can be assessed relative to other streams within a given ecoregion by practitioners who are not associated with USEPA or its partners in the Wadeable Stream Assessment project.</p> <p>Benthic macroinvertebrate assemblages for the Wadeable Streams Assessment were evaluated using a multimetric macroinvertebrate index of biotic condition and a predictive model of taxonomic composition. This model uses a set of least disturbed sites and variables related to natural gradients (e.g. elevation, stream size, stream gradient, latitude, longitude, etc.) to define a taxonomic composition that would be expected in the absence of anthropogenic stressors. The number of expected taxa actually observed at a site is compared to the total number of expected taxa as an Observed:Expected ratio (O/E index). This O/E model was initially developed in Europe and Australia (River Invertebrate Prediction and Classification System, RIVPACS), but is reportedly becoming more commonly used in the U.S.</p> <p style="text-align: center;"><i>(continued on next page)</i></p> | |

| | | |
|---|---|----------------------|
| Name | Wadeable Stream Assessment: Field Operations Manual | Catalog No. 6 |
| Description/ Summary (continued) | <p>Physical habitat data was used to define four condition indicators: streambed excess fine sediment, in-stream habitat cover complexity, riparian vegetation, and riparian human disturbance. Streambed excess fine sediment was assessed using a Relative Bed Stability (RBS) index (Faustini, 2008; Kaufman et al., 2008; 2009), which is a ratio of the median stream reach or riffle particle size diameter divided by the critical bed particle diameter based on streambed shear stress at bankfull flows. In-stream fish habitat cover complexity was based on a measure that sums the amount of instream habitat within one (1) meter of the water surface (Kaufmann et al., 1999). The cover and complexity of riparian vegetation was based on visual estimates of areal vegetative cover and type of vegetation in three strata: canopy, mid-layer, and ground cover (Kaufmann et al., 1999). A Riparian Disturbance Index was used to determine the extent of riparian human disturbance. This index is based on the presence of eleven specific forms of human activities inventoried at 22 separate locations along the sample stream reach, which are weighted according to their proximity to the stream channel (Kaufmann et al. (1999).</p> <p>In addition to field methodology, there is additional information on data-management, safety and health, and other logistical aspects integrated into the methods and overall operational scenario. Specific analytical water chemistry laboratory protocols and benthic macroinvertebrate laboratory protocols are provided in USEPA (2004b) and USEPA (2004c), respectively.</p> | |
| Expertise Required | Not stated. However, proposed sampling, sorting, data reduction, and analysis of biological community assemblages should only be undertaken by persons with appropriate levels of expertise and training. | |
| Time Necessary to Conduct Assessment | Field sampling = 1 day; 2 to 3 persons | |
| Seasonality | Stream sampling for the Wadeable Streams Assessment survey was conducted during a summer index period between 2000 and 2004. | |
| Related Procedures/References | <p>Faustini, J. M. P.R. Kaufmann, and D.P. Larsen. 2008. Using a Relative Bed Stability Index to define reference conditions for assessing anthropogenic sedimentation, American Geophysical Union, Fall Meeting 2008.</p> <p>Kaufmann, P.R., P. Levine, E.G. Robison, C. Seeliger, and D.V. Peck. 1999. Quantifying physical habitat in wadeable streams. EPA/620/R-99/003. U.S. Environmental Protection Agency, Washington, D.C.</p> <p>Kaufmann, P.R., and E.G. Robison. 1998. Physical habitat characterization, Section 7 in J.M. Lazorchak et al., (eds.), Environmental monitoring and assessment program surface waters, field operations and methods for measuring the ecological condition of wadeable streams. EPA/620/R-94/004F. U.S. Environmental Protection Agency, Washington, D.C.</p> <p>Kaufmann, P.R., J.M. Faustini, D.P. Larsen, and M.A. Shirazi. 2008. A roughness-corrected index of relative bed stability for regional stream surveys. <i>Geomorphology</i> 99: 150-170.</p> <p>Lazorchak, J.M., D.J. Klemm, and D.V. Peck. 1998. Environmental monitoring and assessment program surface waters, field operations and methods for measuring the ecological condition of wadeable streams. EPA/620/R-94/004F. U.S. Environmental Protection Agency, Washington, D.C.</p> <p>Kaufmann, P.R., D.P. Larsen, and J.M. Faustini. 2009. Bed stability and sedimentation associated with human disturbances in Pacific Northwest streams. <i>Journal of the American Water Resources Association</i> 45(2): 434-459.</p> | |

| | | |
|--|---|----------------------|
| Name | Wadeable Stream Assessment: Field Operations Manual | Catalog No. 6 |
| Related Procedures/References (continued) | <p>Peck, D.V., J.M. Lazorchak, and D.J. Klemm (editors). Unpublished 2001 draft. Environmental Monitoring and Assessment Program -Surface Waters: Western Pilot Study Field Operations Manual for Wadeable Streams. EPA/XXX/X-XX/XXXX. U.S. Environmental Protection Agency, Washington, D.C.</p> <p>USEPA. 2004b. Wadeable Stream Assessment: Benthic Laboratory Methods. EPA841- B-04-007. U.S. Environmental Protection Agency, Office of Water and Office of Research and Development, Washington, DC.</p> <p>USEPA. 2004c. National Wadeable Stream Assessment: Water Chemistry Laboratory Manual. EPA841- B-04-008. U.S. Environmental Protection Agency, Office of Water and Office of Research and Development, Washington, DC.</p> | |
| Other/Notes | | |

| Name | Watershed Assessment of River Stability and Sediment Supply (WARSSS) | | Catalog No. 7 |
|---------------------------------|--|--|----------------------|
| Primary Author/ Agency | Rosgen, D. 2007. Watershed Assessment of River Stability and Sediment Supply (WARSSS). Wildland Hydrology. Fort Collins, CO. 193 pp. | | |
| Electronic Resource | http://www.epa.gov/warsss/ | | |
| Intended Use/Purpose | Non-Regulatory Condition Assessment (of sediment supply and channel stability); Inventory; Ambient Monitoring. | | |
| Target Resource Type | Not stated. | | |
| Scale/Unit of Assessment | Three-phases: I) Watershed-level reconnaissance; II) Watershed-level inventory; III) Stream reaches, specific hillslopes, etc. | | |
| Geographic Applicability | Nationwide. | | |
| General Level of Effort | Intensive (1 day± in the field by a trained or experienced crew of 2 or more persons) | | |
| Assessment Parameters | Channel/Valley Morphology: | Bankfull stream discharge; Rosgen stream classification; regional curves (bankfull dimensions vs drainage area); bankfull width & depth; radius of curvature; bank height; bank height ratio; cross-sectional channel dimensions; entrenchment ratio; floodprone area width; maximum depth; sinuosity; longitudinal profile; meander length; meander belt width; valley slope; modified Pfankuch channel stability index; bank erosion hazard index (BEHI); near-bank stress (NBS); percent & type of channel alteration; percent of channel blockage (including woody debris, structures, etc.); substrate particle size (pebble count); water surface slope; channel habitat units [bed forms]; pool length & spacing; pool length:riffle width ratio; channel evolutionary stage. | |
| | Physical Habitat: | Percent altered riparian vegetation; length of channel with altered riparian vegetation. | |
| | Water Quality: | Suspended sediment load & bedload [measured using methods in Edwards and Glysson (1999)]. | |
| | Biology: | Riparian species composition and percent coverage per strata. | |
| | Other: | Stream order; watershed area; watershed land use. | |
| Resolution | Qualitative (descriptive); Semi-Quantitative (ordinal scale, rank, etc.); Quantitative (actual measurement or estimate). | | |
| Output | Condition Assessment (of sediment supply and channel stability); Index (e.g. numeric score); Raw data. | | |
| Reference | Regional reference conditions required, but not built-in to the assessment. | | |
| QA/QC | Not stated. | | |

| Name | Watershed Assessment of River Stability and Sediment Supply (WARSSS) | | Catalog No. 7 |
|---|--|--|---------------|
| Description/ Summary | <p>The Watershed Assessment of River Stability and Sediment Supply (WARSSS) was developed by Dave Rosgen with the support of the U.S. Environmental Protection Agency (USEPA). USEPA has developed an internet-based assessment tool using WARSS, which is the principle source of this review.</p> <p>WARSSS utilizes a three-phase approach to assess both suspended and bedload sediment in rivers and streams. Collectively, execution of all three phases of WARSSS may take numerous months and include a multitude of data intensive field investigations and analyses. Results of the assessment can be used to evaluate known or suspected sediment problems, develop sediment remediation and management components of watershed plans, develop sediment TMDLs (Total Maximum Daily Loads), and other uses.</p> <p>Phase I is a Reconnaissance Level Assessment (RLA) that utilizes remote sensing data, published maps, and existing watershed data (e.g. topographic maps, recent and historical aerial photographs, land use/cover and soils maps) to provide a rapid, qualitative assessment of potential sediment sources throughout a watershed.</p> <p>Phase II of the WARSSS is a Rapid Resource Inventory for Sediment & Stability Consequence (RRISSC). The RRISSC phase requires analysis of the type and extent of land uses, the erosion potential of the landscape and channel, and the relationship of potential sediment sources to hillslope, hydrologic and channel processes beginning with target areas identified during the Phase I RLA. A step-by-step risk rating system using a series of worksheets, tables, and relationships of key erosional/depositional process variables is utilized to identify low, moderate, and high risk conditions. The final summary of potential sediment and stream channel stability risk identifies specific areas and stream reaches that may need either mitigation and/or more detailed assessment.</p> <p>The Phase III Prediction Level Assessment (PLA) relies largely on field measurements and is the most detailed level of assessment intended for areas identified as high-risk in the RRISSC. During the PLA, reference conditions are used to determine departure from natural rates of sediment and/or natural channel stability. The PLA analysis ultimately provides data to facilitate the design of well-targeted, site-specific and process-specific management prescriptions. Effectiveness monitoring is critical to compare predicted and observed values and can also be used to determine the effectiveness of the mitigation.</p> <p>The USEPA internet site for WARSSS includes step-by-step instructions for each element of each phase of the assessment, including worksheets, tables, figures, graphs, etc. Background information is provided to familiarize the reader with water quality and biological effects of excessive sediment in rivers and streams. Three case studies are also provided, along with numerous links to additional resources, a glossary, and a considerable bibliography.</p> | | |
| Expertise Required | WARSSS is described as requiring expert judgment that is best undertaken by technical personnel very familiar with sediment sources, processes, and effects. | | |
| Time Necessary to Conduct Assessment | Three-phases: I) >1 day, depending on the size of the watershed being evaluated; II) >1 week, depending on the size of the watershed being evaluated; III) >1 month, depending on the size of the watershed being evaluated. | | |
| Seasonality | Not stated. | | |
| Related Procedures/ References | Edwards, T.K. and G.D. Glysson. 1999. Field Methods for Measurement of Fluvial Sediment, Techniques of Water-Resources Investigations, Book 3, Chapter 2, U.S. Geological Survey. Reston, VA. | | |
| Other/Notes | | | |

| | | |
|---------------------------------|---|--|
| Name | [Vermont] Stream Geomorphic Assessment Protocol Handbooks | Catalog No. 8 |
| Primary Author/ Agency | Vermont Agency of Natural Resources Kline, M., C. Alexander, S. Pomeroy, S. Jaquith, G. Springston, B. Cahoon, and L. Becker. Various Dates (2003, rev. 2004). Stream Geomorphic Assessment Protocol Handbooks. Vermont Agency of Natural Resources, Waterbury, VT. www.vtwaterquality.org/rivers.htm | |
| Electronic Resource | http://www.anr.state.vt.us/dec/waterq/rivers/htm/rv_geoassesspro.htm | |
| Intended Use/Purpose | Non-Regulatory Condition Assessment; Inventory; Ambient Monitoring. | |
| Target Resource Type | Wadeable Streams | |
| Scale/Unit of Assessment | Varies: stream reach to watershed scales | |
| Geographic Applicability | Vermont | |
| General Level of Effort | Easy (rapid); Moderate; Intensive (1 day± in the field by a trained or experienced crew of 2 or more persons). | |
| Assessment Parameters | Channel/Valley Morphology: | Channel hydraulic geometry (plan, pattern, and profile); stream classification; bank slope & bank materials; substrate particle size; rapid geomorphic assessment. |
| | Physical Habitat: | Woody debris tally; rapid visual-based habitat assessment (RBP); riparian buffer width. |
| | Water Quality: | -- |
| | Biology: | -- |
| | Other: | Watershed land use/land cover; river corridor land use). |
| Resolution | Qualitative (descriptive); Semi-Quantitative (ordinal scale, rank, etc.); and Quantitative (actual measurement or estimate). | |
| Output | Semi-quantitative indices representing various geomorphic or physical habitat components; Qualitative Descriptions; and Raw, quantitative data. | |

| | | |
|-----------------------------|---|----------------------|
| Name | [Vermont] Stream Geomorphic Assessment Protocol Handbooks | Catalog No. 8 |
| Reference | <p>Internal: Hydraulic geometry relationships (i.e. regional curves) have been developed and continue to be refined, based on data submitted by users of the Protocols. The Vermont Department of Environmental Conservation has a reference reach program that collects data on geomorphic reference streams statewide, and reports containing reference data from Vermont and other regions by stream type have been drafted.</p> <p>External: Reference stream type must be identified in Phase 1. The reference stream type is defined as the natural stream type that would exist in the absence of anthropogenic changes to the channel, floodplain, and/or watershed. Reference stream type is often based primarily on characteristics of the valley, geology, and climate. Verification and refinement of the reference stream type is made by observing sediment and hydrologic characteristics, as well as channel, floodplain, and terrace land forms during Phases 2 and 3.</p> | |
| QA/QC | <p>The Protocols stress that users should establish a quality assurance (QA) program for each phase of assessment. It further outlines three key components of a good QA program and provides detailed descriptions and recommendations for each: training, data review, and use of a data management system.</p> | |
| Description/ Summary | <p>The purpose of the Stream Geomorphic Assessment Protocols is to provide a method for gathering scientifically sound information that can be used for watershed planning and detailed characterization of riparian and in-stream habitat, stream-related erosion, and flood hazards. The Vermont Agency of Natural Resources (VANR) designed the series of three protocol handbooks to consolidate what had traditionally been distinct river and watershed assessment and resource management programs. Collectively, execution of all phases of the Protocols result in an exhaustive, comprehensive documentation of physical and geomorphic attributes of a stream and its watershed.</p> <p>The Protocols are predicated on a geomorphic stream classification system that VANR developed based on Schumm (1977), Rosgen (1994; 1996), and Montgomery and Buffington (1997) that can be used to generally characterize: 1) the relationship of the stream with its floodplain; 2) the respective roles of bed form, relative channel depth, and stream gradient in sediment transport processes; 3) the size and quantity of sediment in transport; 4) the boundary resistance of the stream bed and banks; and 5) hydrologic runoff characteristics. VANR also developed a channel evolution model adapted from Schumm et al. (1984), Rosgen (1996), and Thorne et al. (1997). Both the classification and the channel evolution model help to frame a "sensitivity rating" that represents a stream's potential rate of change in response to either watershed or local disturbance. Parameters used to rate sensitivity include: 1) erodibility of channel boundary materials; 2) sediment and flow regimes (volume and runoff characteristics); 3) confinement (valley width/channel width); and 4) stage of channel evolution (degree of departure from reference stream type conditions).</p> <p>After first introducing fluvial geomorphic processes, including sediment transport, channel evolution, etc., the Protocols provide three separate, but interrelated approaches for assessing geomorphic and physical habitat conditions of stream reaches and watersheds. Phase 1 is based on remote sensing and very limited, reconnaissance-level, field visits where valley types are identified and geologic conditions investigated to identify provisional stream types. Departure from reference conditions can be postulated based on watershed and stream corridor land use and channel or floodplain modifications. Phase 1 assessments are useful to help prioritize stream reaches for potential Phase 2 assessment, and they also serve as cataloguing databases where the results of Phase 2 and 3 assessments can be entered and tracked on a watershed scale over time.</p> <p>Phase 2 assessments include channel and floodplain cross-sections and stream substrate characterization, all of which is used to identify existing stream type and on-going channel adjustment processes. Qualitative field evaluations of erosion and depositional processes, changes in channel and floodplain geometry, and riparian land use/land cover are used to assess stream geomorphic condition, physical habitat, adjustment processes, reach sensitivity (described previously), and stage of channel evolution. Rapid Habitat Assessment (RHA) and Rapid Geomorphic Assessment (RGA) index values are</p> <p style="text-align: center;"><i>Continued on next page</i></p> | |
| Description/ | | |

| | | |
|---|---|----------------------|
| Name | [Vermont] Stream Geomorphic Assessment Protocol Handbooks | Catalog No. 8 |
| Summary (continued) | <p>also calculated in Phase 2. The RHA is the 10-metric habitat assessment index included as part of the U.S. EPA rapid bioassessment protocols (Barbour et al., 1999). The RGA is based on assessment of 4 to 6 categorical or ordinal metrics that are summed to result in a single index score. Field data sheets and computer database tools have been developed to facilitate Phase 2 data reduction and reporting. The Phase 2 assessment is ideal for identifying stream reaches for protection and restoration projects and the completion of more intensive Phase 3 surveys.</p> <p>Like Phase 2 assessments, Phase 3 assessments are also completed on a stream reach or sub-reach scale. Phase 3 assessments include the use of field survey equipment and other accurate measuring devices and methods to quantify measurements of channel dimension, pattern, profile, and sediments. These are typically undertaken to support requirements for design and implementation of restoration projects. The VANR also uses Phase 3 assessment protocols to develop reference tools (such as regional hydraulic geometry curves). Spreadsheet and database tools have been developed to facilitate Phase 3 data reduction and reporting.</p> <p>Appendices in the Handbooks provide field data forms, database recommendations and instructions, technical information, and detailed techniques and methods for various components of stream geomorphic assessment.</p> | |
| Expertise Required | Technical training is required; Field assistance from VANR specialists is offered on an as available basis. | |
| Time Necessary to Conduct Assessment | Varies Phase 1: based on size of watershed and level of detail; Phase 2: 1 to 2 days per mile of stream length; Phase 3: 3 to 4 days. | |
| Seasonality | Not stated. | |
| Related Procedures/References | <p>Barbour, MT. J. Gerritsen, B.D. Snyder, and J.B. Stribling. 1999. Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition. EPA 841-B-99-002. U.S. Environmental Protection Agency, Office of Water, Washington, D.C.</p> <p>Montgomery, D., and J. Buffington. 1997. Channel-reach Morphology in Mountain Drainage Basins. Geological Society of America Bulletin; v. 109; no. 5; pp 596-611.</p> <p>Rosgen, D.L. 1994. A classification of natural rivers. Catena: 22 169-199.</p> <p>Rosgen D.L. 1996. Applied Fluvial Morphology. Wildland Hydrology. Pagosa Springs, CO.</p> <p>Schumm, S.A. 1969. River Metamorphosis. Proceedings of the American Society of Civil Engineers, Journal of the Hydraulics Division, vol. 95, 255-273.</p> <p>Schumm, S.A. 1977. The Fluvial System. John Wiley and Sons, New York.</p> <p>Schumm, S.A., M.D. Harvey, and C.C. Watson. 1984. Incised Channels Morphology, Dynamics, and Control. Water Resources Publications, Littleton, CO.</p> <p>Thorne, C.R., R.D. Hey, and M.D. Newson. 1997. Applied Fluvial Geomorphology for River Engineering and Management. John Wiley and Sons, Chichester, UK.</p> | |
| Other/Notes | | |

| | | |
|----------------------------------|--|----------------------|
| Name | A Physical Habitat Index for Freshwater Wadeable Streams in Maryland | Catalog No. 9 |
| Primary Author/ Agency: | Maryland Department of Natural Resources (MDNR) Paul, M.J, J.B. Stribling, R.J. Klauda, P.F. Kazyak, M.T. Southerland, and N.E. Roth. 2002. A Physical Habitat Index for Freshwater Wadeable Streams in Maryland. CBWP-MANTA-EA-03-4, Maryland Department of Natural Resources, Monitoring and Non-Tidal Assessment Division, Annapolis, MD. 150 pp. | |
| Electronic Resource: | http://www.dnr.state.md.us/streams/mbss/mbss_pubs.html#technical | |
| Intended Use/Purpose: | Non-Regulatory Condition Assessment; Inventory; Ambient Monitoring. | |
| Target Resource Type: | Wadeable Streams | |
| Scale/Unit of Assessment: | Stream reach of unspecified length. | |
| Geographic Applicability: | Maryland | |
| General Level of Effort: | Easy | |
| Assessment Parameters: | <p><u>Coastal Plain:</u></p> <p>Channel/Valley Morphology: Bank stability.</p> <p>Physical Habitat: In-stream wood; in-stream habitat quality (percent of habitat types present: riffle, run/glide, deep pools, shallow pools, undercut banks, and overhanging cover); epibenthic substrate; shading.</p> <p>Water Quality: - -</p> <p>Biology: - -</p> <p>Other: Remoteness (distance to a road).</p> <p><u>Piedmont:</u></p> <p>Channel/Valley Morphology: Bank stability.</p> <p>Physical Habitat: In-stream wood; in-stream habitat quality; epibenthic substrate; shading; embeddedness; riffle quality.</p> <p>Water Quality: - -</p> <p>Biology: - -</p> <p>Other: Remoteness (distance to a road).</p> <p style="text-align: right;"><i>continued on next page</i></p> | |

| | | |
|---|---|----------------------|
| Name | A Physical Habitat Index for Freshwater Wadeable Streams in Maryland | Catalog No. 9 |
| Assessment Parameters: (continued) | <p><u>Blue Ridge, Ridge and Valley, and Appalachian Plateau:</u></p> <p>Channel/Valley Morphology: Bank stability</p> <p>Physical Habitat: Epibenthic substrate; shading; riparian width.</p> <p>Water Quality: - -</p> <p>Biology: - -</p> <p>Other: Remoteness (distance to a road).</p> | |
| Resolution: | Semi-Quantitative (ordinal scale, rank, etc.); Quantitative (actual measurement or estimate) | |
| Output: | Index (e.g. numeric score) ~ Physical Habitat Index | |
| Reference: | Internal (e.g. Index calibrated to existing local or regional reference data) | |
| QA/QC: | Not stated. | |
| Description/ Summary: | <p>The MDNR Physical Habitat Index (PHI) is not itself a procedure for collecting data. Instead, it is a procedure for analyzing physical habitat data into an index capable of predicting biological stream conditions in Maryland. It is specifically reviewed in this report to illustrate a method of calibrating physical stream assessment data with regional biological stream conditions to develop a physical stream assessment protocol with significant independent utility as a tool to predict biological conditions. The PHI has been subsequently modified by MDNR as a physical habitat assessment component for the Maryland Biological Stream Survey (MBSS) sampling protocols (MDNR, 2007).</p> <p>MDNR developed PHI as a multi-metric physical habitat index capable of discriminating reference stream conditions from degraded stream conditions in Maryland. This work updates and revises a provisional PHI developed by MDNR in 1999 (Hall et al., 1999). The PHI was developed by using biological, chemical, land use, and physical stream habitat data that had been collected throughout the State of Maryland from 1994-2000 using methods described in Roth et al. (1999). Streams were classified based on physiographic setting, and selected criteria were used to represent reference and degraded stream conditions (principally land use). Biological data was specifically avoided during selection of reference sites in order to independently assess the discriminatory efficiency of the PHI and avoid the circularity caused by including biological data in a tool to predict biological conditions. Candidate stream habitat metrics were then identified and tested for their ability to discriminate between reference and degraded conditions. The most discriminating and least redundant metrics were then assembled into a final revised PHI (Paul et al., 2002). Different PHI metrics are used for each of three stream classes based on physiography (see Assessment Parameters above).</p> <p>Some PHI metrics are recorded as counts, measurements, or estimates made in the field, while others are rated using standardized MBSS rating methods. Still others are simple presence/absence observations. The method used for collecting data in the field for each metric differs based according to guidance provided in MDNR (2007).</p> <p>Paul et al. (2002) tested sediment texture and relative bed stability (ratio of the median sediment particle diameter to that diameter moved during channel forming flows (Kaufmann et al., 1999)) in 30 streams (15 Piedmont and 15 Coastal Plain) as potential additional metrics that could predict biological integrity for the PHI. While both of these metrics had significant correlations with a benthic IBI, the sample streams lacked the data necessary to compute the PHI and evaluate, whether they could improve discriminatory or predictive value of the PHI.</p> | |
| Expertise Required: | MDNR (2007) states that only persons who have received MBSS training and have demonstrated proficiency performing MBSS physical habitat assessments should conduct MBSS physical habitat assessments. | |

| | | |
|--|---|----------------------|
| Name | A Physical Habitat Index for Freshwater Wadeable Streams in Maryland | Catalog No. 9 |
| Time Necessary to Conduct Assessment: | Not stated. | |
| Seasonality: | MDNR (2007) states that most MBSS physical habitat assessment information is collected during the Summer index period (March 1 to April 30). However, a number of important measures are rated during the Spring index period (June 1 to September 30). | |
| Related Procedures/References: | <p>Kaufmann, P.R., P. Levine, E.G. Robison, C. Seeliger, and D.V. Peck. 1999. Quantifying Physical Habitat in Streams. U.S. Environmental Protection Division, Office of Research and Development, Washington, DC.. EPA/620/R-99/003.</p> <p>MDNR. 2007. Maryland Biological Stream Survey: Sampling Manual Field Protocols. Maryland Department of Natural Resources, Monitoring and Non-Tidal Assessment Division, CBWP-MANTA-EA-07-01, Annapolis, MD.</p> <p>Roth, N.E., M.T. Southerland, G. Mercurio, J.C. Chaillou, P.F. Kazyak, S.S. Stranko, A.P. Prochaska, D.G. Heimbuch, and J.C. Seibel. 1999. State of the Streams: 1995-1997 Maryland Biological Stream Survey Results. Prepared by Versar, Inc., Post, Buckley, Schuh, and Jernigan, Inc., and Maryland Department of Natural Resources, Monitoring and Non-Tidal Assessment Division. CBWP-MANTA-EA-99-6.</p> | |
| Other/Notes: | Paul et al. (2002) report that the final PHIs were unrelated to watershed area and had an overall discrimination efficiency of 80%. The PHI's were also significantly correlated with indices of biotic integrity for both benthic macroinvertebrates (BIBI) and fish (FIBI). However, the strength of these correlations varied across physiographic regions and even river basins within physiographic regions. | |

| | | |
|---------------------------------|--|-----------------------|
| Name | Physical Habitat and Water Chemistry Assessment Protocol for Wadeable Streams Monitoring Sites | Catalog No. 10 |
| Primary Author/ Agency | Minnesota Pollution Control Agency MPCA. 2002. Physical Habitat and Water Chemistry Assessment Protocol for Wadeable Streams Monitoring Sites. Minnesota Pollution Control Agency, Biological Monitoring Program, December 2002, St. Paul, MN. | |
| Electronic Resource | http://www.pca.state.mn.us/water/biomonitoring/bio-streams-fish.html#sops | |
| Intended Use/Purpose | Ambient Monitoring; WQ Standards. | |
| Target Resource Type | Wadeable streams | |
| Scale/Unit of Assessment | Stream reach, 35X mean stream width (minimum 150 meters, maximum 500 meters) | |
| Geographic Applicability | Minnesota | |
| General Level of Effort | Moderate. | |
| Assessment Parameters | <p>Channel/Valley Morphology: Stream discharge; water depth; mean distance between stream meanders (aka meander wavelength); mean distance between riffles;</p> <p>Physical Habitat: Depth of fines + water (fines >2 mm diameter); embeddedness (to nearest 25%); dominant substrate class (est.); percent algae (est.); percent-cover of fish concealment structures; percent-cover of streambank with exposed soil; total number of channel habitat units (riffles, pools, runs, bends, and log jams); riparian vegetative cover (densiometer); riparian buffer width.</p> <p>Water Quality: Air temperature; water temperature; conductivity; dissolved oxygen; turbidity; pH; transparency; total phosphorus; total suspended solids; ammonia; nitrite-nitrate.</p> <p>Biology: - -</p> <p>Other: Dominant riparian land use</p> | |
| Resolution | Semi-Quantitative (ordinal scale, rank, etc.); Quantitative (actual measurement or estimate). | |
| Output | Index (e.g. numeric score) ~ Stream Habitat Assessment (MPCA, 2007). Raw data. | |
| Reference | N/A (The objective of the method or procedure is not presented in the context of defining the condition of a resource. However, it may be used to identify or establish reference conditions.) | |
| QA/QC | Inexperienced field crew members must receive training. Requisite self-checks whereby field crew personnel cross-reference data collected by other crew members; crew leaders must periodically verify that crew members are adhering to protocol. | |

| | | |
|---|--|-----------------------|
| Name | Physical Habitat and Water Chemistry Assessment Protocol for Wadeable Streams Monitoring Sites | Catalog No. 10 |
| Description/ Summary | <p>The Minnesota Pollution Control Agency (MPCA), Biological Monitoring Program developed the Physical Habitat and Water Chemistry Assessment Protocol for Wadeable Stream Monitoring Sites to support assessment of water quality and development of biological criteria for Minnesota streams. These procedures are also applicable for EMAP stations and sites suspected of being impacted by a source of pollution.</p> <p>Quantitative stream habitat data is collected using a transect-point method modified from "Guidelines for evaluating fish habitat in Wisconsin streams" (Simonson et al., 1993). Thirteen equally spaced transects are established perpendicular to stream flow in the sample reach, and measurements or observations of habitat features are recorded from 0.3 m x 0.3 m quadrats set at four equally spaced points (1/5, 2/5, 3/5, and 4/5 of wetted stream width) and the channel thalweg along each transect. Key habitat features include variables describing channel morphology, substrate, cover, and riparian condition (see Assessment Parameters above).</p> <p>Data forms are provided and must be filled out individually for each transect. A single Station Features data sheet records the length and location (spacing) of major morphological and habitat features within the sample reach, including riffles, runs, pools, meander bends, islands, log jams, beaver dams, and other such features that may affect channel morphology, such as bridges, culverts, dams, and tributaries.</p> <p>MPCA also has a Stream Habitat Assessment (SHA) protocol (MPCA, 2007) based on the Ohio Environmental Protection Agency's Qualitative Habitat Evaluation Index (QHEI) (Rankin, 1989). The SHA assigns scores for many of the stream metrics assessed during the Physical Habitat and Water Chemistry Assessment Protocol (MPCA, 2002) based on aggregate classes of potential results for each metric. The SHA adds a few additional metrics (e.g. surrounding land use within 2-3 square miles of assessment reach) and uses ratios of some existing metrics in order to assign scores (e.g. maximum thalweg depth: shallowest thalweg depth, pool width: riffle width). The maximum SHA score is 100.</p> | |
| Expertise Required | <p>Field technicians must have at least one year of college education and coursework in environmental and/or biological science. Field crew leaders must be a professional aquatic biologist with a minimum of a Bachelor of Science degree in aquatic biology or closely related specialization, and six months field experience sampling physical stream habitat.</p> | |
| Time Necessary to Conduct Assessment | <p>Not stated.</p> | |
| Seasonality | <p>Summer index period: mid-June thru mid-September</p> | |
| Related Procedures/References | <p>MPCA. 2007. Stream Habitat Assessment Protocol for Stream Monitoring Sites. Minnesota Pollution Control Agency, Biological Monitoring Program, March 2007, St. Paul, MN.</p> <p>MPCA. 2009. Reconnaissance Procedures for Initial Visit to Stream Monitoring Sites. Minnesota Pollution Control Agency, Biological Monitoring Program, February 2009, St. Paul, MN.</p> <p>Rankin, E.T. 1989. The Qualitative Habitat Evaluation Index (QHEI): Rationale, Methods, and Application. Ohio Environmental Protection Agency, Division of Water Quality Planning & Assessment, Ecological Assessment Section. Columbus, OH. 73 pp.</p> <p>Simonson, T.D., J. Lyons, and P.D. Kanehl. 1993. Guidelines for Evaluating Fish Habitat in Wisconsin Streams. Gen. Tech. Rpt NC-164, USFS North Central Experiment Station, St. Paul, MN. 36 pp.</p> | |
| Other/Notes | <p>The MPCA Protocol provides a good example of a semi-quantitative physical stream assessment protocol used in a biological monitoring program.</p> | |

| | | |
|---------------------------------|--|-----------------------|
| Name | Field evaluation manual for Ohio's primary headwater habitat streams | Catalog No. 11 |
| Primary Author/ Agency | Ohio Environmental Protection Agency OEPA. 2002a. Field evaluation manual for Ohio's primary headwater habitat streams, Version 1.0, July 2002. Ohio Environmental Protection Agency, Division of Surface Water, Columbus, Ohio | |
| Electronic Resource | http://www.epa.ohio.gov/dsw/wqs/headwaters/index.aspx | |
| Intended Use/Purpose | Non-Regulatory Condition Assessment; Ambient Monitoring | |
| Target Resource Type | Headwater streams with a drainage area less than 1 square mile (ephemeral, intermittent, or perennial) | |
| Scale/Unit of Assessment | Stream reach, 200 feet, or shorter if necessary to avoid channel confluences | |
| Geographic Applicability | Ohio | |
| General Level of Effort | Easy to Moderate A three-tiered protocol is presented with corresponding levels of effort 1) Rapid habitat evaluation referred to as the Headwater Habitat Evaluation Index (HHEI); and two levels of biological assessment, 2) Family-level taxonomic identification; and 3) Genus-species level taxonomic identification. | |
| Assessment Parameters | <p>Channel/Valley Morphology: Bankfull width; channel substrate composition (selected from nine possible categories); maximum pool depth.</p> <p>Physical Habitat: Riparian buffer width; percent open canopy.</p> <p>Water Quality: Temperature; pH; conductivity; dissolved oxygen.</p> <p>Biology: Fish; salamanders; benthic macroinvertebrates (as necessary).</p> <p>Other: Floodplain land use; development pressure.</p> | |
| Resolution | Dependent on which of three-tier level of assessment is undertaken: Qualitative (descriptive) Semi-Quantitative (ordinal scale, rank, etc.) Quantitative (actual measurement or estimate) | |
| Output | Index (e.g. numeric score); Qualitative Description; Raw data; and Programmatic or Regulatory Support Information (WQ standards) | |
| Reference | Internal (e.g. Index calibrated to existing local or regional reference data) | |
| QA/QC | Not stated. | |

| | | |
|---|--|-----------------------|
| Name | Field evaluation manual for Ohio's primary headwater habitat streams | Catalog No. 11 |
| Description/ Summary | <p>The Field Evaluation Manual for Ohio's Primary Headwater Habitat Streams is intended to promote standardized assessment of actual and expected biological conditions in primary headwater habitat (PHWH) streams in Ohio. The methods outlined in the Manual are designed solely to statistically differentiate among three designated uses of PHWH streams in Ohio, as defined in State water quality standards:</p> <p style="padding-left: 40px;">Class III PHWH Stream (cool-cold water adapted native fauna); Class II PHWH Stream (warm water adapted native fauna); Class I PHWH Stream (ephemeral stream, normally dry channel).</p> <p>Chemical, biological, and physical habitat evaluations were conducted in PHWH streams throughout Ohio to assess seasonal trends in benthic macroinvertebrate assemblage. Statistical analysis is provided in OEPA (2002b; 2002c; 2002d).</p> <p>The Headwater Habitat Evaluation Index (HHEI) is a rapid habitat evaluation tool based on three physical measurements found to be highly correlated with biological measures of PHWH stream quality in Ohio: i) channel substrate composition; ii) maximum pool depth; and iii) average bankfull width (OEPA, 2002d). The HHEI rapid assessment tool is most predictive when "modified" channels are separated from natural channels having little or no evidence of channel modification. Specific methods are presented for each of the above referenced parameters. Index scores are compared to categories defining each of the above referenced classes of PHWH Streams.</p> <p>All PHWH evaluations also include assessment of riparian zone and floodplain quality (i.e. width and land use), flow regime, sinuosity, and gradient, although none of these factors are included in the calculation of the HHEI score. All of these parameters are simply categorical check-boxes.</p> <p>If the HHEI assessment is questionable, or additional support for the designated use category determined using the HHEI is desired, one can conduct a Headwater Macroinvertebrate Field Evaluation Index (HMFEI) and a rapid bioassessment of vertebrates (salamanders) using one of two tiers of effort presented in the Manual. Specific sampling protocols for each are dutifully referenced. If watershed size is greater than 1.0 square mile or natural deep pools are greater than 40 cm regardless of watershed size, a Qualitative Habitat Evaluation Index (QHEI) evaluation should be completed (Rankin, 1989).</p> <p>Data forms and detailed instructions are provided. There is also a suggested step-by-step procedure for executing an entire assessment, and there is a decision making flowchart to determine appropriate PHWH stream class using the HHEI.</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Varies; dependent on which of three-tier level of assessment is undertaken. | |
| Seasonality | June to September is optimal for biological component(s) of the assessment | |

| | | |
|--------------------------------------|---|-----------------------|
| Name | Field evaluation manual for Ohio's primary headwater habitat streams | Catalog No. 11 |
| Related Procedures/References | <p>OEPA. 2002b. Technical support document for Ohio's primary headwater streams: fish and amphibian assemblages. Ohio Environmental Protection Agency, Division of Surface Water, Columbus, Ohio.</p> <p>OEPA. 2002c. Technical support document of Ohio's primary headwater streams benthic: macroinvertebrate assemblage. Ohio Environmental Protection Agency, Division of Surface Water, Columbus, Ohio.</p> <p>OEPA. 2002d. Ohio EPA Primary Headwater Habitat Initiative Data Compendium, 1999-2000 Habitat, Chemistry, and Stream Morphology Data. Ohio Environmental Protection Agency, Division of Surface Water, Columbus, Ohio.</p> <p>Rankin, E. 1989. The qualitative habitat evaluation index (QHEI): Rational, methods, and applications. Ohio Environmental Protection Agency, Division of Surface Water, Columbus, Ohio.</p> | |
| Other/Notes | <p>An attempt to relate Rosgen stream class with PHWH stream class was inconclusive; attributed by the authors to most likely be because the Rosgen system was not calibrated to the small watershed size (<1.0 square mile) of PHWH streams (OEPAC, 2002).</p> | |

| | | |
|---------------------------------|---|-----------------------|
| Name | The Qualitative Habitat Evaluation Index (QHEI): Rationale, Methods, and Application | Catalog No. 12 |
| Primary Author/ Agency | <p>Ohio Environmental Protection Agency</p> <p>OEPA. 2006. Methods for Assessing Habitat in Flowing Waters: Using the Qualitative Habitat Evaluation Index (QHEI). OEPA Technical Bulletin EAS/2006-06-1, Ohio Environmental Protection Agency, Division of Surface Water, Ecological Assessment Section. Columbus, OH. 26 pp.</p> <p>Rankin, E.T. 1989. The Qualitative Habitat Evaluation Index (QHEI): Rationale, Methods, and Application. Ohio Environmental Protection Agency, Division of Water Quality Planning & Assessment, Ecological Assessment Section. Columbus, OH. 73 pp.</p> | |
| Electronic Resource | <p>http://www.epa.ohio.gov/dsw/bioassess/BioCriteriaProtAqLife.aspx</p> | |
| Intended Use/Purpose | <p>Ambient Monitoring; WQ Standards.</p> | |
| Target Resource Type | <p>Wadeable and non-wadeable streams, although correlations with a fish IBI in Ohio has been found to be weaker in small streams.</p> | |
| Scale/Unit of Assessment | <p>Stream reach of unspecified length.</p> | |
| Geographic Applicability | <p>Ohio. However, its use is reported to now include some adjacent states.</p> | |
| General Level of Effort | <p>Easy (rapid)</p> | |
| Assessment Parameters | <p>Channel/Valley Morphology: Sinuosity (categorical classes); presence/absence or recovery state following channelization; channel stability; bank stability; channel gradient; substrate (type/size class, origin, & quality); predominance and development of riffle/pool complexes; pool/glide and riffle/run quality (max pool or glide depth, riffle width & depth, run depth, riffle/run substrate size class, riffle/run embeddedness, flow velocity class).</p> <p>Physical Habitat: In-stream cover (type and percent-cover class); riparian buffer width; floodplain cover type.</p> <p>Water Quality: - -</p> <p>Biology: - -</p> <p>Other: Watershed area.</p> | |
| Resolution | <p>Qualitative (descriptive); Semi-Quantitative (ordinal scale, rank, etc.).</p> | |
| Output | <p>Index (e.g. numeric score); Qualitative Description.</p> | |

| | | |
|---|---|-----------------------|
| Name | The Qualitative Habitat Evaluation Index (QHEI): Rationale, Methods, and Application | Catalog No. 12 |
| Reference | N/A (The objective of the method or procedure is not presented in the context of defining the condition of a resource. However, it may be used to identify or establish reference conditions.) | |
| QA/QC | Rankin (1989) stresses that regular training is a necessity to minimize bias and ensure comparability of assessments among field biologists. Field data sheet headers require that survey crew members indicate their level of QHEI training. At least one crew member must have completed OEPA QHEI training (OEPA, 2006). | |
| Description/ Summary | <p>The Qualitative Habitat Evaluation Index (QHEI) is an index of macro-habitat quality intended to assess stream habitat that is generally accepted to influence fish communities and which is also important to other aquatic life (Rankin, 1989). It was designed as a measure that would require a minimal amount of time and with a minimum of field measurements, but also relies upon experienced field biologists to execute the evaluation within acceptable ranges of accuracy and precision.</p> <p>The QHEI is based on emergent properties of habitat (e.g. sinuosity, pool/riffle development) rather than the individual metrics that shape these properties (e.g. current velocity, depth, substrate size, etc.). A field data sheet, modified in OEPA (2006), provides qualitative condition descriptors for 1 to 7 variables under each of six stream properties. The field surveyor matches the condition description for each variable with observed conditions in the field and checks the appropriate box. Each box includes an affiliated point score. Point scores are totaled for each metric to provide subtotals related to the above six stream properties. The sum of all metric subtotals provides the total QHEI score, which has a maximum of 100. More detailed definitions of terms used on the field data sheet, including broader descriptions and illustrations or drawings of each variable, are provided by OEPA (2006).</p> <p>The QHEI was found to be significantly different among Ohio ecoregions and significantly correlated with fish IBI (Rankin, 1989). However, the correlation with the fish IBI was weaker in wadeable and headwater streams relative to larger channels requiring boat access. Rankin (1989) suggests that due to the inherent interconnectedness of smaller channels with their watersheds and riparian zones, disturbances outside of the stream channel itself may exert a more prominent impact on the biological community, thus affecting IBI more than QHEI and thereby adversely affecting the correlation of the two. Rankin (1989) also notes that general basin characteristics and overall habitat quality exert a greater influence on fish communities than does site specific habitat, such as that assessed using the QHEI. Thus, he concludes, the QHEI (or any other site specific habitat measure) is not inclusive enough to be an absolute site specific predictor of fish communities without further consideration of basin-wide or reach-wide influences on stream biota (Rankin, 1989).</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | Not stated. | |
| Related Procedures/ References | Ohio EPA. 1989. Biological criteria for the protection of aquatic life: Volume III. Standardized biological field sampling and laboratory methods for assessing fish and macroinvertebrate communities. Ohio Environmental Protection Agency, Columbus, OH. | |
| Other/Notes | | |

| | | |
|---------------------------------|--|-----------------------|
| Name | Guidelines for Evaluating Fish Habitat in Wisconsin Streams | Catalog No. 13 |
| Primary Author/ Agency | USFS Simonson, T.D., J. Lyons, and P.D. Kanehl. 1993. Guidelines for Evaluating Fish Habitat in Wisconsin Streams. Gen. Tech. Rpt NC-164, USFS North Central Experiment Station, St. Paul, MN. 36 pp. | |
| Electronic Resource | http://www.treesearch.fs.fed.us/pubs/10228 | |
| Intended Use/Purpose | Inventory; Ambient Monitoring. | |
| Target Resource Type | Perennial wadeable streams (ideally >1.5m wide with watersheds >13km ²) | |
| Scale/Unit of Assessment | Stream reach, 35X low flow wetted width (minimum 100 meters) | |
| Geographic Applicability | Wisconsin | |
| General Level of Effort | Moderate. | |
| Assessment Parameters | <p>Channel / Valley Morphology: Stream discharge; stage; velocity; wetted width; water depth; channel gradient; mean distance between bends (aka meander wave length); length and spacing of channel habitat units (aka bed forms); percent substrate particle size classes (est.).</p> <p>Physical Habitat: Bank vegetative protection; embeddedness (est.); fine sediment depth; percent cover and types of fish concealment structures; riparian buffer width; canopy cover (densiometer).</p> <p>Water Quality: Dissolved oxygen; temperature; conductivity, turbidity.</p> <p>Biology: - -</p> <p>Other: Stream order; riparian land use; watershed area.</p> | |
| Resolution | Semi-Quantitative (ordinal scale, rank, etc.); Quantitative (actual measurement or estimate). | |
| Output | Subjective Index (e.g. numeric score); Qualitative Description; Programmatic or Regulatory Support Information | |
| Reference | Not stated. However, the River Fish Habitat Rating (FHR) index was internally calibrated to the Wisconsin fish IBI. | |
| QA/QC | Not stated. | |

| | | |
|---|--|-----------------------|
| Name | Guidelines for Evaluating Fish Habitat in Wisconsin Streams | Catalog No. 13 |
| Description/ Summary | <p>Simonson et al. (1993) recommend that habitat data be collected using the basic framework of the transect method suggested by Platts et al. (1983), where sampling is based on transects spaced two times the mean wetted stream width throughout the sample reach, for a total of at least 18 sample transects per reach. Accuracy of sampling small streams (<10m wide) is not compromised by sampling transects spaced every three times the mean wetted width, but the authors do not recommend any fewer than 18 transects on larger channels (Simonson et al., 1993).</p> <p>Stream habitat characteristics are measured or estimated from one or more locations relative to each transect: 1) within a specified distance above and below the transect, 2) along the transect (e.g., 5m total belt width), or 3) at positions along the transect line, typically four equally spaced positions across the channel, plus the thalweg. Methods to measure or estimate each habitat characteristic are suggested, and the authors also report the accuracy and precision of each method based on their own analysis of survey results.</p> <p>Simonson et al. (1993) provide field data sheets and also discuss data management and analysis. The authors also present Fish Habitat Rating (FHR) indices based on actual field measurements as a means to compare habitat surveys by rating the physical habitat of streams and rivers to support diverse, healthy fish communities. Two different FHR indices are presented one for streams less than 10 meters wide, and a second for rivers 10 to 50 meters wide. The Stream FHR is based on seven selected variables or ratios that are rated poor, fair, good, or excellent based on reference conditions provided in the Guidelines: 1) riparian buffer width, 2) bank erosion, 3) pool area, 4) width/depth ratio, 5) riffle-to-riffle ratio or bend-to-bend ratio (average distance between riffles or bends divided by average stream width), 6) percent fine sediment, and 7) cover for fish. Points are allocated to each quality category and then summed to obtain a total Stream FHR index. The River FHR is based on five selected variables or ratios, including bank stability, maximum thalweg depth, riffle-to-riffle ratio or bend-to-bend ratio, percent rocky substrate, and cover for fish.</p> | |
| Expertise Required: | Not stated. | |
| Time Necessary to Conduct Assessment | 2 to 4 hours. | |
| Seasonality | Baseflow conditions, ideally during Summer. | |
| Related Procedures/References | Platts, W.S., W.F. Megahan, and G.W. Minshall. 1983. Methods for Evaluating Stream, Riparian, and Biotic Conditions. USDA Forest Service Intermountain Forest and Range Experiment Station, General Technical Report INT-138, Ogden, UT. 70 pp. | |
| Other/Notes | The Wisconsin Department of Natural Resources "Guidelines for Evaluating Habitat of Wadeable Streams" closely mirrors Simonson et al. (1993). | |

| | | |
|----------------------------------|--|-----------------------|
| Name | Physical Habitat of Aquatic Ecosystems (Texas) | Catalog No. 14 |
| Primary Author/ Agency | Texas Commission on Environmental Quality (TCEQ) TCEQ. 2007. Physical Habitat of Aquatic Ecosystems, Chapter 9 in Surface Water Quality Monitoring Procedures, Vol. 2: Methods for Collecting and Analyzing Biological Assemblage and Habitat Data. RG-416, Texas Commission on Environmental Quality, Monitoring Operations Division, June 2007. | |
| Electronic Resource | http://www.tceq.state.tx.us/comm_exec/forms_pubs/pubs/rg/rg-416/index.html | |
| Intended Use/Purpose | Non-Regulatory Condition Assessment; Inventory; Ambient Monitoring; WQ Standards. | |
| Target Resource Type | Wadeable and Non-wadeable streams. This habitat assessment procedure may be used unmodified in non-flowing streams if water is present in pools covering >50 percent of the sample reach (~intermittent with pools). | |
| Scale/Unit of Assessment: | Wadeable Streams: Stream reach, 40x average stream width; not less than 150m and not more 500m (avoiding significant tributary confluences, bridge crossings, etc.) Non-wadeable Streams: Stream reach encompassing one full meander; not less than 500m and not more than 1km (avoiding significant tributary confluences, bridge crossings, etc.) | |
| Geographic Applicability | Texas | |
| General Level of Effort | Easy to Moderate | |
| Assessment Parameters | <p>Channel / Valley Morphology: Stream discharge; wetted channel width; water depth; channel flow status; channel habitat units [aka bed forms]; maximum pool width; maximum pool depth; maximum pool length; number of riffles; number of flow obstructions; percent bank erosion (est.); dominant substrate particle size class (est.); percent of substrate that is gravel or larger (> 2mm) (est.); channel gradient; bank angle; number and definition of stream bends.</p> <p>Physical Habitat: Percent-cover and type of vegetation on stream banks and in riparian zone; percent canopy cover (densiometer); riparian buffer width; percent and type of in-stream cover; ordinal est. of algae and macrophyte percent-cover.</p> <p>Water Quality: Temperature, pH; dissolved oxygen; specific conductance; salinity.</p> <p>Biology: - -</p> <p>Other: Stream order; watershed area; categorical riparian zone aesthetics.</p> | |
| Resolution | Semi-quantitative | |
| Output | Index (e.g. numeric score)~ Habitat Quality Index | |
| Reference | Internal to the Habitat Quality Index. | |

| | | |
|---|--|-----------------------|
| Name | Physical Habitat of Aquatic Ecosystems (Texas) | Catalog No. 14 |
| QA/QC | Completion of an annual self-audit report (administrative and record keeping); an annual technical systems audit, both in the field and laboratory; and TCEQ approval of a Quality Assurance Project Plan. Biological voucher specimens and use of specific taxonomic keys are required. | |
| Description/ Summary | <p>TCEQ uses habitat data collected according to these methods, in conjunction with fish and benthic macroinvertebrate community surveys, to provide a holistic evaluation of the health of stream biological assemblages and to develop future indices of aquatic life use. Fish (TCEQ, 2007, Ch. 3) are sampled using both electrofishing and seining, and data is evaluated using a regionalized fish IBI for Texas streams (Linam et al., 2002). Benthic macroinvertebrate (TCEQ, 2007, Ch. 5) sampling is conducted following USEPA RBP protocols (Barbour et al., 1999) and assessed as a benthic IBI. In-situ physiochemical water quality is monitored according to TCEQ (2008, Ch. 3).</p> <p>Sampling is conducted from 5 to 11 channel cross-sections equally spaced throughout the reach. Part I of the assessment utilizes Stream Physical Characteristics Worksheets to record in-stream channel measurements, stream morphology, and riparian environment attributes for each transect or for the entire reach, following methods generally derived from USEPA EMAP protocols (Kaufmann and Robison, 1998). These measurements and estimates are averaged and summarized to complete the Summary of Physical Characteristics of Water Body in Part II. Then in Part III, a Habitat Quality Index (HQI) is calculated based on the values summarized in Part II.</p> | |
| Expertise Required | Training is offered by TCEQ, and required of all monitoring participants periodically. However, the regularity of requisite training is not specified. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | The TCEQ Physical Habitat procedures are intended to be conducted as part of biological assessments, and those assessments should be undertaken during the index period between March 15 and October 15. If only one assessment can be undertaken at a monitoring station, biological data should be collected between July 1 and September 30. | |
| Related Procedures/ References | <p>Barbour, M.T., J. Gerritsen, B.D. Snyder, and J.B. Stribling. 1999. Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition. EPA 841-B-99-002. U.S. Environmental Protection Agency, Office of Water, Washington, D.C.</p> <p>Kaufmann, P.R. and E.G. Robison. 1998. Physical Habitat Characterization, Section 7 in J.M. Lazorchak, D.J. Klemm, and D.V. Peck (eds), Environmental Monitoring and Assessment Program-Surface Waters: Field Operations and Methods for Measuring the Ecological Condition of Wadeable Streams. U.S. Environmental Protection Agency, EPA/620/R-94/004F, Washington, D.C.</p> <p>Linam, G.W., L.J. Kleinsasser, and K.B. Mayes. 2002. Regionalization of the Index of Biotic Integrity for Texas Streams. River Studies Report No. 17., Texas Parks and Wildlife Department, Austin, Texas.</p> <p>TCEQ. 2008. Surface Water Quality Monitoring Procedures, Vol. 1: Physical and Chemical Monitoring Methods. RG-415, Texas Commission on Environmental Quality, Water Quality Planning Division, October 2008.</p> | |
| Other/Notes | | |

| | | |
|---------------------------------|---|--|
| Name | [Kansas] Subjective Evaluation of Aquatic Habitats | Catalog No. 15 |
| Primary Author/ Agency | Kansas Department of Wildlife & Parks KDWP. 2004. Subjective Evaluation of Aquatic Habitats. Kansas Department of Wildlife & Parks, Environmental Services Section, revised 2004. Topeka, KS. | |
| Electronic Resource | http://www.kdwp.state.ks.us/news/Other-Services/Environmental-Reviews/Aquatic-Field-Habitat-Evaluations | |
| Intended Use/Purpose | Non-Regulatory Condition Assessment; Inventory; Ambient Monitoring. | |
| Target Resource Type | Streams: Ephemeral, Intermittent, or Perennial | |
| Scale/Unit of Assessment | Not stated. | |
| Geographic Applicability | Kansas | |
| General Level of Effort | Easy (rapid) | |
| Assessment Parameters | Channel/Valley Morphology: | Channel modification; sinuosity (via map); percent of historical floodplain available of inundation; dominant substrate class; number of substrate types; pool:riffle sequencing; bank erosion. |
| | Physical Habitat: | embeddedness (class est.); in-stream cover types and percent cover (aka fish concealment structures); canopy cover (est.); percent of historical floodplain covered by native vegetation. |
| | Water Quality: | Condition classes based on professional judgment: dissolved oxygen / biochemical oxygen demand; nutrient enrichment; pesticides; turbidity; temperature; other. |
| | Biology: | Fish community characteristics (professional judgment); benthic invertebrates (dominant taxa); freshwater mussels (presence/absence); amphibians (presence/absence); other aquatic vertebrates (presence/absence). |
| | Other: | Stream type (ephemeral, intermittent, or perennial); floodplain land use classes; watershed land use classes. |
| Resolution | Qualitative (descriptive); Semi-Quantitative (ordinal scale, rank, etc.). | |
| Output: | Index (e.g. numeric score) | |
| Reference | Best Professional Judgment | |
| QA/QC | Not stated. | |

| | | |
|---|--|-----------------------|
| Name | [Kansas] Subjective Evaluation of Aquatic Habitats | Catalog No. 15 |
| Description/ Summary | <p>The Kansas Department of Wildlife & Parks (KDWP), Subjective Evaluation of Aquatic Habitats utilizes a four groups of individual parameters that are scored and then summed to provide a total stream habitat index (R-value). The R-value index is in turn associated with four overall stream habitat condition classes: excellent, good, fair, and poor.</p> <p>The number of points possible varies among the groups, from 50 points for the Physical Habitat Key to 15 points each for both the Biological Component Key and the Water Quality Component key. Each parameter within each group is scored based on qualitative, categorical, ranked conditions or classes as described in the document and outlined on the field data sheet. The Water Quality Component Key and the Biological Component Key, which includes a fish community parameter and a benthic invertebrate parameter, as well as a presence/absence of freshwater mussels, amphibians and other aquatic vertebrates, are not to be included in the final R-value rating if the stream is dry or inadequate water is present.</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | Not stated. | |
| Related Procedures/ References | | |
| Other/Notes | <p>The U.S. Army Corps of Engineers, Kansas City District "Draft Kansas Stream Mitigation Guidance," (rev. December 31, 2009) utilizes the KDWP R-value stream habitat index as one factor for determining the "Existing Condition" of streams either proposed to be impacted or to be used for compensatory mitigation as part of Clean Water Act, Section 404 permit applications. The "Draft Kansas Stream Mitigation Guidance" is a standard operating procedure modeled after the USACE Charleston District SOP reviewed herein.</p> | |

| | | |
|---------------------------------|---|-----------------------|
| Name | [PIBO] Effectiveness monitoring for streams and riparian areas: sampling protocol for stream channel attributes | Catalog No. 16 |
| Primary Author/ Agency | U.S. Forest Service Heitke, J.D., E.J. Archer, D.D. Dugaw, B.A. Bouwes, E.A. Archer, R.C. Henderson, and J.L. Kershner. 2008. Effectiveness monitoring for streams and riparian areas: sampling protocol for stream channel attributes. PACFISH/INFISH Biological Opinion (PIBO) Effectiveness Monitoring Program, Multi-federal Agency Monitoring Program; Logan, UT. Unpublished paper on file at: http://www.fs.fed.us/biology/fishecology/emp . | |
| Electronic Resource | http://www.fs.fed.us/biology/fishecology/emp | |
| Intended Use/Purpose | Inventory | |
| Target Resource Type | Wadeable Streams | |
| Scale/Unit of Assessment | Stream reach, minimum length of 20X bankfull width based on width classes (525 feet min length) | |
| Geographic Applicability | Interior Columbia River basin ~ Washington, Oregon, and most of Idaho, as well as western Montana, northeastern Nevada, and northwestern Wyoming | |
| General Level of Effort | Moderate to Intensive | |
| Assessment Parameters | <p>Channel/Valley Morphology: Channel gradient; bankfull width; bankfull depth; width/depth ratio; entrenchment ratio; reach length & valley length [allows for calculation of sinuosity]; substrate particle size (pebble counts); pool length & residual pool depth; undercut depth; bank type; bank material; bank angle; bank stability.</p> <p>Physical Habitat: Woody debris tally; percent surface fines on pool tails.</p> <p>Water Quality: Conductivity; alkalinity</p> <p>Biology: Benthic macroinvertebrates.</p> <p>Other: - -</p> | |
| Resolution | Quantitative (actual measurement or estimate) | |
| Output | Raw data | |
| Reference | N/A (The objective of the method or procedure is not presented in the context of defining the condition of a resource. However, it may be used to identify or establish reference conditions.) | |
| QA/QC | Not stated. | |

| | | |
|---|--|-----------------------|
| Name | [PIBO] Effectiveness monitoring for streams and riparian areas: sampling protocol for stream channel attributes | Catalog No. 16 |
| Description/ Summary | <p>The primary objective of the PACFISH/INFISH (PIBO) Effectiveness Monitoring Program is to determine whether priority biological and physical attributes, processes, and functions of riparian and aquatic systems are being degraded, maintained, or restored on federally managed lands within the interior Columbia River basin. This document describes the standardized methods that PIBO compiled following ten years of use, evaluation, and peer review, as well as a set of summary statistics for each attribute.</p> <p>The PIBO Effectiveness Monitoring protocols utilize transect-based methods for measuring physical habitat and geomorphic metrics. Stepwise instructions are thorough and include illustrative figures for clarification. Although many of the methods reported for specific metrics are modifications of methods proposed by others (e.g. Platts et al, 1987), the PIBO Effectiveness Monitoring protocols have typically further refined such methods to reduce bias and increase measurement precision. There is also a section devoted to explaining a proper method to photo-document the sample reach.</p> <p>Equipment lists, field data forms, decontamination procedures, and data management is discussed. There are also alternative sampling methods provided for sampling stream reaches affected by beaver.</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | Not stated. | |
| Related Procedures/ References | <p>Henderson, R.C.; E.K. Archer, B.A. Bouwes, M.S. Coles-Ritchie, and J.L. Kershner. 2005. PACFISH/INFISH Biological Opinion (PIBO): Effectiveness Monitoring Program seven-year status report 1998 through 2004. Gen. Tech. Rep. RMRS-GTR-162. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 16 pp.</p> <p>Kershner, J.L., E.K. Archer, M. Coles-Ritchie, E.R. Cowley, R.C. Henderson, K. Kratz, C.M. Quimby, D.L. Turner, L.C. Ulmer, M.R. Vision. 2004. Guide to effective monitoring of aquatic and riparian resources. General Technical Report RMRS-GTR-121. U.S. Department of Agriculture Forest Service, Fort Collins, CO.</p> <p>Platts, W.S., C. Armour, G.D. Booth, M. Bryant, J.L. Bufford, P. Cuplin, S. Jensen, G. W. Lienkaemper, G.W. Minshall, S.P. Monsen, R.L. Nelson, J.R. Sedell, and J.S. Tuhy. 1987. Methods for Evaluating Riparian Habitats with Applications to Management. U.S. Forest Service, Intermountain Research Station General Technical Report INT-221. 177 pp.</p> <p>USFS. 2004. Effectiveness monitoring for streams and riparian areas within the Pacific Northwest: stream channel methods for core attributes. Aquatic and Riparian Effectiveness Monitoring Program (AREMP) & PACFISH/INFISH (PIBO) Effectiveness Monitoring Program, Multi-Federal Agency Monitoring Programs. U.S. Department of Agriculture, Forest Service. Unpublished paper available at: http://www.reo.gov/monitoring/reports/watershed-reports-publications.shtml</p> | |
| Other/Notes | | |

| | | |
|---------------------------------|--|-----------------------|
| Name | R1/R4 (Northern /Intermountain Regions) Fish and Fish Habitat Standard Inventory Procedures Handbook | Catalog No. 17 |
| Primary Author/ Agency | U.S. Forest Service Overton, C.K., S.P. Wollrab, B.C. Roberts, and M.A. Radko. 1997. R1/R4 (Northern and Intermountain Regions) fish and fish habitat standard inventory procedures handbook. Gen. Tech. Rep. INT-GTR-346. U.S.D.A. Forest Service, Intermountain Research Station, Odgen, UT. 80pp. | |
| Electronic Resource | http://www.fs.fed.us/rm/pubs_int/int_gtr346.pdf | |
| Intended Use/Purpose | Inventory | |
| Target Resource Type | Perennial streams identifiable on U.S. Geologic Survey 1:24,000 topographic quad maps | |
| Scale/Unit of Assessment | Stream reach of unspecified length that is defined by confluences or changes in classified reach type (i.e. Montgomery and Buffington (1993) valley segments). | |
| Geographic Applicability | Northern Region (R1) and Intermountain Region (R4) of the USFS, which includes all or parts of WA, OR, ID, MT, ND, SD, WY, UT, NV, east-central CA] | |
| General Level of Effort | Three sampling schemes are presented with corresponding levels of effort ranging from Level I (least intensive) to Level III (most intensive). | |
| Assessment Parameters | <p>Channel/Valley Morphology: Stream discharge; classification of stream reach type as A, B, or C (synonymous with Montgomery and Buffington's (1993) valley segments); Rosgen stream classification; channel gradient; valley confinement; bankfull width and depth (optional); percent undercut banks; channel habitat units [aka bed forms] and lengths; wetted channel width, average water depth; average maximum depth of pocket pools; maximum pool depth; pool crest depth; substrate particle size class (est. or pebble count); percent surface fine sediment (<6mm); bank stability (classes); woody debris tally; riparian community type classification.</p> <p>Physical Habitat: Woody debris tally; riparian community type classification.</p> <p>Water Quality: Temperature.</p> <p>Biology: Fish abundance.</p> <p>Other: - -</p> | |
| Resolution | Qualitative (descriptive; categorical), Semi-Quantitative (ordinal scale, rank, etc.), and Quantitative (actual measurement or estimates). | |
| Output | Raw data / data sheets | |
| Reference | N/A (The objective of the method or procedure is not presented in the context of defining the condition of a resource. However, it may be used to identify or establish reference conditions.) | |

| | | |
|---|--|-----------------------|
| Name | R1/R4 (Northern /Intermountain Regions) Fish and Fish Habitat Standard Inventory Procedures Handbook | Catalog No. 17 |
| QA/QC | The Standard Inventory Procedures outlines recommended training procedures for field crews that includes exercises both in the field and in the office entering data. It is also recommended that field crews break every 2 to 3 hours and review field data sheets for missing data, illegible entries, misplaced decimal points, etc. Data forms and equipment lists are provided. | |
| Description/ Summary | <p>Most morphological and physical habitat metrics required in the Standard Inventory Procedures are undertaken using visual estimation methods or selected from standardized lists of types or classes. These metrics are therefore primarily qualitative or semi-quantitative. For example, a detailed, hierarchical channel habitat type classification (aka bed forms) is provided in tabular form, explained in the text, and illustrated with photographs and diagrams. This classification and attendant metrics to further characterize habitat types (e.g. pool depth, pool crest depth, step pool total, etc.) provide the primary focus of the physical and morphological component of the Standard Inventory Procedures. Relative fish abundance by species and size/age class is determined using the direct enumeration snorkeling technique of Thurow (1994), and is the primary quantitative component of the Standard Inventory Procedures.</p> <p>The final 15 to 30 minutes of the field survey should be spent writing a narrative description of the site, including observed land management activities, natural limitations to fish migration, sediment sources and other site observations that might not be captured by field sampling.</p> | |
| Expertise Required | Sample metrics were specifically selected, in part, for the ease with which inexperienced field technicians could be taught the sampling methods, resulting in reasonable expectations for accurate, consistent data. | |
| Time Necessary to Conduct Assessment | Field sampling = 1 day, 2 to 3 persons | |
| Seasonality | Methods are designed for sampling fish and fish habitat at stream baseflow, thus after peak snowmelt. However, caution needs to be taken to avoid sampling streams during spring and summer runs of spawning chinook salmon- a listed endangered species. Where fish surveys will be conducted, sampling should occur in July and August. | |
| Related Procedures/ References | <p>Montgomery, D.R., and J.M. Buffington. 1993. Channel classification, prediction of channel response and assessment of channel condition. Report TFW-SI-110-93-002, Timber/Fish/Wildlife Agreement, Washington, 96 pp.</p> <p>Thurow, R.F. 1994. Underwater methods for study of salmonids in the Intermountain West. Gen. Tech. Rpt. INT-GTR-307, U.S.D.A. Forest Service, Intermountain Research Station, 30 pp.</p> | |
| Other/ Notes | | |

| | | |
|---------------------------------|--|-----------------------|
| Name | Effectiveness monitoring for streams and riparian areas within the Pacific Northwest: stream channel methods for core attributes | Catalog No. 18 |
| Primary Author/ Agency | U.S. Forest Service USFS. 2004. Effectiveness monitoring for streams and riparian areas within the Pacific Northwest: stream channel methods for core attributes. Aquatic and Riparian Effectiveness Monitoring Program (AREMP) & PACFISH/INFISH (PIBO) Effectiveness Monitoring Program, Multi-Federal Agency Monitoring Programs. U.S. Department of Agriculture, Forest Service. Unpublished paper available at: http://www.reo.gov/monitoring/reports/watershed-reports-publications.shtml | |
| Electronic Resource | http://www.reo.gov/monitoring/reports/watershed-reports-publications.shtml | |
| Intended Use/Purpose | Inventory; Ambient Monitoring. | |
| Target Resource Type | Wadeable Streams | |
| Scale/Unit of Assessment | Stream reach, 20X bankfull width based on width classes (minimum 525 feet) | |
| Geographic Applicability | Washington, Oregon, and most of Idaho, as well as western Montana, northeastern Nevada, northwestern Wyoming, and northern California (~interior Columbia River watershed, plus areas west of the Cascade Mountains). | |
| General Level of Effort | Moderate | |
| Assessment Parameters: | Channel/Valley Morphology: Bankfull width; water surface slope; substrate particle size; pool length & residual pool depth. Physical Habitat: Woody debris tally; percent surface fines on pool tails. Water Quality: Conductivity. Biology: Benthic macroinvertebrates. Other: - - | |
| Resolution | Quantitative (actual measurement or estimate) | |
| Output | Raw data | |
| Reference | N/A (The objective of the method or procedure is not presented in the context of defining the condition of a resource. However, it may be used to identify or establish reference conditions.) | |
| QA/QC | Not stated. | |

| | | |
|---|---|-----------------------|
| Name | Effectiveness monitoring for streams and riparian areas within the Pacific Northwest: stream channel methods for core attributes | Catalog No. 18 |
| Description/ Summary | <p>The Aquatic and Riparian Effectiveness Monitoring Program (AREMP) is a multi-federal agency monitoring program to assess the condition of watersheds within the Northwest Forest Plan area (federally managed lands "west of the Cascades"). The primary objective of the PACFISH/INFISH (PIBO) Effectiveness Monitoring Program is to determine whether priority biological and physical attributes, processes, and functions of riparian and aquatic systems are being degraded, maintained, or restored on federally managed lands within the interior Columbia River basin. This document describes the standardized methods that AREMP and PIBO compiled following ten years of use, evaluation, and peer review for a set of core stream channel attributes.</p> <p>The Core Attributes methods utilize transect-based methods for measuring physical habitat and geomorphic metrics. Stepwise instructions are thorough and include illustrative figures for clarification. This is, however, simply a collection of recommended metrics. There is no discussion of data management, QA/QC, data analysis, or any other component typical of a condition assessment procedure. The intent of this document is to simply identify the core metrics shared by the AREMP and PIBO long-term monitoring programs.</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | Not stated. | |
| Related Procedures/ References | <p>Heitke, J.D., E.J. Archer, D.D. Dugaw, B.A. Bouwes, E.A. Archer, R.C. Henderson, and J.L. Kershner. 2008. Effectiveness monitoring for streams and riparian areas: sampling protocol for stream channel attributes. PACFISH/INFISH Biological Opinion (PIBO) Effectiveness Monitoring Program, Multi-federal Agency Monitoring Program; Logan, UT. Unpublished paper on file at: http://www.fs.fed.us/biology/fishecology/emp.</p> <p>Henderson, R.C.; E.K. Archer, B.A. Bouwes, M.S. Coles-Ritchie, and J.L. Kershner. 2005. PACFISH/INFISH Biological Opinion (PIBO): Effectiveness Monitoring Program seven-year status report 1998 through 2004. Gen. Tech. Rep. RMRS-GTR-162. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 16 p.</p> <p>Kershner, J.L., E.K. Archer, M. Coles-Ritchie, E.R. Cowley, R.C. Henderson, K. Kratz, C.M. Quimby, D.L. Turner, L.C. Ulmer, M.R. Vision. 2004. Guide to effective monitoring of aquatic and riparian resources. General Technical Report RMRS-GTR-121. U.S. Department of Agriculture Forest Service, Fort Collins, CO.</p> | |
| Other/Notes | | |

| | | |
|---------------------------------|--|-----------------------|
| Name | A Manual of Procedures for Sampling Surface Waters [Arizona] | Catalog No. 19 |
| Primary Author/ Agency | Arizona Department for Environmental Quality ADEQ. 2005. A Manual of Procedures for Sampling Surface Waters, L. Lawson (ed.), Arizona Department for Environmental Quality, Hydrologic Support and Assessment Section. Phoenix, AZ. | |
| Electronic Resource | http://www.azdeq.gov/environ/water/assessment/download/sampling.pdf | |
| Intended Use/Purpose | Non-Regulatory Condition Assessment; Inventory; Ambient Monitoring. | |
| Target Resource Type | Wadeable Streams | |
| Scale/Unit of Assessment | Stream reach, 20-30X bankfull width or two complete meander lengths (minimum 100 meters "for large streams") | |
| Geographic Applicability | Arizona | |
| General Level of Effort | Intensive (1 day± in the field by a trained or experienced crew of 2 or more persons) | |
| Assessment Parameters: | <p>Channel/Valley Morphology: Stream discharge; stream type classification (Rosgen, 1996); stream type evolutionary stage; longitudinal channel profile; channel cross-section (bankfull cross-sectional area, bankfull width, bankfull depth; floodprone width); bank height ratio (Rosgen, 2001a); bank erodibility hazard index (Rosgen, 1996; 2001b); substrate particle size (pebble count in riffles, pools, and zig-zag & sieve sample); channel habitat units (aka bed forms); near bank stress; channel pattern / planform (sinuosity, belt width, radius of curvature, meander wave length); entrenchment ratio; sediment competence; pool facet slope analysis; Pfankuch channel stability (modified from Pfankuch (1975).</p> <p>Physical Habitat: Linear habitat complexity index (based on run+glide, riffle, and pool lengths); Habitat Assessment Index; Proper Functioning Condition for riparian wetlands (Prichard et al., 1998); riparian percent canopy gaps (densiometer); riparian vegetative community type.</p> <p>Water Quality: Dissolved oxygen; specific conductivity; pH; temperature; turbidity; redox; bacteria.</p> <p>Biology: Benthic macroinvertebrates; diatoms; percent cover of algae & aquatic macrophytes; riparian vegetation percent cover per strata (trees, shrubs, ground cover) (est.); dominant trees per size class.</p> <p>Other: Potential sources of non-point source pollution.</p> | |
| Resolution | Semi-Quantitative (ordinal scale, rank, etc.); Quantitative (actual measurement or estimate). | |
| Output | Condition Assessment; Index (e.g. numeric score); Raw data. | |

| | | |
|---|--|-----------------------|
| Name | A Manual of Procedures for Sampling Surface Waters [Arizona] | Catalog No. 19 |
| Reference | N/A (The objective of the method or procedure is not presented in the context of defining the condition of a resource. However, it may be used to identify or establish reference conditions.) | |
| QA/QC | Not stated. | |
| Description/ Summary | <p>The Arizona Department of Environmental Quality (ADEQ), Manual of Procedures for the Sampling of Surface Waters is an exhaustive collection of very specific methods, protocols, administrative policies, and QA/QC considerations that covers all facets Arizona's surface water sampling program. Section 1 outlines Pre-Trip Administrative Activities, including safety procedures and data forms, while Section 2 describes Equipment Calibration and Cleaning Procedures. Section 3 addresses Field Procedures and is divided into three parts. Part A covers Basic Field Procedures and details activities directly involved in collecting field data for water quality, bacteria, macroinvertebrates and algae. Part B, Geomorphology Procedures, describes activities that assess the physical properties of stream channels. Part C, Habitat Assessments Procedures, describes the methods used to collect and assess habitat and the biological condition of wadeable streams. Section 4 of the Manual addresses Post-Trip Procedures, and Section 5 discusses Data Management. Finally, Section 6 provides Supporting Material as an appendix to the Manual.</p> <p>Biological components of the ADEQ Manual include bacteria, macroinvertebrates, and diatoms. ADEQ has developed benthic IBI's for cold-water streams (above 5,000 feet elevation) and warm-water streams (below 5,000 feet elevation). Macroinvertebrate sampling is to be conducted in perennial streams only. Formulas to calculate IBI's are provided.</p> <p>The Geomorphology Procedures in Part B of Section 3 are based on or derived from Rosgen (1996) and many measures and interpretive ratios are taken directly from various Rosgen publications. Numerous charts, tables, graphs, and illustrations taken from Rosgen training course materials are also provided in the manual, and surveying methods from Harrelson et al. (1994) are referenced and summarized. Most of the geomorphology parameters specified in the ADEQ Manual result in raw quantitative data, although there are numerous commonly used interpretive ratios and indices based on these data.</p> <p>The Habitat Assessment procedures provided in Part C of Section 3 are intended to aid the interpretation of benthic macroinvertebrate bioassessments. There are field data sheets provided in Part C for water chemistry, discharge, field observations about hydrology, biology, and general condition of the stream reach, as well as non-point source observations, the ADEQ Habitat Assessment Index, and riparian community assessment. The Habitat Assessment Index is based on USEPA RBP (Barbour et al., 1999) and USEPA EMAP protocols for rapid habitat assessment (Lazorchak et al., 1998).</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | Macroinvertebrate sampling should occur in baseflow conditions following winter runoff; generally April-May for desert streams and May-June for mountain streams. | |

| | | |
|--------------------------------------|---|-----------------------|
| Name | A Manual of Procedures for Sampling Surface Waters [Arizona] | Catalog No. 19 |
| Related Procedures/References | <p>Barbour, M.T., J. Gerritsen, B.D. Snyder, and J.B. Stribling. 1999. Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition. EPA 841-B-99-002. USEPA Office of Water, Washington, D.C.</p> <p>Harrelson, CC., C.L. Rawlins, and J.P. Potyondy. 1994. Stream Channel Reference Sites: An Illustrated Guide to Field Technique. General Technical Report RM-245, U.S. Forest Service Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.</p> <p>Lazorchak, J.M., A.T Herlihy, and J. Green. 1998. Rapid habitat and Visual Stream Assessments, Section 14 in J.M. Lazorchak et al. (Eds) EMAP- Surface Waters: Field Operations and Methods for Measuring the Ecological Condition of Wadeable Streams. EPA/620/R-94/004F, U.S. Environmental protection Agency, Washington, D.C.</p> <p>Moody, T.O. and W. Odem. 1999. Regional relationships for bankfull stage in natural channels of Central and Southern Arizona. Prepared for the U.S. Forest Service, Albuquerque, NM by Northern Arizona University, Flagstaff, AZ.</p> <p>Pfankuch, D.J. 1975. Stream reach inventory and channel stability evaluation: A watershed management procedure. U.S. Forest Service Northern Region, R1-75-002.</p> <p>Prichard, D., J. Anderson, C. Correll, J. Fogg, K. Gebhardt, R. Krapf, S. Leonard, B. Mitchell, and J. Staats. 1998. Riparian area management: A user guide to assessing Proper Functioning Condition and the supporting science fo lotic areas. Technical Reference 1737-15, BLM/RS/ST-98/001+1737, U.S. Bureau of Land Management, Denver, CO.</p> <p>Rosgen, D.L. 1996. Applied River Morphology. Wildland Hydrology. Pagosa Springs, CO.</p> <p>Rosgen, D.L. 2001a. A stream channel stability assessment methodology, pgs II-18 to II-26 in Proceedings of the Seventh Federal Interagency Sedimentation Conference, March 25-29, 2001, Reno, NV.</p> <p>Rosgen, D.L. 2001b. A practical method of computing streambank erosion rate, pgs II-9 to II-17 in Proceedings of the Seventh Federal Interagency Sedimentation Conference, March 25-29, 2001, Reno, NV.</p> | |
| Other/Notes | Moody and Odem (1999) compiled regional hydraulic curves for Arizona and New Mexico. | |

| | | |
|---------------------------------|--|-----------------------|
| Name | Stream Condition Inventory (SCI) Technical Guide | Catalog No. 20 |
| Primary Author/ Agency | U.S. Forest Service Frazier, J.W., K.B. Roby, J.A. Boberg, K. Kenfield, J.B. Reiner, D.L. Azuma, J.L. Furnish, B.P. Staab. 2005. Stream Condition Inventory (SCI) Technical Guide. USDA Forest Service, Pacific Southwest Region - Ecosystem Conservation Staff. Vallejo, CA. | |
| Electronic Resource | http://www.fs.fed.us/r5/publications/water_resources/sci/techguide-v5-08-2005-a.pdf | |
| Intended Use/Purpose | Inventory; Ambient Monitoring. | |
| Target Resource Type | Wadeable perennial streams with channel gradients $\leq 10\%$. The SCI Technical Guide adds that some SCI methods are applicable to intermittent streams, but others are not. | |
| Scale/Unit of Assessment | Stream reach (recommended minimum length is 500 meters; 100 meter reach is acceptable if neither large woody debris nor pools are key attributes) | |
| Geographic Applicability | California | |
| General Level of Effort | Intensive (1 day \pm in the field by a trained or experienced crew of 2 or more persons) | |
| Assessment Parameters | <p>Channel/Valley Morphology: Channel cross-sectional dimensions; width:depth ratio; entrenchment; water surface gradient; bank stability (percent cover of vegetation, rock, downed wood, or other erosion resistant material); bank angle; substrate particle size distribution; bankfull stage; number and length of channel habitat units [aka bed forms]; residual pool depth; streamshore water depth; pool sediment $\sim V*w$ (optional).</p> <p>Physical Habitat: Woody debris tally; pool tail surface fine sediment; stream shading (solar insolation);</p> <p>Water Quality: Temperature; conductivity; total alkalinity.</p> <p>Biology: Macroinvertebrates; aquatic fauna (herptofauna and fish).</p> <p>Other: - -</p> | |
| Resolution | Quantitative (actual measurement or estimate). | |
| Output | Raw data | |
| Reference | <p>The Technical Guide refers to regional reference streams for which inventories using SCI can provide useful comparison to non-reference conditions. However, the protocol itself does not result in a "condition index" that is based on an internal calibration to these reference streams.</p> <p>However, Appendix A of the Technical Guide presents a brief analysis of SCI data comparing conditions between <i>a priori</i> classification of reference and non-reference streams throughout USFS Region 5.</p> | |

| | | |
|---|--|-----------------------|
| Name | Stream Condition Inventory (SCI) Technical Guide | Catalog No. 20 |
| QA/QC | All crew members must complete both introductory and refresher training sessions that include a combination of classroom and field exercises. All field data is to be checked by the crew leader while still in the field to ensure that all data sheets are legible and complete. Specific QA/QC documentation forms are provided to track QA/QC measures, including training documentation, field survey checklists, field oversight, and data entry. | |
| Description/ Summary | <p>The purpose of the USFS Pacific Southwest Region Stream Condition Inventory (SCI) is to collect intensive and repeatable data from stream reaches to document existing conditions and make reliable comparisons over time within or between stream reaches. It is designed to assess effectiveness of management actions on streams and to document temporal changes in stream conditions of unmanaged watersheds.</p> <p>The protocol stresses quantifiable, objective measurements of 17 core attributes and one optional attribute, but also adds that still additional optional attributes related to specific biota or stream characteristics may be needed to meet local inventory and monitoring objectives. Collecting SCI data in the field is accomplished using a multiple-pass sequence throughout the sample stream reach. The sample protocol provided in the Technical Guide is based on a four-pass sequence, where some of the above referenced attributes are measured and documented during each successive pass. Sample procedures for some specific attributes that could require potentially heavy or cumbersome equipment are described using more simplistic methods to ease transport into remote sample locations. One example includes the use of line levels and stadia rods in lieu of heavy tripods and a total station or automated level for channel surveying. Recommended sequential sample methods are described, including specific task instructions, necessary equipment, and data forms for each pass.</p> | |
| Expertise Required | Not stated, but refer to training requirements in QA/QC above. | |
| Time Necessary to Conduct Assessment | <p>The Technical Guide suggests that up to 2-3 days could be required to initially establish and survey a sample reach, depending on travel time and crew experience. An undefined, but shorter amount of time is necessary to re-sample the same reach.</p> <p>The optional V*w pool sediment attribute is acknowledged to be a very intensive inventory, and is in fact cited as requiring 1-3 days to sample only this attribute, depending on reach length.</p> | |
| Seasonality | Not stated. | |
| Related Procedures/ References | Applicable references are provided for each of the 18 stream inventory attributes described in the Technical Guide. | |
| Other/Notes | | |

| | | |
|---------------------------------|---|---|
| Name | Idaho Small Stream Ecological Assessment Framework | Catalog No. 21 |
| Primary Author/ Agency | Idaho Department of Environmental Quality Grafe, C.S. (ed.). 2002a. Idaho Small Stream Ecological Assessment Framework: An Integrated Approach. Idaho Department of Environmental Quality; Boise, Idaho. | |
| Electronic Resource | http://www.deq.idaho.gov/water/data_reports/surface_water/monitoring/publications.cfm | |
| Intended Use/Purpose | Non-Regulatory Condition Assessment; Inventory; Ambient Monitoring. | |
| Target Resource Type | Wadeable streams (generally <5 th order; wetted width <15 feet at baseflow) | |
| Scale/Unit of Assessment | Not stated (see IDEQ, 2007). | |
| Geographic Applicability | Idaho | |
| General Level of Effort | Moderate | |
| Assessment Parameters: | Channel/Valley Morphology: | Substrate particle size analysis (i.e. number of Wolman size classes); channel shape (undercut). |
| | Physical Habitat: | In-stream cover; woody debris tally; percent fines less than 2mm in wetted stream width; embeddedness; percent bank cover; percent canopy cover; disruptive pressures (qualitative variable used to determine seasonal human impacts on riparian zones); zone of influence (riparian zone width). |
| | Water Quality: | -- |
| | Biology: | Macroinvertebrates; fish. |
| | Other: | -- |
| Resolution | Semi-Quantitative (ordinal scale, rank, etc.); Quantitative (actual measurement or estimate). | |
| Output | Condition Assessment; Index (e.g. numeric score); Raw data. | |
| Reference | Internal. | |
| QA/QC | Not stated (see IDEQ, 2007). | |

| | | |
|---|---|-----------------------|
| Name | Idaho Small Stream Ecological Assessment Framework | Catalog No. 21 |
| Description/ Summary | <p>The Idaho Small Stream Ecological Assessment Framework describes the development and integration of three multimetric indexes that the Idaho Department of Environmental Quality (IDEQ) uses to assess aquatic life use support for small Idaho streams. The indexes were developed based on rapid bioassessment concepts developed by USEPA (Barbour et al. 1999). Specific field sampling protocols are described in IDEQ (2007). IDEQ uses different monitoring and assessment protocols depending on water body size, and has developed a three-parameter index to distinguish “small streams” from “rivers.” These parameters include stream order, average width at base flow, and average depth at base flow. Generally, streams that are less than fifth order, less than 15 feet in average base flow wetted width, and less than an average of 0.4 meters deep at base flow are considered small streams by IDEQ. Grafe (2002b) discusses aquatic life use support protocols for use on Idaho rivers.</p> <p>The Stream Macroinvertebrate Index (SMI) uses nine benthic macroinvertebrate metrics to calculate uniquely referenced index values for each of three different Idaho bioregions (Northern Mountains, Central and Southern Mountains, and Basins). These individual metrics include: total taxa, Ephemeroptera taxa, Plecoptera taxa, Trichoptera taxa, percent Plecoptera, Hilsenhoff Biotic Index, percent five dominant taxa, scraper taxa, and clinger taxa. Jessup and Gerritsen (2002) describe the development of the SMI in detail.</p> <p>The Stream Fish Index (SFI) utilizes two different sets of metrics to characterize water quality condition for montane-forested and desert basin-rangeland classifications. The rangeland metrics include: percent cold water individuals, Jaccard’s community similarity coefficient, percent omnivores and herbivores, percent cyprinids as longnose dace, percent of fish with certain abnormalities (deformities, eroded fins, lesions, and tumors), and catch per unit effort. The metrics in the forested classification are comprised of: number of cold water native species, percent cold water individuals, percent sensitive native individuals, number of sculpin age classes (unless sample is comprised solely of salmonids), number of salmonid age classes, and catch per unit effort. Mebane (2002) describes the development of the SFI in detail.</p> <p>The Stream Habitat Index (SHI) is calibrated to Idaho ecoregions and utilizes ten habitat measures that statistically had the highest correlation with either human disturbance or biological condition. Fore and Bollman (2002) describe the development of the SHI in detail.</p> <p>Each of the above referenced three index scores are adjusted to a common scale using a 1, 2, 3 scoring system, and then averaged to provide a single score representing stream ecological condition.</p> | |
| Expertise Required | Not stated (see IDEQ, 2007). | |
| Time Necessary to Conduct Assessment | Not stated (see IDEQ, 2007). | |
| Seasonality | Not stated (see IDEQ, 2007). | |

| | | |
|--------------------------------------|---|-----------------------|
| Name | Idaho Small Stream Ecological Assessment Framework | Catalog No. 21 |
| Related Procedures/References | <p>Barbour, M.T., J. Gerritsen, B.D. Snyder, and J.B. Stribling. 1999. Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition. EPA 841-B-99-002. USEPA Office of Water, Washington, D.C.</p> <p>Fore, L. and W. Bollman. 2002. Stream habitat index. Chapter 5, In C.S. Grafe (ed). Idaho Small Stream Ecological Assessment Framework: An Integrated Approach. Idaho Department of Environmental Quality. Boise, Idaho.</p> <p>Grafe, C.S. (ed). 2002b. Idaho River Ecological Assessment Framework: An Integrated Approach. Idaho Department of Environmental Quality. Boise, Idaho.</p> <p>IDEQ. 2007. Beneficial Use Reconnaissance Program Field Manual for Streams. Idaho Department of Environmental Quality, Beneficial Use Reconnaissance Program Technical Advisory Committee. Boise, Idaho.</p> <p>Jessup, B. and J. Gerritsen. 2002. Stream macroinvertebrate index. Chapter 3, In C.S. Grafe (ed). Idaho Small Stream Ecological Assessment Framework: An Integrated Approach. Idaho Department of Environmental Quality. Boise, Idaho.</p> <p>Mebane, C.A. 2002. Stream fish index. Chapter 4, In C.S. Grafe (ed). Idaho Small Stream Ecological Assessment Framework: An Integrated Approach. Idaho Department of Environmental Quality. Boise, Idaho.</p> | |
| Other/Notes | | |

| | | |
|---------------------------------|---|-----------------------|
| Name | Idaho River Ecological Assessment Framework | Catalog No. 22 |
| Primary Author/ Agency | Idaho Department of Environmental Quality Grafe, C.S. (ed). 2002b. Idaho River Ecological Assessment Framework: An Integrated Approach. Idaho Department of Environmental Quality. Boise, Idaho. | |
| Electronic Resource | http://www.deq.idaho.gov/water/data_reports/surface_water/monitoring/publications.cfm | |
| Intended Use/Purpose | Non-Regulatory Condition Assessment; Inventory; Ambient Monitoring. | |
| Target Resource Type | Non-wadeable rivers (≥fifth order, >15 feet in average base flow wetted width, and >0.4 meters average depth at base flow) | |
| Scale/Unit of Assessment | Not stated (see IDEQ, 2007). | |
| Geographic Applicability | Idaho | |
| General Level of Effort | Not stated. | |
| Assessment Parameters | Channel/Valley Morphology: -- | |
| | Physical Habitat: -- | |
| | Water Quality: Temperature; dissolved oxygen; biochemical oxygen demand; pH; total solids; ammonia + nitrate nitrogen; total phosphorus; fecal coliform. | |
| | Biology: Macroinvertebrates; fish; diatoms. | |
| | Other: -- | |
| Resolution | Semi-Quantitative (ordinal scale, rank, etc.); Quantitative (actual measurement or estimate). | |
| Output | Condition Assessment; Index (e.g. numeric score); Raw data. | |
| Reference | Internal. | |
| QA/QC | Not stated (see IDEQ, 2007). | |

| | | |
|---|--|-----------------------|
| Name | Idaho River Ecological Assessment Framework | Catalog No. 22 |
| Description/ Summary | <p>The Idaho Department of Environmental Quality (IDEQ) uses biological indicators, physicochemical data and numeric water quality criteria to assess aquatic life use support for rivers. The Idaho River Ecological Assessment Framework describes the development and integration of the River Macroinvertebrate Index (RMI), River Fish Index (RFI), and River Diatom Index (RDI) that IDEQ uses to assess cold water aquatic life use support determinations in Idaho rivers. The River Physicochemical Index (RPI), another interpretive tool, is also discussed.</p> <p>IDEQ uses different monitoring and assessment protocols depending on water body size, and has developed a three-parameter index to distinguish “small streams” from “rivers.” These parameters include stream order, average width at base flow, and average depth at base flow. Generally, streams that are at least fifth order, greater than 15 feet in average base flow wetted width, and greater than an average of 0.4 meters deep at base flow are considered rivers by IDEQ. Grafe (2002a) discusses aquatic life use support protocols for use on small Idaho streams.</p> <p>The River Macroinvertebrate Index (RMI) is a multimetric index consisting of five macroinvertebrate metrics: taxa richness, EPT richness, percent dominance, percent Elmidae (riffle beetles), and percent predators. This macroinvertebrate index is basically a variation of the framework designed for small streams (Jessup and Gerritsen, 2002) and is applicable to Idaho rivers throughout the state. Royer and Mebane (2002) raise some interesting considerations applicable to identifying biological reference conditions for 0 large rivers.</p> <p>The River Fish Index (RFI) is a quantitative fish index applicable to cold water rivers of the interior Columbia River basin (Idaho, Montana, Oregon, Washington, and Wyoming). The index is comprised of the following metrics: number of cold water native species, number of sculpin age classes or percent sculpin (data dependent), percent sensitive native individuals, percent cold water individuals, percent tolerant individuals, number of non-indigenous species, number of selected salmonid age classes, number of cold water individuals per minute of electrofishing, percent carp (if carp introduced), and anomalies. Mebane (2002) describes the RFI in detail.</p> <p>The River Diatom Index (RDI) consists of seven attributes of relative abundance including percent: sensitive to disturbance, very tolerant of disturbance, nitrogen heterotrophs, polysaprobic, requiring high oxygen, very motile, and deformed valves. The RDI also includes two measures of taxon richness: eutrophic and alkaliphilic species. The index significantly correlated with measures of human disturbance at the site and at the level of the catchment. Fore and Grafe (2002) describe the RDI in detail.</p> <p>The River Physicochemical Index (RPI) is based on the Oregon Water Quality Index (Cude, 1998; 2001). This index has been tested and used extensively in Oregon to assess water quality conditions. The RPI consists of eight water quality parameters: Sub-index scores for each variable are calculated using complex regressions for data that falls within a set range for each of the variables and threshold scores for data outside of that range (Cude, 1998). The individual sub-indexes are then averaged to give a single index value. Brandt (2002) describes the applicability of the Oregon Water Quality Index to Idaho rivers.</p> <p>IDEQ integrates the RMI, RDI, and RFI index scores using a rating and averaging approach. Index scores are adjusted to a common scale using a 1, 2, 3 scoring system. The converted scores are then averaged to provide a single score. The RPI is not integrated in the averaging process, but may provide additional information in interpreting physicochemical data.</p> | |
| Expertise Required | Not stated (see IDEQ, 2007). | |
| Time Necessary to Conduct Assessment | Not stated (see IDEQ, 2007). | |
| Seasonality | Not stated (see IDEQ, 2007). | |

| | | |
|--------------------------------------|--|-----------------------|
| Name | Idaho River Ecological Assessment Framework | Catalog No. 22 |
| Related Procedures/References | <p>Brandt, D. 2002. River physiochemical index. Chapter 6, In C.S. Grafe (ed.) Idaho River Ecological Assessment Framework: An Integrated Approach. Idaho Department of Environmental Quality; Boise, Idaho.</p> <p>Cude, C.G. 1998. Oregon water quality index: a tool for evaluating water quality management effectiveness. Oregon Department of Environmental Quality, Laboratory Division, Water Quality Monitoring Section. Portland, OR. 20 pp.</p> <p>Cude, C.G. 2001. Oregon water quality index: a tool for evaluating water quality management effectiveness. Journal of the American Water Resources Association 37(1):125-137.</p> <p>Fore, L.S. and C.S. Grafe. 2002. River diatom index. Chapter 5, In C.S. Grafe (ed.) Idaho River Ecological Assessment Framework: An Integrated Approach. Idaho Department of Environmental Quality; Boise, Idaho.</p> <p>Grafe, C.S. (ed.). 2002a. Idaho Small Stream Ecological Assessment Framework: An Integrated Approach. Idaho Department of Environmental Quality; Boise, Idaho.</p> <p>Jessup, B. and J. Gerritsen. 2002. Stream macroinvertebrate index. Chapter 3, In C.S. Grafe (ed.) Idaho Small Stream Ecological Assessment Framework: An Integrated Approach. Idaho Department of Environmental Quality. Boise, Idaho.</p> <p>Mebane, C.A. 2002. River fish index. Chapter 4, In C.S. Grafe (ed.) Idaho River Ecological Assessment Framework: An Integrated Approach. Idaho Department of Environmental Quality; Boise, Idaho.</p> <p>Royer, T.V. and C.A. Mebane. 2002. River macroinvertebrates index. Chapter 3, In C.S. Grafe (ed.) Idaho River Ecological Assessment Framework: An Integrated Approach. Idaho Department of Environmental Quality; Boise, Idaho.</p> | |
| Other/Notes | | |

| | | |
|---------------------------------|---|-----------------------|
| Name | Beneficial Use Reconnaissance Program Field Manual for Streams | Catalog No. 23 |
| Primary Author/ Agency | Idaho Department of Environmental Quality IDEQ. 2007. Beneficial Use Reconnaissance Program Field Manual for Streams. Idaho Department of Environmental Quality, Beneficial Use Reconnaissance Program Technical Advisory Committee. Boise, Idaho. | |
| Electronic Resource | http://www.deq.idaho.gov/water/data_reports/surface_water/monitoring/overview.cfm#beneficial | |
| Intended Use/Purpose | Non-Regulatory Condition Assessment; Inventory; Ambient Monitoring. | |
| Target Resource Type | Wadeable streams | |
| Scale/Unit of Assessment | Stream reach, 30X bankfull width (minimum 100 meters) | |
| Geographic Applicability | Idaho | |
| General Level of Effort | Moderate; Intensive (1 day± in the field by a trained or experienced crew of 2 or more persons) | |
| Assessment Parameters: | <p>Channel/Valley Morphology: Stream discharge; width/depth ratio (wetted and bankfull dimensions); entrenchment ratio; sinuosity; channel habitat units [aka bed forms]; elevation; channel gradient; bank angle; bank undercut distance; substrate particle size distribution (pebble counts); Rosgen channel classification.</p> <p>Physical Habitat: Woody debris tally; shade/canopy cover (densiometer); bank cover and stability (percent cover of vegetation, rock, downed wood, or other erosion resistant material); Pool Quality Index (pool length, substrate, overhead cover, submerged cover, percentage of undercut banks, maximum pool depth, maximum pool width, and depth at pool tail out); rapid habitat assessment (modified from Hayslip, 1993).</p> <p>Water Quality: Temperature; specific conductivity; bacteria (E. coli).</p> <p>Biology: Macroinvertebrate assemblages; periphyton assemblages; fish assemblages; amphibians.</p> <p>Other: Stream order.</p> | |
| Resolution | Semi-Quantitative (ordinal scale, rank, etc.); Quantitative (actual measurement or estimate). | |
| Output | Raw data (Grafe et al. (2002) describe data analysis and interpretation of BURP data.) | |

| | | |
|---|--|-----------------------|
| Name | Beneficial Use Reconnaissance Program Field Manual for Streams | Catalog No. 23 |
| Reference | <p>N/A (The objective of the BURP Field Manual itself does not address reference conditions per se.)</p> <p>Grafe (2002a; 2002b) describes the development and integration of various condition indexes that IDEQ uses to assess aquatic life use support for Idaho streams and rivers, and these indexes have been developed and calibrated based on internal reference data from either Idaho ecoregions or bioregions, as applicable.</p> | |
| QA/QC | <p>IDEQ ensures quality BURP data by providing centralized training for BURP crews, annual BURP Coordinator workshops, strict adherence to the Field Manual, consistent crew supervision, compilation and adherence to annual work plans, conducting comprehensive annual field audits, and following a quality assurance plan that addresses such issues as data handling, voucher specimens, and equipment calibration.</p> | |
| Description/ Summary | <p>The Idaho Beneficial Use Reconnaissance Program (BURP) conducts stream monitoring activities to support assessments of biological assemblages and physical habitat structure, which in turn supports characterization of individual stream integrity and the total quality of Idaho's waters. The BURP Field Manual is presented consistent with the four phases of BURP field activities: (1) Planning; (2) Preparing for field activities; (3) Field sampling, including detailed protocol descriptions; and (4) Follow-up and reporting.</p> <p>The field sampling protocols, which are generally transect based, are presented in a recommended sequence for performing monitoring activities.</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | June to September | |
| Related Procedures/ References | <p>Grafe, C.S. (ed.). 2002a. Idaho Small Stream Ecological Assessment Framework: An Integrated Approach. Idaho Department of Environmental Quality; Boise, Idaho.</p> <p>Grafe, C.S. (ed.). 2002b. Idaho River Ecological Assessment Framework: An Integrated Approach. Idaho Department of Environmental Quality; Boise, Idaho.</p> <p>Grafe, C. S., M. McIntyre, C. Mebane and D. Mosier. 2002. Water Body Assessment Guidance (Second Edition). Idaho Department of Environmental Quality. Boise, ID.</p> <p>Hayslip, G.A. (ed.). 1993. Region 10 in-stream biological monitoring handbook for wadeable streams in the Pacific Northwest. EPA 910/9-92-013, U.S. Environmental Protection Agency, Region 10. Seattle, WA.</p> | |
| Other/Notes | | |

| | | |
|---------------------------------|---|-----------------------|
| Name | Aquatic Inventories Project Methods for Stream Habitat Surveys | Catalog No. 24 |
| Primary Author/ Agency | Oregon Department of Fish and Wildlife Moore, K., K. Jones, J. Dambacher, C. Stein, et al. 2008. Aquatic Inventories Project: Methods for Stream Habitat Surveys, Version 17.1, May 2008. Oregon Department of Fish and Wildlife, Aquatic Inventories Project, Conservation and Recovery Program, Corvallis, OR. | |
| Electronic Resource | http://www.science.oregonstate.edu/~madsen/TIESNA2009/Habitat_protocol.pdf | |
| Intended Use/Purpose | Inventory; Ambient Monitoring. | |
| Target Resource Type | Streams (No further clarification provided. However, there are procedural references specific to dry channels, suggesting that intermittent streams may also be inventoried using these methods). | |
| Scale/Unit of Assessment | Stream reach of unspecified length that is defined based on confluences with named tributaries, changes in valley and channel form, major changes in vegetation type, or changes in land use or ownership. Appendices suggest that the sample stream reach should be 1000 meters. | |
| Geographic Applicability | Oregon | |
| General Level of Effort | Moderate; Intensive (1 day± in the field by a trained or experienced crew of 2 or more persons) | |
| Assessment Parameters | <p>Channel/Valley Morphology: Stream discharge; water surface gradient; length, wetted width, and sub-type of each channel habitat unit [aka bed forms]; maximum pool depth; pool crest depth; substrate particle size classes (est.); boulder count (greater than 0.5 m average diameter located within or at margins of bankfull channel); percent active eroding banks (est.); percent undercut banks (est.); elevation; categorical valley type based on valley width index (ratio of the active channel width to the valley width); bankfull width; channel height above bankfull depth; floodprone width; terrace height (height from the streambed to the top of the first terrace above the floodprone height); terrace width; riparian zone gradient.</p> <p>Physical Habitat: Woody debris tally; channel shade (via clinometer); general riparian community structure (size class and type).</p> <p>Water Quality: Temperature.</p> <p>Biology: Fish; amphibians; riparian vegetation (belt transect 5m x 30m perpendicular to each side of the stream): percent-cover trees (est.), percent-cover shrubs (est.), percent cover herbaceous layer (est.); tree count (stem density) per size class.</p> <p>Other: Stream order; drainage density; watershed area; watershed land use.</p> | |
| Resolution | Semi-Quantitative (ordinal scale, rank, etc.); Quantitative (actual measurement or estimate). | |
| Output | Raw data. | |
| Reference | N/A (The objective of the method or procedure is not presented in the context of defining the condition of a resource. However, it may be used to identify or establish reference conditions.) | |
| QA/QC | Not stated. | |

| | | | |
|---|---|--|-----------------------|
| Name | Aquatic Inventories Project Methods for Stream Habitat Surveys | | Catalog No. 24 |
| Description/ Summary | <p>The Aquatic Inventories Project is designed to provide quantitative information on habitat condition for streams throughout Oregon. The Methods for Stream Habitat Surveys systematically identifies and quantifies valley and stream geomorphic features, resulting in a matrix of measurements and spatial relationships that can be generalized into frequently occurring valley and channel types.</p> <p>The Methods procedure requires completion of five (5) data sheets: 1) Stream Reach, 2) Unit-1, 3) Unit-2, 4) Wood, and 5) Riparian. Most channel morphology and physical habitat parameters are measured or estimated at either every channel habitat unit or every nth channel habitat unit, where $n \leq 10$. Channel habitat units (aka bed forms) are themselves classified in the field according to defined sub-types that share relatively homogeneous bed form, flow characteristics, and water surface slope. For example, six sub-types of pools are defined in the Methods.</p> <p>Data forms and instructions/guidelines for estimating or measuring each parameter are provided.</p> | | |
| Expertise Required | Field work consistent with the Methods for Stream Habitat Surveys is intended to be carried out by a crew of two persons. | | |
| Time Necessary to Conduct Assessment | Not stated. | | |
| Seasonality | Not stated. | | |
| Related Procedures/References | | | |
| Other/Notes | | | |

| | | |
|---------------------------------|--|-----------------------|
| Name | Stream Inventory Handbook: Level I & II | Catalog No. 25 |
| Primary Author/ Agency | U.S. Forest Service USFS. 2009. Stream Inventory Handbook: Level I & II, Version 2.9. U.S. Forest Service, Pacific Northwest Region, Region 6. | |
| Electronic Resource | http://www.fs.fed.us/r6/water/fhr/sida/handbook/Stream-Inv-2009.pdf | |
| Intended Use/Purpose | Inventory; Ambient Monitoring | |
| Target Resource Type | Wadeable streams (ephemeral, intermittent, or perennial) | |
| Scale/Unit of Assessment | Watershed; and/or Stream reach: A reach is a relatively homogeneous section of stream containing attributes of common character. The recommended minimum length for all reaches is 0.5 miles. All riffles (fast water) must be treated as "measured riffles" in any reach less than 0.5-mile long. | |
| Geographic Applicability | Oregon and Washington | |
| General Level of Effort | Intensive (1 day± in the field by a trained or experienced crew of 2 or more persons) | |
| Assessment Parameters | <p>Channel/Valley Morphology: Stream discharge; length, wetted width, maximum depth, and average depth of each channel habitat unit [aka bed forms]; pool crest depth; pool forming feature (opt.); Rosgen stream type; valley form (opt.); thalweg length (longitudinal profile); bankfull width; average and maximum bankfull depth; floodprone area width; bank stability; substrate particle size classes (est.); particle size distribution (pebble count); mapped valley width; mapped channel length; mapped valley length; mapped channel gradient; measured channel gradient (opt.); mapped sinuosity; elevation (min/max).</p> <p>Physical Habitat: Woody debris tally; inner riparian zone width (average width along both banks from bankfull to a distinct change in vegetation); successional class of riparian vegetation (based on vegetative type and size class); dominant overstory & understory riparian species.</p> <p>Water Quality: Long-term thermograph (mid-June to late September).</p> <p>Biology: Fish; amphibians.</p> <p>Other: Stream order (opt.); watershed area.</p> | |
| Resolution | Quantitative (actual measurement or estimate) | |
| Output | Raw data | |
| Reference | N/A (The objective of the method or procedure is not presented in the context of defining the condition of a resource. However, it may be used to identify or establish reference conditions.) | |

| | | |
|---|---|-----------------------|
| Name | Stream Inventory Handbook: Level I & II | Catalog No. 25 |
| QA/QC | <p>QA/QC requirements are detailed and extensive for each of four phases of implementation of a monitoring program using the USFS Region 6 Stream Inventory Handbook: (1) Program Administration, (2) Pre-Inventory Training, (3) Field Inventory Training; and (4) Post-Inventory Training. There is both regional and national forest-level training required that includes the Handbook protocols themselves, map and aerial photograph interpretation, equipment use and maintenance, taxonomic identification of fish and amphibians, data management, data entry, data analysis, and report writing. Each national forest must also establish a “test reach” for forest-level training.</p> | |
| Description/ Summary | <p>The USFS Region 6 Stream Inventory Handbook: Level I & II is designed on a hierarchical scale. Level I is the basic in-office procedure which identifies standard attributes of the watershed/stream to be analyzed. Level II is an extensive stream channel, riparian vegetation, aquatic habitat condition and biotic inventory conducted on a watershed scale. The Level II inventory includes both requisite core attributes that are necessary to evaluate the condition of the stream and optional attributes. It has been reviewed and is compatible with similar aquatic inventories developed by state agencies, specifically the Oregon Department of Fish and Wildlife (ODFW) and Timber, Fish and Wildlife (TFW) in Washington State. It has been developed as the aquatic companion to the USFS Integrated Resource Inventory, and is comparable with other USFS stream inventories developed in Regions 1, 4, and 5. It contains the “Core Data Standards” developed by an interagency team for implementation of the Northwest Forest Plan.</p> <p>There are two (2) forms to be completed during the Level I in-office inventory and seven (7) to be completed in the field during Level II inventory. Existing information about the stream and watershed to be inventoried is compiled in Level I including existing maps, historic land use and/or aerial photographs, remote sensing data, and previous inventories and/or investigations. Preliminary study stream reaches are also identified in Level I based on changes in mapped valley width, mapped channel gradient, mapped sinuosity, or streamflow inferred by the confluence of large tributaries.</p> <p>All Level II inventory parameters, except stream discharge and particle size distribution, are measured in at least ten (10) pools (scour, plunge, & dam) and ten (10) fast water riffles (turbulent & non-turbulent) in the reach. Channel habitat unit lengths must be measured in every habitat unit throughout the sample reach. The Handbook provides very detailed instructions for measuring each parameter and includes detailed field data sheets.</p> | |
| Expertise Required | See QA/QC above. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | Minimum baseflow conditions. | |
| Related Procedures/ References | | |
| Other/Notes | | |

| | | |
|---------------------------------|--|-----------------------|
| Name | Functional Assessment Approach for High Gradient Streams: West Virginia | Catalog No. 26 |
| Primary Author/ Agency | U.S. Army Corps of Engineers, Huntington District USACE Huntington District. 2007. Functional Assessment Approach for High Gradient Streams: West Virginia. June 2007, U.S. Army Corps of Engineers, Huntington District, Huntington, WV. | |
| Electronic Resource | http://www.lrh.usace.army.mil/permits/ | |
| Intended Use/Purpose | Regulatory Assessment (Clean Water Act, Section 404); Compensatory Mitigation Protocol | |
| Target Resource Type | Headwater Streams: Ephemeral, Intermittent, & Low-order Perennial Characterized by high gradient (channel slope ranges from 4% to 10%), low sinuosity, with common to many step pools (Rosgen type A, Aa, or Aa+ streams) | |
| Scale/Unit of Assessment | Stream reach of unspecified length. | |
| Geographic Applicability | West Virginia | |
| General Level of Effort | Easy (rapid) | |
| Assessment Parameters | <p>Channel/Valley Morphology: Watershed gradient; categorical channel alteration; channel gradient & number of step pools; substrate particle size (est.).</p> <p>Physical Habitat: Woody debris tally.</p> <p>Water Quality: - -</p> <p>Biology: Percent-cover trees; percent-cover shrubs; percent-cover herbaceous layer; Number of native species in upper-most vegetative strata.</p> <p>Other: Watershed land use/ land cover (est.); percent-cover soil detritus.</p> | |
| Resolution | Qualitative (descriptive; categorical); Semi-Quantitative (ordinal scale, rank, etc.). | |
| Output | Condition Assessment; Index (e.g. numeric score); Programmatic or Regulatory Support Information. | |
| Reference | Internal, but based on "field observations, professional judgment, published literature," and similar assessment indicators from other regions and ecosystems | |
| QA/QC | Not stated. However, the documentation indicates that no field studies have been conducted specifically to calibrate the metrics or indicators used in the Approach. | |

| | | |
|---|---|-----------------------|
| Name | Functional Assessment Approach for High Gradient Streams: West Virginia | Catalog No. 26 |
| Description/ Summary | <p>The “Functional Assessment Approach for High Gradient Streams: West Virginia” is considered by the USACE, Huntington District to be an interim approach that involves a visual evaluation of the physical and biological structure of the assessment site. The assessment itself uses a set of eleven (11) metrics that are scored based on ordinal or categorical descriptions and then aggregated in model equations to represent indicators of four (4) defined functions: hydrology, biogeochemical cycling, plant community functions, and wildlife habitat. Each function is described in the documentation, and rationale for including the subset of metrics used to generate an indicator score for each function, scaled from zero to 1.0, is also provided.</p> <p>The Approach documentation specifies that “decisions about how to use the numbers [output] are a matter of policy,” and are not specified in the document.</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | Not stated. | |
| Related Procedures/ References | None. | |
| Other/Notes | <p>The organizational presentation of the Approach document and the structure of the specific model equations that represent each function are very similar to those commonly used in regional guidebooks for hydrogeomorphic (HGM) functional assessment of wetlands.</p> <p>In early 2010, the IFAA was reportedly in the process of being significantly revised by the USACE Engineer Research and Development Center in Vicksburg, Mississippi.</p> | |

| | | |
|---------------------------------|--|-----------------------|
| Name | West Virginia Stream and Wetland Valuation Metric | Catalog No. 27 |
| Primary Author/ Agency | West Virginia Interagency Review Team West Virginia Interagency Review Team. 2010. West Virginia Stream and Wetland Valuation Metric, Version 1.1. March 2010. USACE Huntington District, USACE Pittsburgh District, USEPA, USFWS, USDA NRCS, West Virginia Department of Environmental Protection, and West Virginia Division of Natural Resources. | |
| Electronic Resource | http://www.lrh.usace.army.mil/permits/ | |
| Intended Use/Purpose | Regulatory Assessment (Clean Water Act, Section 404); Compensatory Mitigation Protocol. | |
| Target Resource Type | Wadeable Streams: Ephemeral, Intermittent, or Perennial | |
| Scale/Unit of Assessment | Not stated. However, the benthic macroinvertebrate sampling protocol upon which the West Virginia Stream Condition Index is based utilizes a sample stream reach of 100 meters. | |
| Geographic Applicability | West Virginia | |
| General Level of Effort | Moderate | |
| Assessment Parameters | <p>Channel/Valley Morphology: Channel alteration (H, L)¹; frequency of riffles or bends (H); sinuosity (L); bank stability (H, L); pool substrate characterization (L); Velocity/depth combinations (H); pool variability (L).</p> <p>Physical Habitat: Epifaunal substrate/available cover (H, L); embeddedness (H); sediment deposition (H, L); channel flow status (H, L); bank vegetative protection (H, L); riparian zone width (H, L).</p> <p>Water Quality: Specific conductivity; pH; dissolved oxygen.</p> <p>Biology: Benthic macroinvertebrates.</p> <p>Other: --</p> <p>¹ All Channel/Valley Morphology and Physical Habitat parameters listed above are included as part of the USEPA RBP stream habitat assessment index. H = applicable in high gradient streams; L = applicable in low gradient streams.</p> | |
| Resolution | Semi-Quantitative (ordinal scale, rank, etc.); Quantitative (actual measurement or estimate). | |
| Output | Condition Assessment; Index (e.g. numeric score); Programmatic or Regulatory Support Information. | |
| Reference | Internal (e.g. Index calibrated to existing local or regional reference data). | |
| QA/QC | Not stated. | |

| | | |
|---|---|-----------------------|
| Name | West Virginia Stream and Wetland Valuation Metric | Catalog No. 27 |
| Description/ Summary | <p>The West Virginia Stream and Wetland Metric Valuation (SWMV) was developed to provide regulatory agencies in West Virginia with an assessment method suitable to consistently evaluate proposed impacts to jurisdictional streams and wetlands considering all forms of compensatory mitigation, including mitigation banks, in-lieu fee programs, and permittee responsible mitigation. Only the stream component of SWMV will be addressed here.</p> <p>The SWMV synthesizes correlations derived from multiple established individual assessment methodologies, including the stream habitat assessment component of the USEPA Rapid Bioassessment Protocols (Barbour et al., 1999), the West Virginia Stream Condition Index (Barbour et al., 2000), and a water quality data sheet utilized by the West Virginia Department of Environmental Protection. The SWMV utilizes these data to generate an index ranging from 0 to 1.0 to represent the physical, chemical, and biological integrity of the stream being assessed.</p> <p>The RBP stream habitat assessment is a visual-based rapid assessment that relies upon visual characterizations of ten stream features in order to categorize the quality of those features as either poor, marginal, suboptimal, or optimal. The range of quality from poor to optimal is further defined on a point scale from 0 to 20 for each stream habitat parameter assessed. A few stream habitat parameters used in the assessment vary based on whether the stream has a high gradient and therefore dominated by riffle/run habitat types and coarse substrate, or a low gradient dominated by glide/pool habitats and typically finer substrates. The water quality parameters of concern in the SWMV include pH, specific conductivity, and dissolved oxygen. Both the physical habitat assessment and the water quality data are required for ephemeral, intermittent, or perennial streams. The West Virginia Stream Condition Index (WVSCI) is based on six (6) biological metrics calculated from benthic macroinvertebrate assemblages collected using the WVDEP Save Our Streams protocol (WVDEP, 2010), and is to be included only on intermittent or perennial streams.</p> <p>The SWMV includes a Microsoft Excel spreadsheet that automates the calculation of both sub-indices for each of the indicators (physical, chemical, and biological), as well as the overall condition index. The user enters data for each indicator in the designated portion of the spreadsheet, including the 10 individual parameter scores of the RBP stream habitat assessment (physical indicators), measured water quality data for pH, specific conductivity, and dissolved oxygen (chemical indicators), and the WVSCI index score (biological indicator). The spreadsheet aggregates the subindices into an overall condition index ranging from 0 (poor condition) to 1.0 (best condition). All calculations are internal to the spreadsheet, and cannot be modified by the user.</p> <p>Data may be entered not only for the stream proposed to be impacted, but also for the proposed mitigation site. Additionally, inferences may be drawn to anticipate conditions in the mitigation stream five-years from the date of mitigation. The difference in index score between existing conditions at the mitigation site and anticipated conditions forms the basis upon which determinations of the necessary mitigation stream length may be drawn. There are also considerations built into the spreadsheet to account for anticipated temporal loss of ecosystem functions (i.e. time to maturity of a mitigation site).</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | <p>Not stated.</p> <p>Barbour et al. (2000) found no distinct differentiation based seasonality of data used to develop the WVSCI, which was collected from May to September between 1996 and 1997. However, the authors opined that narrowing the collection period to a range from late spring to early summer would reduce variability and thereby improve the assessments.</p> | |

| | | |
|--------------------------------------|--|-----------------------|
| Name | West Virginia Stream and Wetland Valuation Metric | Catalog No. 27 |
| Related Procedures/References | <p>Barbour, M.T., J. Gerritsen, B.D. Snyder, and J.B. Stribling. 1999. Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition. EPA 841-B-99-002. USEPA Office of Water, Washington, D.C.</p> <p>Barbour, M.T., J. Burton, and J. Gerritsen. 2000. A Stream Condition Index for West Virginia Wadeable Streams. March 28, 2000 (revised July 21, 2000). EPA 68-C7-0014. U.S. Environmental Protection Agency, Region 3 Environmental Services Division and U.S. Environmental Protection Agency, Office of Science and Technology, Office of Water.</p> <p>WVDEP. 2010. West Virginia Save Our Streams. West Virginia Department of Environmental Protection, Division of Water and Waste Management, http://www.wvdep.org/item.cfm?ssid=11&ss1id=202.</p> | |
| Other/Notes | <p>The stream portion of the Stream and Wetland Valuation Metric is anticipated to be superceded by completion of a Comprehensive Stream Assessment Methodology being developed by the USACE Engineer Research and Development Center.</p> | |

| | | |
|---------------------------------|---|--|
| Name | Unified Stream Methodology | Catalog No. 28 |
| Primary Author/ Agency | U.S. Army Corps of Engineers, Norfolk District and Virginia Department of Environmental Quality, January 2007 | |
| Electronic Resource | http://www.nao.usace.army.mil/technical%20services/Regulatory%20branch/USM.asp | |
| Intended Use/Purpose | Regulatory Assessment (Clean Water Act, Section 404; Virginia Water Protection Permit Program); Compensatory Mitigation Protocol. | |
| Target Resource Type | Wadeable streams: Ephemeral, Intermittent, or Perennial. | |
| Scale/Unit of Assessment | Stream reach defined by changes in channel condition, riparian buffer, in-stream habitat, or channel alteration. | |
| Geographic Applicability | Virginia | |
| General Level of Effort | Easy | |
| Assessment Parameters | <u>Wadeable perennial or intermittent streams - Reach Condition Index (based on visual observation):</u> | |
| | Channel / Valley Morphology: | Channel condition (cross-sectional channel stability; preponderance of sediment deposition; vegetative bank coverage; bank erosion); Channel alteration (preponderance of anthropogenic channel disturbance, such as channelization, rip-rap, road crossings, etc.). |
| | Physical Habitat: | Riparian buffers (canopy coverage; number of well represented vegetative strata); in-stream habitat (percent coverage of in-stream habitat, including substrate size variation, flow velocity and depth, woody and leafy debris, undercut banks, etc.). |
| | Water Quality: | -- |
| | Biology: | -- |
| | Other: | -- |
| Resolution | Qualitative (descriptive); Semi-Quantitative (ordinal scale, rank, etc.) | |
| Output | Subjective Index (e.g. numeric score); Qualitative Description; Programmatic or Regulatory Support Information | |
| Reference | Measured External Reference Required (e.g. site specific / project specific reference). | |
| QA/QC | Not stated. | |

| | | |
|---|--|-----------------------|
| Name | Unified Stream Methodology | Catalog No. 28 |
| Description/ Summary | <p>The Unified Stream Methodology (USM) provides a rapid method to assess stream compensatory mitigation requirements for proposed projects seeking authorization to impact jurisdictional streams, as well as the number of credits generated by proposed mitigation projects. The first step in USM is to define the exiting condition of the proposed project stream by calculating a Reach Condition Index (RCI). The RCI is based on condition indices of four factors, each of which is scored according to categorical or ordinal descriptions provided: (1) Channel condition (based on channel evolutionary stage; morphological response following perturbation); (2) Riparian buffer (weighted average percent cover of various vegetative cover types within 100 feet of stream reach); (3) In-stream habitat (relative quantity and variety of natural physical structures in the stream that provide habitat for aquatic organisms); and (4) Channel alteration (direct impacts to the stream as a result of anthropogenic activities). Descriptions provided in the USM of each parameter and condition class thereof are augmented with color photographs representing each condition class.</p> <p>Scoring of the Channel condition factor of the RCI is weighted 2X any other single factor to reflect the importance of physical stability on overall channel condition. Scores for each of the above referenced four factors are summed and then divided by five (5) to obtain the RCI. The RCI is then multiplied by a categorical Impact Factor (IF) that increases with the perceived severity of stream impact type, and the linear length of stream impact in order to determine the compensation requirements necessary to offset proposed impacts.</p> <p>The number of mitigation credits allocated to proposed mitigation measures is based on categorical descriptions of mitigation activities described in the USM. Restoration measures are defined consistent with Rosgen (1997), and receive the greatest mitigation credit per unit stream length. Stream enhancement activities and riparian buffer improvements are likewise described and allocated corresponding credits. Additional "adjustment factors" can be used to further augment mitigation credit generation if certain "exceptional or site specific circumstances" warrant. These include the presence of or benefits to rare, threatened or endangered species or their habitats; livestock exclusion fencing; and watershed preservation.</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | Not stated. | |
| Related Procedures/References | Rosgen, D.L. 1997. A geomorphological approach to restoration of incised rivers. Pgs 12-22 in S.S.Y. Wang, E.J. Langendoen and F.D. Shields, Jr. (eds.), Proceedings of the Conference on Management of Landscapes Disturbed by Channel Incision., University of Mississippi, Oxford, MS. | |
| Other/Notes | | |

| | | |
|---------------------------------|--|-----------------------|
| Name | USACE Charleston District, Standard Operating Procedure: Compensatory Mitigation | Catalog No. 29 |
| Primary Author/ Agency | U.S. Army Corps of Engineers, Charleston District USACE, Charleston District. 2002. Standard Operating Procedure: Compensatory Mitigation. RD-SOP-02-01, September 19, 2002. U.S. Army Corps of Engineers, Charleston District, Charleston, SC. [NOTE: the Charleston SOP is currently being updated, as of February 2010]. | |
| Electronic Resource | http://www.sac.usace.army.mil/?action=mitigation.home | |
| Intended Use/Purpose | Regulatory Assessment (Clean Water Act, Section 404); Compensatory Mitigation Protocol | |
| Target Resource Type | Intermittent Streams; Perennial Streams; and Riparian Zones | |
| Scale/Unit of Assessment | Stream reach of unspecified length | |
| Geographic Applicability | South Carolina | |
| General Level of Effort | Varies; The SOP refers to other guidance for assessment and monitoring methods. | |
| Assessment Parameters | <p><u>Varies, but may include:</u></p> <p>Channel/Valley Morphology: Stream discharge; channel cross-sections & longitudinal profiles [dimension, pattern and profile]; measures of channel and streambank stability (methods undefined); substrate and sediment characteristics (undefined).</p> <p>Physical Habitat: - -</p> <p>Water Quality: Temperature; dissolved oxygen; turbidity.</p> <p>Biology: As applicable: Fish; benthic macroinvertebrates; riparian vegetation.</p> <p>Other: - -</p> | |
| Resolution | Qualitative (descriptive; categorical); Semi-Quantitative (ordinal scale, rank, etc.); and/or Quantitative (actual measurement or estimate). | |
| Output | Programmatic or Regulatory Support Information | |
| Reference | <p>Measured External Reference Required (e.g. site specific / project specific reference)</p> <p>Reference is not necessarily required to place the project stream into regional context based on physical or biological condition, but rather to suggest specific design and/or success criteria for proposed mitigation projects.</p> | |
| QA/QC | Not stated. | |

| | | |
|---|---|-----------------------|
| Name | USACE Charleston District, Standard Operating Procedure: Compensatory Mitigation | Catalog No. 29 |
| Description/ Summary | <p>The Charleston SOP provides a basic written framework to improve predictability and consistency in the development, review, and approval of compensatory mitigation plans submitted as part of the CWA 404 regulatory program within the USACE Charleston District. While the SOP does not provide stream restoration design criteria, it repeatedly references Rosgen methods (Rosgen, 1996) and allocates mitigation credits based in part on the “priority level” of restoration as described in Rosgen (1996). The SOP refers to the use of an external reference site from which design criteria and success standards may be drawn, and refers to Rosgen (1996), the Federal Stream Restoration Working Group (1998), NRCS (1996), and the North Carolina Stream Restoration Institute at North Carolina State University for stream restoration methods and tools. The Charleston SOP also refers to Harrelson et al. (1994) for appropriate stream surveying procedures.</p> <p>Proposed stream mitigation plans must include, among other programmatic elements, surveys of baseline conditions and post-construction conditions; measurable and quantifiable success criteria; and a monitoring plan (5-year minimum) that encompasses both physical and biological metrics. The SOP refers to Rosgen (1996) and the Federal Stream Restoration Working Group (1998) for specific stream monitoring methods.</p> <p>The Charleston SOP states that the goal of compensatory mitigation shall be the restoration and maintenance of the chemical, physical, and biological integrity of the Nation's waters by replacing unavoidably lost wetland or stream functions as close as possible to the impact site. However, the SOP is mostly an administrative tool for allocating mitigation credits and outlining programmatic requirements for mitigation projects. It utilizes a set of matrices to determine the number of mitigation credits necessary to compensate for proposed adverse impacts to aquatic resources, and a second set of similar matrices to estimate the number of mitigation credits generated by a proposed mitigation plan. Each matrix includes a number of factors that are scored independently and then summed to reach a per unit mitigation credit lost or gained. This per unit value is then multiplied by the linear length of stream either impacted or restored (enhanced) to determine a total number of mitigation credits lost or generated, respectively. Most evaluative factors are scored categorically according to condition classes defined in the SOP itself. Some are conceptually rooted in ecological or functional condition of the resources (e.g. Existing Condition of the resource to be impacted; Net Improvement at a mitigation site), while others address programmatic priorities of the CWA 404 regulatory program and/or value judgments of the agency or agencies that play a role in its administration (e.g. Lost Type or Dominant Impact of the resource to be impacted; Control or Location of the proposed mitigation site).</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | Not stated. | |
| Related Procedures/References | <p>Federal Interagency Stream Restoration Working Group. 1998. Stream Corridor Restoration; Principles, Processes, and Practices. National Technical Information Service, Springfield, Virginia.</p> <p>Harrelson, CC., C.L. Rawlins, and J.P. Potyondy. 1994. Stream Channel Reference Sites: An Illustrated Guide to Field Technique. General Technical Report RM-245, U.S. Forest Service Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.</p> <p>NRCS. 1996. Streambank and shoreline protection. In Engineering field handbook, Part 650, Chapter 16, United States Department of Agriculture, Natural Resources Conservation Service .</p> <p>Rosgen, D.L. 1996. Applied River Morphology. Wildland Hydrology Books, Pagosa Springs, Colorado.</p> | |

| | | |
|---------------------------------|---|-----------------------|
| Name | [Kentucky] Draft Stream Relocation/Mitigation Guidelines | Catalog No. 30 |
| Primary Author/ Agency | Kentucky Division of Water KDOW. 2007. Draft Stream Relocation/Mitigation Guidelines, revised October 15, 2007. Kentucky Natural Resources and Environmental Protection Cabinet, Division of Water, Frankfort, KY. | |
| Electronic Resource | http://www.water.ky.gov/permitting/wqcert/ | |
| Intended Use/Purpose | Regulatory Assessment (Clean Water Act, Section 401 Water Quality Certification); Compensatory Mitigation Protocol. | |
| Target Resource Type | Wadeable Streams: Intermittent and Perennial | |
| Scale/Unit of Assessment | Stream reach of unspecified length | |
| Geographic Applicability | Kentucky | |
| General Level of Effort | Moderate to Intensive (1 day± in the field by a trained or experienced crew of 2 or more persons) | |
| Assessment Parameters | <p>Channel/Valley Morphology: Bankfull stream discharge; Level II stream type (Rosgen, 1996); dimensionless critical shear stress & shear stress values; longitudinal channel profile (bankfull water surface elevation, channel gradient, valley gradient, pool and riffle gradient); planform (sinuosity, belt width, radius of curvature, meander wave length, floodprone area width); channel cross-sections (channel width & depth in riffles & pools, bankfull cross-sectional area, bankfull width, wetted perimeter, entrenchment ratio, hydraulic radius; floodprone area); substrate particle size (pebble count & sieve sample); riffle:pool ratio & placement.</p> <p>Physical Habitat: - -</p> <p>Water Quality: - -</p> <p>Biology: Determined on a case-by-case basis.</p> <p>Other: Watershed area.</p> | |
| Resolution | Semi-Quantitative (ordinal scale, rank, etc.); Quantitative (actual measurement or estimate) | |
| Output | Condition Assessment; Index (e.g. numeric score); Raw data. | |
| Reference | Measured External Reference Required (Site specific). | |
| QA/QC | | |

| | | |
|---|---|-----------------------|
| Name | [Kentucky] Draft Stream Relocation/Mitigation Guidelines | Catalog No. 30 |
| Description/ Summary | <p>The Draft Stream Relocation/Mitigation Guidelines from the Kentucky Division of Water (KDOW) provides detailed guidance on mitigation requirements and monitoring and assessment requirements for stream relocations and mitigation projects in the Commonwealth of Kentucky. Mitigation requirements themselves are based on ratios, dependent on the type of mitigation actions proposed. For example, stream enhancement measures will require a greater linear stream length of mitigation relative to stream restoration activities used to mitigate equivalent impacts.</p> <p>Although monitoring and assessment requirements are generally provided in outline form, the requirements themselves are discussed in detail, and suitable methods are referenced. Requisite data to support stream relocation or mitigation projects include longitudinal channel profiles for the impact reach, reference stream segment, and post-construction channel. Planform information must also be presented for both the reference stream segment and post-construction channel. Channel cross-sections must be collected from meander bends and straight reaches of the channel in both the reference stream segment and post-construction channel. The Guidelines refer to Harrelson et al. (1994) for appropriate stream surveying procedures.</p> <p>Requisite monitoring parameters are clearly indicated in the Guidelines and include most of the above referenced parameters, in addition to riparian vegetation (density, percent cover, and dominance) and the rapid stream Habitat Assessment Index from the USEPA RBP (Barbour et al., 1999). Tentative habitat criteria relating the RBP Habitat Assessment Index to biological conditions for each of Kentucky's ecoregions has been compiled, and is presented in Chapter 6 of KDOW (2002). When biological monitoring is required for stream relocation or mitigation projects, standard methods in KDOW (2002) must be followed.</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | Not stated. | |
| Related Procedures/References | <p>Barbour, M.T., J. Gerritsen, B.D. Snyder, and J.B. Stribling. 1999. Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition. EPA 841-B-99-002. USEPA Office of Water, Washington, D.C.</p> <p>Harrelson, CC., C.L. Rawlins, and J.P. Potyondy. 1994. Stream Channel Reference Sites: An Illustrated Guide to Field Technique. General Technical Report RM-245, U.S. Forest Service Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.</p> <p>KDOW. 2002. Methods for Assessing Biological Integrity of Surface Waters. July 2002. Kentucky Natural Resources and Environmental Protection Cabinet, Division of Water, Frankfort, KY.</p> | |
| Other/Notes | <p>The above KDOW referenced web site also includes links to documents reporting regional bankfull channel characteristics (aka hydraulic regional curves) for each ecoregion in Kentucky. Some of these documents also include stream channel morphological data collected from select designated KDOW biological reference streams and conclude with a discussion on how the regional relationships may be used during stream assessment and restoration design.</p> | |

| | | | |
|---------------------------------|--|--|-----------------------|
| Name | Stream Assessment Protocol for Headwater Streams in the Eastern Kentucky Coalfield Region (eKY Protocol) | | Catalog No. 31 |
| Primary Author/ Agency | <p>U.S. Army Corps of Engineers, Louisville District [based in large part on work by Kentucky Division of Water]</p> <p>Sparks, E.J., J. Townsend, T. Hagman, and D. Messer. 2003a. Stream Assessment Protocol for Headwater Streams in the Eastern Kentucky Coalfield Region. Aquatic Resource News: A Regulatory Newsletter 2(1), U.S. Army Corps of Engineers, Institute for Water Resources, Alexandria, VA.</p> <p>Sparks, E.J., T.E. Hagman, D. Messer, and J.M. Townsend. 2003b. Eastern Kentucky Stream Assessment Protocol: Utility in Making Mitigation Decisions. Aquatic Resource News: A Regulatory Newsletter 2(2), U.S. Army Corps of Engineers, Institute for Water Resources, Alexandria, VA.</p> | | |
| Electronic Resource | <p>http://www.usace.army.mil/CECW/Pages/aqua_news.aspx</p> <p>See also Pond and McMurray (2002), http://www.water.ky.gov/sw/swmonitor/sop/</p> | | |
| Intended Use/Purpose | Regulatory Assessment (Clean Water Act, Section 404); Compensatory Mitigation Protocol. | | |
| Target Resource Type | Headwater Streams, either intermittent or perennial ~ 1 st and 2 nd order streams with a drainage area of generally <3 to 5 square miles [actual reference and test sites used to develop the Macroinvertebrate Bioassessment Index (MBI) ranged from 0.25 to 3.5 square miles] | | |
| Scale/Unit of Assessment | Stream reach, 100-meters | | |
| Geographic Applicability | Eastern Kentucky Coalfield Region, including portions of three Level III ecoregions: Southwestern Appalachians, 68; Central Appalachians, 69; and Western Allegheny Plateau, 70. | | |
| General Level of Effort | Easy to Moderate - The eKY Protocol utilizes both biotic and abiotic indices to reach an "Ecological Integrity Index," but allows for only the abiotic factors to be evaluated in the absence of comparable biotic data or when there is less time available for assessment (e.g. preliminary site visit). | | |
| Assessment Parameters | <p>Channel/Valley Morphology: - -</p> <p>Physical Habitat: Riparian zone width; embeddedness; rapid visual-based habitat assessment (RBP).</p> <p>Water Quality: Conductivity.</p> <p>Biology: Benthic macroinvertebrates (optional).</p> <p>Other: - -</p> | | |
| Resolution | <p><u>Sparks et al. (2003a)</u> Semi-Quantitative (ordinal scale, rank, etc.) Quantitative (actual measurement or estimate)</p> | <p><u>Pond and McMurray (2002)</u> Semi-Quantitative (ordinal scale, rank, etc.) Quantitative (actual measurement or estimate)</p> | |
| Output | Condition Assessment; Index (e.g. numeric score); Programmatic or Regulatory Support Information. | | |

| | | |
|---|---|-----------------------|
| Name | Stream Assessment Protocol for Headwater Streams in the Eastern Kentucky Coalfield Region (eKY Protocol) | Catalog No. 31 |
| Reference | Internal (e.g. Index calibrated to existing local or regional reference data); based on Pond and McMurray (2002) <i>a priori</i> classification of sites as representative of <i>least disturbed conditions</i> in the region during compilation of the MBI. | |
| QA/QC | Not stated. | |
| Description/ Summary | <p>Sparks et al. (2003a) utilized the Eastern Kentucky macroinvertebrate biological index (MBI) compiled by the Kentucky Division of Water (Pond and McMurray, 2002) to develop the eKY Protocol specifically for the U.S. Army Corps of Engineers, Louisville District in its administration of Section 404 of the Clean Water Act (CWA). Physical habitat metrics collected by Pond and McMurray (2002) during the development of the bioassessment index were mostly transect-based estimates, but not completely quantitative measurements. Three of these metrics, plus one water quality metric, collectively differentiated <i>a priori</i> reference and test sites with 98% accuracy: percent embeddedness, canopy cover, conductivity, and rapid habitat assessment score (Pond and McMurray, 2002). Pond and McMurray (2002) also evaluated a family-level MBI (F-MBI) and found a strong relationship between the F-MBI and the original genus level MBI.</p> <p>Recommendations for using the eKY Protocol include three components: characterization, assessment, and analysis (Sparks et al., 2003a). Characterization includes a checklist specific to the CWA 404 program for documenting potential consequences of a proposed dredge and fill project on the aquatic environment and describes the physical characteristics of the headwater stream ecosystem and surrounding landscape. Assessment involves calculation of the Ecological Integrity Index (EII) for both existing conditions and anticipated post-project conditions. Analysis includes utilization of the assessment results to evaluate the proposed project under the CWA 404(b)(1) Guidelines and to help define potential compensatory mitigation needs, if applicable.</p> <p>Sparks et al. (2003b) provide examples of how the eKY Protocol is used to evaluate projects in the CWA 404 regulatory program, including how assessment results are used to determine mitigation ratios. EII spreadsheet calculators and mitigation ratio calculators are available on the USACE, Louisville District web site, including spreadsheets developed to account for the temporal loss of ecosystem functions between project site impact and implementation of mitigation. Although these examples do not specifically illustrate the protocol's application in designing mitigation projects, Sparks et al. (2003b) stress that such projects should be designed using "sound principles of fluvial geomorphology...based on [project specific] reference reaches."</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | Most robust level of assessment is ideally based on macroinvertebrates sampled during the spring index period (mid-February to late-May). | |
| Related Procedures/ References | <p>Barbour, M.T., J. Gerritsen, B.D. Snyder, and J.B. Stribling. 1999. Rapid bioassessment protocols for use in streams and wadeable rivers: periphyton, benthic macroinvertebrates, and fish, second edition. EPA 841-B-99-002. Office of Water, U.S. Environmental Protection Agency, Washington, D.C.</p> <p>Pond, G.J. and S.E. McMurray. 2002. A Macroinvertebrate Bioassessment Index for Headwater Streams of the Eastern Coalfield Region, Kentucky. Kentucky Division of Water, Water Quality Branch, Frankfort, KY. 56 pp.</p> | |
| Other/Notes | | |

| | | |
|---------------------------------|---|-----------------------|
| Name | Stream Mitigation Guidelines [NC] | Catalog No. 32 |
| Primary Author/ Agency | U.S. Army Corps of Engineers, Wilmington District USACE. 2003. Stream Mitigation Guidelines. U.S. Army Corps of Engineers, Wilmington District, Wilmington, NC. | |
| Electronic Resource | http://www.saw.usace.army.mil/wetlands/Mitigation/stream_mitigation.html | |
| Intended Use/Purpose | Regulatory Assessment (USACE CWA Section 404; NCDWQ CWA Section 401); Compensatory Mitigation Protocol | |
| Target Resource Type | Non-tidal Streams | |
| Scale/Unit of Assessment | Stream reach of unspecified length | |
| Geographic Applicability | North Carolina | |
| General Level of Effort | Easy (rapid); Moderate. | |
| Assessment Parameters | <p><u>Stream Quality Assessment index (based on visual observation):</u></p> <p>Channel/Valley Morphology: Entrenchment; presence of adjacent floodplain; sinuosity; evidence of channel incision or widening; presence of major bank failures; presence of flow / persistence of pools; evidence of human alteration; rooting depth and density on banks; dominant substrate size class and diversity of size classes; riffle and pool abundance, depth and frequency.</p> <p>Physical Habitat: Riparian buffer width; presence of groundwater discharge; presence of adjacent wetlands; sediment input; in-stream habitat complexity; canopy coverage; embeddedness.</p> <p>Water Quality: Evidence of nutrient or chemical discharges.</p> <p>Biology: Invertebrates' abundance, taxa richness, and sensitivity; types of amphibians present; fish abundance and taxa diversity; wildlife use of stream and riparian zone.</p> <p>Other: Impact by agriculture, livestock, or timber production.</p> | |
| Resolution | Qualitative (descriptive) Semi-Quantitative (ordinal scale, rank, etc.) Quantitative (actual measurement or estimate) | |
| Output | Condition Assessment; Index (e.g. numeric score); Qualitative Description; Raw data Programmatic or Regulatory Support Information | |

| | | |
|---|---|-----------------------|
| Name | Stream Mitigation Guidelines [NC] | Catalog No. 32 |
| Reference | <p>Condition assessment for large streams is based on ecoregional data collected by the NCDWQ bioassessment program. Site specific physical and morphological data is required from an external site specific reference stream reach.</p> <p>Post-construction benthic macroinvertebrate sampling must also include a sample station upstream of the mitigation stream section (NCDWQ, 2001). In some cases, one of NCDWQ's regional biological reference stations will also be required for monitoring.</p> | |
| QA/QC | <p>NCDWQ has specific requirements for the development of a quality assurance plan for benthic macroinvertebrate sampling that must be first coordinated with NCDWQ. The QA plan must include standard operating procedures that clearly demonstrate the ability of those involved with collection, taxonomic analyses, and reporting of results (NCDWQ, 2001).</p> | |
| Description/ Summary | <p>Although the USACE Wilmington District bases stream mitigation requirements for CWA 404 permits on ratios, the integration of stream assessment information, tools, and guidance from various State and Federal sources that are included in the Stream Mitigation Guidelines (and directly referenced on the USACE Wilmington District's web site) warrants its inclusion in this review.</p> <p>Final compensatory mitigation requirements for streams in the USACE Wilmington District consist of mitigation ratios that are generally based on the existing stream channel conditions and four levels or types of mitigation activities described in the Guidance. These categorical levels vary by the proposed mitigation actions' degree of complexity and include geomorphic stream channel considerations, biological considerations, and water quality (chemical) considerations.</p> <p>Existing stream conditions for large streams and rivers (wetted width ≥ 4 meters) are assessed based on bioclassification criteria and rating protocols developed for some of the major ecoregions in North Carolina by the North Carolina Division of Water Quality (NCDWQ). These criteria themselves are based primarily on benthic macroinvertebrates community composition, but habitat quality and fish community conditions are also used to assess quality conditions for large streams and rivers. There are five (5) stream quality condition classes based on these criteria.</p> <p>The condition of small perennial streams (wetted width < 3 meters) is assessed using a Stream Quality Assessment Worksheet that provides an index based on scores from observations of 23 metrics apportioned into four categories: (1) physical conditions, (2) channel stability, (3) habitat, and (4) biology.</p> <p>Monitoring requirements in the USACE Wilmington District recommend stream dimension, pattern, and profile surveying using methods from Harrelson et al. (1994). Additional requisite monitoring elements are based on the type and spatial extent of mitigation activities conducted, but may include biological sampling (NCDWQ, 2001), channel stability analysis, and/or riparian vegetation survival and growth. Specific evaluation criteria for mitigation sites are provided.</p> | |
| Expertise Required | Not stated. | |
| Time Necessary to Conduct Assessment | Not stated. | |
| Seasonality | <p>NCDWQ (2001) recommends that benthic macroinvertebrate samples be collected during the summer (June – September) for mitigation projects in the mountain and piedmont ecoregions (including the Sand Hills), but during the winter (January - March) for mitigation projects in coastal plain swamp streams.</p> | |

| | | |
|--------------------------------------|--|-----------------------|
| Name | Stream Mitigation Guidelines [NC] | Catalog No. 32 |
| Related Procedures/References | <p>Doll, B.A., G.L. Grabow, K.R. Hall, J. Halley, W.A. Harman, G.D. Jennings, and D.E. Wise. 2003. Stream Restoration: A Natural Channel Design Handbook. NC Stream Restoration Institute, NC State University. http://www.bae.ncsu.edu/programs/extension/wqg/srp/guidebook.html</p> <p>Harrelson, CC., C.L. Rawlins, and J.P. Potyondy. 1994. Stream Channel Reference Sites: An Illustrated Guide to Field Technique. General Technical Report RM-245, U.S. Forest Service Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.</p> <p>NCDWQ. 2001. Interim, Internal Technical Guide: Benthic Macroinvertebrate Monitoring Protocols for Compensatory Stream Restoration Projects, North Carolina Division of Water Quality, 401/Wetlands Unit. December, 2001, Raleigh, NC.</p> | |
| Other/Notes | <p>Appendices to the Stream Mitigation Guidelines include hydraulic regional curves for North Carolina, as well as a fact sheet describing "Application of the Rosgen Stream Classification System to North Carolina." Links to the internet sites of North Carolina state agencies involved in stream assessment, monitoring, and mitigation are provided.</p> <p>North Carolina State University maintains a Stream Restoration Program (NCSRP) consisting largely of faculty of the Department of Biological and Agricultural Engineering, as well as North Carolina Sea Grant and off-campus Extension faculty. The goal of NCSRP is to improve water quality and aquatic ecology through research, demonstration projects, and education/training. Among the many technical resources compiled by NCSRP, Doll et al. (2003) compiled a handbook on natural channel design for stream restoration that is available on the NCSRP web site.</p> | |