

2. Trends in Greenhouse Gas Emissions

2.1 Recent Trends in U.S. Greenhouse Gas Emissions and Sinks

In 2013, total U.S. greenhouse gas emissions were 6,673.0 MMT or million metric tons CO₂ Eq. Total U.S. emissions have increased by 5.9 percent from 1990 to 2013, and emissions increased from 2012 to 2013 by 2.0 percent (127.9 MMT CO₂ Eq.). The increase from 2012 to 2013 was due to an increase in the carbon intensity of fuels consumed to generate electricity due to an increase in coal consumption, with decreased natural gas consumption. Additionally, cold winter conditions lead to an increase in fuels for the residential and commercial sectors for heating. In 2013 there also was an increase in industrial production across multiple sectors resulting in increases in industrial sector emissions. Lastly, transportation emissions increased as a result of a small increase in vehicle miles traveled (VMT) and fuel use across on-road transportation modes. Since 1990, U.S. emissions have increased at an average annual rate of 0.3 percent. Figure 2-1 through Figure 2-3 illustrate the overall trend in total U.S. emissions by gas, annual changes, and absolute changes since 1990.

Figure 2-1: U.S. Greenhouse Gas Emissions by Gas

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.




Figure 2-2: Annual Percent Change in U.S. Greenhouse Gas Emissions





Figure 2-3: Cumulative Change in Annual U.S. Greenhouse Gas Emissions Relative to 1990

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.



Overall, from 1990 to 2013, total emissions of CO₂ increased by 381.5 MMT CO₂ Eq. (7.4 percent), while total emissions of CH₄ decreased by 109.2 MMT CO₂ Eq. (14.6 percent), and total emissions of N₂O increased 25.3 MMT CO₂ Eq. (7.7 percent). During the same period, aggregate weighted emissions of HFCs, PFCs, SF₆, and NF₃ rose by 74.3 MMT CO₂ Eq. (72.9 percent). Despite being emitted in smaller quantities relative to the other principal greenhouse gases, emissions of HFCs, PFCs, SF₆, and NF₃ are significant because many of them have extremely high GWPs and, in the cases of PFCs, SF₆, and NF₃, long atmospheric lifetimes. Conversely, U.S. greenhouse gas emissions were partly offset by C sequestration in managed forests, trees in urban areas, agricultural soils, and landfilled yard trimmings. These were estimated to offset 13.2 percent of total emissions in 2013.

As the largest contributor to U.S. greenhouse gas emissions, carbon dioxide (CO₂) from fossil fuel combustion has accounted for approximately 77 percent of global warming potential (GWP) weighted emissions for the entire time series since 1990, from 75 percent of total GWP-weighted emissions in 1990 to 77 percent in 2013. Emissions from this source category grew by 8.8 percent (417.0 MMT CO₂ Eq.) from 1990 to 2013 and were responsible for most of the increase in national emissions during this period. From 2012 to 2013, these emissions increased by 2.6 percent (131.7 MMT CO₂ Eq.). Historically, changes in emissions from fossil fuel combustion have been the dominant factor affecting U.S. emission trends.

Changes in CO₂ emissions from fossil fuel combustion are influenced by many long-term and short-term factors, including population and economic growth, energy price fluctuations, technological changes, and seasonal temperatures. On an annual basis, the overall consumption of fossil fuels in the United States fluctuates primarily in response to changes in general economic conditions, energy prices, weather, and the availability of non-fossil alternatives. For example, in a year with increased consumption of goods and services, low fuel prices, severe summer and winter weather conditions, nuclear plant closures, and lower precipitation feeding hydroelectric dams,

there would likely be proportionally greater fossil fuel consumption than in a year with poor economic performance, high fuel prices, mild temperatures, and increased output from nuclear and hydroelectric plants.

In the longer-term, energy consumption patterns respond to changes that affect the scale of consumption (e.g., population, number of cars, and size of houses), the efficiency with which energy is used in equipment (e.g., cars, power plants, steel mills, and light bulbs) and behavioral choices (e.g., walking, bicycling, or telecommuting to work instead of driving).

Energy-related CO₂ emissions also depend on the type of fuel or energy consumed and its carbon (C) intensity. Producing a unit of heat or electricity using natural gas instead of coal, for example, can reduce the CO₂ emissions because of the lower C content of natural gas.

A brief discussion of the year to year variability in fuel combustion emissions is provided below, beginning with 2009.

From 2009 to 2010, CO₂ emissions from fossil fuel combustion increased by 3.3 percent, which represents one of the largest annual increases in CO₂ emissions from fossil fuel combustion for the twenty four-year period from 1990 to 2013. This increase is primarily due to an increase in economic output from 2009 to 2010, and increased industrial production and manufacturing output (FRB 2014). Carbon dioxide emissions from fossil fuel combustion in the industrial sector increased by 6.6 percent, including increased emissions from the combustion of fuel oil, natural gas and coal. Overall, coal consumption increased by 5.8 percent, the largest annual increase in coal consumption for the twenty four-year period between 1990 and 2013. In 2010, weather conditions remained fairly constant in the winter and were much hotter in the summer compared to 2009, as heating degree days decreased slightly by 0.4 percent and cooling degree days increased by 17.3 percent to their highest levels in the twenty one-year period from 1990 to 2010. As a result of the more energy-intensive summer weather conditions, electricity sales to the residential and commercial end-use sectors in 2010 increased approximately 6.0 percent and 1.8 percent, respectively.

From 2010 to 2011, CO₂ emissions from fossil fuel combustion decreased by 2.5 percent. This decrease is a result of multiple factors including: (1) a decrease in the carbon intensity of fuels consumed to generate electricity due to a decrease in coal consumption, with increased natural gas consumption and a significant increase in hydropower used; (2) a decrease in transportation-related energy consumption due to higher fuel costs, improvements in fuel efficiency, and a reduction in miles traveled; and (3) relatively mild winter conditions resulting in an overall decrease in energy demand in most sectors. Changing fuel prices played a role in the decreasing emissions. A significant increase in the price of motor gasoline in the transportation sector was a major factor leading to a decrease in energy consumption by 1.2 percent. In addition, an increase in the price of coal and a concurrent decrease in natural gas prices led to a 5.7 percent decrease and a 2.5 percent increase in fuel consumption of these fuels by electric generators. This change in fuel prices also reduced the carbon intensity of fuels used to produce electricity in 2011, further contributing to the decrease in fossil fuel combustion emissions.

From 2011 to 2012, CO₂ emissions from fossil fuel combustion decreased by 3.9 percent, with emissions from fossil fuel combustion at their lowest level since 1994. This decrease from 2011 to 2012 is primarily a result of the decrease in the carbon intensity of fuels used to generate electricity due to a slight increase in the price of coal, and a significant decrease in the price of natural gas. The consumption of coal used to generate electricity decreased by 12.3 percent, while consumption of natural gas for electricity generation increased by 20.4 percent. Also, emissions declined in the transportation sector largely due to a small increase in fuel efficiency across different transportation modes and limited new demand for passenger transportation. In 2012, weather conditions remained fairly constant in the summer and were much warmer in the winter compared to 2011, as cooling degree days increased by 1.7 percent while heating degree days decreased 12.6 percent. This decrease in heating degree days resulted in a decreased demand for heating fuel in the residential and commercial sector, which had a decrease in natural gas consumption of 11.7 and 8.0 percent, respectively.

From 2012 to 2013, CO₂ emissions from fossil fuel combustion increased by 2.6 percent, this increase is primarily a result of the increased energy consumption in the residential and commercial sectors, as heating degree days increased 18.5 percent in 2013 as compared to 2012. The cooler weather led to an increase of 16.9 and 12.4 percent direct use of fuels in the residential and commercial sectors, respectively. In addition, there was an increase of 1.2 and 0.9 percent in electricity consumption in the residential and commercial sectors, respectively, due to regions that heat their homes with electricity. The consumption of natural gas used to generate electricity decreased by 10.2 percent due to an increase in the price of natural gas. Electric power plants shifted some consumption from natural

gas to coal, and as a result increased coal consumption to generate electricity by 4.2 percent. Lastly, industrial production increased 2.9 percent from 2012 to 2013, resulting in an increase in the in CO₂ emissions from fossil fuel combustion from the industrial sector by 4.2 percent.

Table 2-1 summarizes emissions and sinks from all U.S. anthropogenic sources in weighted units of MMT CO₂ Eq., while unweighted gas emissions and sinks in kilotons (kt) are provided in Table 2-2.

Table 2-1: Recent Trends in U.S. Greenhouse Gas Emissions and Sinks (MMT CO₂ Eq.)

Gas/Source	1990	2005	2009	2010	2011	2012	2013
CO₂	5,123.7	6,134.0	5,500.6	5,704.5	5,568.9	5,358.3	5,505.2
Fossil Fuel Combustion	4,740.7	5,747.7	5,197.1	5,367.1	5,231.3	5,026.0	5,157.7
Electricity Generation	1,820.8	2,400.9	2,145.7	2,258.4	2,157.7	2,022.2	2,039.8
Transportation	1,493.8	1,887.8	1,720.3	1,732.0	1,711.5	1,700.8	1,718.4
Industrial	842.5	827.8	727.7	775.7	774.1	784.2	817.3
Residential	338.3	357.8	336.4	334.7	327.2	283.1	329.6
Commercial	217.4	223.5	223.5	220.2	221.0	197.1	220.7
U.S. Territories	27.9	49.9	43.5	46.2	39.8	38.6	32.0
Non-Energy Use of Fuels	117.7	138.9	106.0	114.6	108.4	104.9	119.8
Iron and Steel Production & Metallurgical Coke Production	99.8	66.7	43.0	55.7	60.0	54.3	52.3
Natural Gas Systems	37.6	30.0	32.2	32.3	35.6	34.8	37.8
Cement Production	33.3	45.9	29.4	31.3	32.0	35.1	36.1
Petrochemical Production	21.6	28.1	23.7	27.4	26.4	26.5	26.5
Lime Production	11.7	14.6	11.4	13.4	14.0	13.7	14.1
Ammonia Production	13.0	9.2	8.5	9.2	9.3	9.4	10.2
Incineration of Waste	8.0	12.5	11.3	11.0	10.5	10.4	10.1
Petroleum Systems	4.4	4.9	4.7	4.2	4.5	5.1	6.0
Liming of Agricultural Soils	4.7	4.3	3.7	4.8	3.9	5.8	5.9
Urea Consumption for Non-Agricultural Purposes	3.8	3.7	3.4	4.7	4.0	4.4	4.7
Other Process Uses of Carbonates	4.9	6.3	7.6	9.6	9.3	8.0	4.4
Urea Fertilization	2.4	3.5	3.6	3.8	4.1	4.2	4.0
Aluminum Production	6.8	4.1	3.0	2.7	3.3	3.4	3.3
Soda Ash Production and Consumption	2.7	2.9	2.5	2.6	2.6	2.7	2.7
Ferroalloy Production	2.2	1.4	1.5	1.7	1.7	1.9	1.8
Titanium Dioxide Production	1.2	1.8	1.6	1.8	1.7	1.5	1.6
Zinc Production	0.6	1.0	0.9	1.2	1.3	1.5	1.4
Phosphoric Acid Production	1.6	1.4	1.0	1.1	1.2	1.1	1.2
Glass Production	1.5	1.9	1.0	1.5	1.3	1.2	1.2
Carbon Dioxide Consumption	1.5	1.4	1.8	1.2	0.8	0.8	0.9
Peatlands Remaining Peatlands	1.1	1.1	1.0	1.0	0.9	0.8	0.8
Lead Production	0.5	0.6	0.5	0.5	0.5	0.5	0.5
Silicon Carbide Production and Consumption	0.4	0.2	0.1	0.2	0.2	0.2	0.2
Magnesium Production and Processing	+	+	+	+	+	+	+
<i>Land Use, Land-Use Change, and Forestry (Sink)^a</i>	<i>(775.8)</i>	<i>(911.9)</i>	<i>(870.9)</i>	<i>(871.6)</i>	<i>(881.0)</i>	<i>(880.4)</i>	<i>(881.7)</i>
<i>Wood Biomass and Ethanol Consumption^b</i>	<i>219.4</i>	<i>229.8</i>	<i>250.5</i>	<i>265.1</i>	<i>268.1</i>	<i>267.7</i>	<i>283.3</i>
<i>International Bunker Fuels^c</i>	<i>103.5</i>	<i>113.1</i>	<i>106.4</i>	<i>117.0</i>	<i>111.7</i>	<i>105.8</i>	<i>99.8</i>
CH₄	745.5	707.8	709.5	667.2	660.9	647.6	636.3
Enteric Fermentation	164.2	168.9	172.7	171.1	168.7	166.3	164.5
Natural Gas Systems	179.1	176.3	168.0	159.6	159.3	154.4	157.4
Landfills	186.2	165.5	158.1	121.8	121.3	115.3	114.6
Coal Mining	96.5	64.1	79.9	82.3	71.2	66.5	64.6
Manure Management	37.2	56.3	59.7	60.9	61.4	63.7	61.4

Petroleum Systems	31.5	23.5	21.5	21.3	22.0	23.3	25.2
Wastewater Treatment	15.7	15.9	15.6	15.5	15.3	15.2	15.0
Rice Cultivation	9.2	8.9	9.4	11.1	8.5	9.3	8.3
Stationary Combustion	8.5	7.4	7.4	7.1	7.1	6.6	8.0
Abandoned Underground Coal Mines	7.2	6.6	6.4	6.6	6.4	6.2	6.2
Forest Fires	2.5	8.3	5.8	4.7	14.6	15.7	5.8
Mobile Combustion	5.6	3.0	2.3	2.3	2.3	2.2	2.1
Composting	0.4	1.9	1.9	1.8	1.9	1.9	2.0
Iron and Steel Production & Metallurgical Coke Production	1.1	0.9	0.4	0.6	0.7	0.7	0.7
Field Burning of Agricultural Residues	0.3	0.2	0.3	0.3	0.3	0.3	0.3
Petrochemical Production	0.2	0.1	+	0.1	+	0.1	0.1
Ferroalloy Production	+	+	+	+	+	+	+
Silicon Carbide Production and Consumption	+	+	+	+	+	+	+
Peatlands Remaining Peatlands	+	+	+	+	+	+	+
Incineration of Waste	+	+	+	+	+	+	+
<i>International Bunker Fuels^c</i>	0.2	0.1	0.1	0.1	0.1	0.1	0.1
N₂O	329.9	355.9	356.1	360.1	371.9	365.6	355.2
Agricultural Soil Management	224.0	243.6	264.1	264.3	265.8	266.0	263.7
Stationary Combustion	11.9	20.2	20.4	22.2	21.3	21.4	22.9
Mobile Combustion	41.2	38.1	24.6	23.7	22.5	20.2	18.4
Manure Management	13.8	16.4	17.0	17.1	17.3	17.3	17.3
Nitric Acid Production	12.1	11.3	9.6	11.5	10.9	10.5	10.7
Wastewater Treatment	3.4	4.3	4.6	4.7	4.8	4.9	4.9
N ₂ O from Product Uses	4.2	4.2	4.2	4.2	4.2	4.2	4.2
Adipic Acid Production	15.2	7.1	2.7	4.2	10.2	5.5	4.0
Forest Fires	1.7	5.5	3.8	3.1	9.6	10.3	3.8
Settlement Soils	1.4	2.3	2.2	2.4	2.5	2.5	2.4
Composting	0.3	1.7	1.7	1.6	1.7	1.7	1.8
Forest Soils	0.1	0.5	0.5	0.5	0.5	0.5	0.5
Incineration of Waste	0.5	0.4	0.3	0.3	0.3	0.3	0.3
Semiconductor Manufacture	+	0.1	0.1	0.1	0.2	0.2	0.2
Field Burning of Agricultural Residues	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Peatlands Remaining Peatlands	+	+	+	+	+	+	+
<i>International Bunker Fuels^c</i>	0.9	1.0	0.9	1.0	1.0	0.9	0.9
HFCs	46.6	131.4	142.9	152.6	157.4	159.2	163.0
Substitution of Ozone Depleting Substances ^d	0.3	111.1	136.0	144.4	148.4	153.5	158.6
HCFC-22 Production	46.1	20.0	6.8	8.0	8.8	5.5	4.1
Semiconductor Manufacture	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Magnesium Production and Processing	0.0	0.0	+	+	+	+	0.1
PFCs	24.3	6.6	3.9	4.4	6.9	6.0	5.8
Aluminum Production	21.5	3.4	1.9	1.9	3.5	2.9	3.0
Semiconductor Manufacture	2.8	3.2	2.0	2.6	3.4	3.0	2.9
SF₆	31.1	14.0	9.3	9.5	10.0	7.7	6.9
Electrical Transmission and Distribution	25.4	10.6	7.3	7.0	6.8	5.7	5.1
Magnesium Production and Processing	5.2	2.7	1.6	2.1	2.8	1.6	1.4
Semiconductor Manufacture	0.5	0.7	0.3	0.4	0.4	0.4	0.4
NF₃	+	0.5	0.4	0.5	0.7	0.6	0.6
Semiconductor Manufacture	+	0.5	0.4	0.5	0.7	0.6	0.6
Total Emissions	6,301.1	7,350.2	6,722.7	6,898.8	6,776.6	6,545.1	6,673.0
Total Sinks^a	(775.8)	(911.9)	(870.9)	(871.6)	(881.0)	(880.4)	(881.7)

Net Emissions (Sources and Sinks)	5,525.2	6,438.3	5,851.9	6,027.2	5,895.6	5,664.7	5,791.2
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.

+ Does not exceed 0.05 MMT CO₂ Eq.

^a Parentheses indicate negative values or sequestration. Sinks (i.e., CO₂ removals) are only included in the Net Emissions total. Refer to Table 2-8 for a breakout of emissions and removals for Land Use, Land-Use Change, and Forestry by gas and source category.

^b Emissions from Wood Biomass and Ethanol Consumption are not included specifically in summing energy sector totals. Net carbon fluxes from changes in biogenic carbon reservoirs are accounted for in the estimates for Land Use, Land-Use Change, and Forestry.

^c Emissions from International Bunker Fuels are not included in totals.

^d Small amounts of PFC emissions also result from this source.

Note: Totals may not sum due to independent rounding.

Table 2-2: Recent Trends in U.S. Greenhouse Gas Emissions and Sinks (kt)

Gas/Source	1990	2005	2009	2010	2011	2012	2013
CO₂	5,123,695	6,133,969	5,500,602	5,704,531	5,568,891	5,358,276	5,505,178
Fossil Fuel Combustion	4,740,670	5,747,683	5,197,058	5,367,144	5,231,341	5,026,000	5,157,697
Electricity Generation	1,820,818	2,400,874	2,145,658	2,258,399	2,157,688	2,022,181	2,039,750
Transportation	1,493,758	1,887,799	1,720,314	1,731,971	1,711,538	1,700,782	1,718,406
Industrial	842,473	827,808	727,724	775,674	774,101	784,227	817,252
Residential	338,347	357,827	336,375	334,734	327,211	283,095	329,609
Commercial	217,393	223,453	223,492	220,195	221,022	197,097	220,714
U.S. Territories	27,882	49,923	43,495	46,172	39,781	38,617	31,965
Non-Energy Use of Fuels	117,658	138,877	106,018	114,554	108,359	104,917	119,850
Iron and Steel Production & Metallurgical Coke Production	99,781	66,666	43,029	55,746	60,008	54,327	52,288
Natural Gas Systems	37,645	29,995	32,201	32,334	35,551	34,764	37,808
Cement Production	33,278	45,910	29,432	31,256	32,010	35,051	36,146
Petrochemical Production	21,633	28,124	23,706	27,388	26,396	26,477	26,514
Lime Production	11,700	14,552	11,411	13,381	13,981	13,715	14,072
Ammonia Production	13,047	9,196	8,454	9,188	9,292	9,377	10,152
Incineration of Waste	7,972	12,454	11,295	11,026	10,550	10,363	10,137
Petroleum Systems	4,445	4,904	4,656	4,153	4,467	5,060	6,001
Liming of Agricultural Soils	4,667	4,349	3,669	4,784	3,871	5,776	5,925
Urea Consumption for Non-Agricultural Purposes	3,784	3,653	3,427	4,730	4,029	4,449	4,663
Other Process Uses of Carbonates	4,907	6,339	7,583	9,560	9,335	8,022	4,424
Urea Fertilization	2,417	3,504	3,555	3,778	4,099	4,225	4,011
Aluminum Production	6,831	4,142	3,009	2,722	3,292	3,439	3,255
Soda Ash Production and Consumption	2,741	2,868	2,488	2,612	2,624	2,672	2,712
Ferroalloy Production	2,152	1,392	1,469	1,663	1,735	1,903	1,785
Titanium Dioxide Production	1,195	1,755	1,648	1,769	1,729	1,528	1,608
Zinc Production	632	1,030	943	1,182	1,286	1,486	1,429
Phosphoric Acid Production	1,586	1,395	1,016	1,130	1,198	1,138	1,173
Glass Production	1,535	1,928	1,045	1,481	1,299	1,248	1,160
Carbon Dioxide Consumption	1,472	1,375	1,795	1,206	802	841	903
Peatlands Remaining Peatlands	1,055	1,101	1,024	1,022	926	812	770
Lead Production	516	553	525	542	538	527	525
Silicon Carbide Production and Consumption	375	219	145	181	170	158	169
Magnesium Production and Processing	1	3	1	1	3	2	2
<i>Land Use, Land-Use Change, and Forestry (Sink)^a</i>	<i>(775,835)</i>	<i>(911,929)</i>	<i>(870,879)</i>	<i>(871,609)</i>	<i>(880,999)</i>	<i>(880,394)</i>	<i>(881,732)</i>
<i>Wood Biomass and Ethanol Consumption^b</i>	<i>219,413</i>	<i>229,844</i>	<i>250,491</i>	<i>265,110</i>	<i>268,064</i>	<i>267,730</i>	<i>283,337</i>
<i>International Bunker Fuels^c</i>	<i>103,463</i>	<i>113,139</i>	<i>106,410</i>	<i>116,992</i>	<i>111,660</i>	<i>105,805</i>	<i>99,763</i>
CH₄	29,820	28,314	28,380	26,687	26,437	25,905	25,453
Enteric Fermentation	6,566	6,755	6,908	6,844	6,750	6,653	6,581
Natural Gas Systems	7,165	7,053	6,722	6,382	6,371	6,176	6,295
Landfills	7,450	6,620	6,324	4,873	4,851	4,611	4,585
Coal Mining	3,860	2,565	3,194	3,293	2,849	2,658	2,584
Manure Management	1,486	2,254	2,388	2,437	2,457	2,548	2,456
Petroleum Systems	1,261	939	860	854	878	931	1,009
Wastewater Treatment	626	635	623	619	610	606	601
Rice Cultivation	366	358	378	444	339	372	332
Stationary Combustion	339	296	295	283	283	264	318

Abandoned Underground Coal								
Mines	288	264	254	263	257	249	249	
Forest Fires	101	332	233	190	584	626	233	
Mobile Combustion	225	121	93	92	91	88	86	
Composting	15	75	75	73	75	77	79	
Iron and Steel Production & Metallurgical Coke Production	46	34	17	25	28	29	28	
Field Burning of Agricultural Residues	13	9	12	11	12	12	12	
Petrochemical Production	9	6	2	2	2	3	3	
Ferroalloy Production	1	+	+	+	+	1	+	
Silicon Carbide Production and Consumption	1	+	+	+	+	+	+	
Peatlands Remaining Peatlands	+	+	+	+	+	+	+	
Incineration of Waste	+	+	+	+	+	+	+	
<i>International Bunker Fuels^c</i>	7	5	5	6	5	4	3	
N₂O	1,107	1,194	1,195	1,208	1,248	1,227	1,192	
Agricultural Soil Management	752	817	886	887	892	892	885	
Stationary Combustion	40	68	69	74	71	72	77	
Mobile Combustion	138	128	82	80	76	68	62	
Manure Management	46	55	57	57	58	58	58	
Nitric Acid Production	41	38	32	39	37	35	36	
Wastewater Treatment	11	15	16	16	16	16	17	
N ₂ O from Product Uses	14	14	14	14	14	14	14	
Adipic Acid Production	51	24	9	14	34	19	13	
Forest Fires	6	18	13	11	32	35	13	
Settlement Soils	5	8	8	8	8	8	8	
Composting	1	6	6	5	6	6	6	
Forest Soils	+	2	2	2	2	2	2	
Incineration of Waste	2	1	1	1	1	1	1	
Semiconductor Manufacture	+	+	+	+	1	1	1	
Field Burning of Agricultural Residues	+	+	+	+	+	+	+	
Peatlands Remaining Peatlands	+	+	+	+	+	+	+	
<i>International Bunker Fuels^c</i>	3	3	3	3	3	3	3	
HFCs	M							
Substitution of Ozone Depleting Substances ^d	M	M	M	M	M	M	M	
HCFC-22 Production	3	1	+	1	1	+	+	
Semiconductor Manufacture	+	+	+	+	+	+	+	
Magnesium Production and Processing	0	0	+	+	+	+	+	
PFCs	M							
Aluminum Production	M	M	M	M	M	M	M	
Semiconductor Manufacture	M	M	M	M	M	M	M	
SF₆	1	+	+	+	+	+	+	
Electrical Transmission and Distribution	1	+	+	+	+	+	+	
Magnesium Production and Processing	+	+	+	+	+	+	+	
Semiconductor Manufacture	+	+	+	+	+	+	+	
NF₃	+							
Semiconductor Manufacture	+	+	+	+	+	+	+	

+ Does not exceed 0.5 kt.

M Mixture of multiple gases

^a Refer to Table 2-8 for a breakout of emissions and removals for Land Use, Land-Use Change, and Forestry by gas and source category.

^b Emissions from Wood Biomass and Ethanol Consumption are not included specifically in summing energy sector totals. Net carbon fluxes from changes in biogenic carbon reservoirs are accounted for in the estimates for Land Use, Land-Use Change, and Forestry

^c Emissions from International Bunker Fuels are not included in totals.

^d Small amounts of PFC emissions also result from this source.

Note: Totals may not sum due to independent rounding. Parentheses indicate negative values or sequestration.

Emissions of all gases can be summed from each source category into a set of five sectors defined by the Intergovernmental Panel on Climate Change (IPCC). Over the twenty four-year period of 1990 to 2013, total emissions in the Energy, Industrial Processes and Product Use, and Agriculture sectors grew by 346.2 MMT CO₂ Eq. (6.5 percent), 17.0 MMT CO₂ Eq. (5.0 percent), and 67.0 MMT CO₂ Eq. (14.9 percent), respectively. Emissions from the Waste sector decreased by 67.7 MMT CO₂ Eq. (32.9 percent). Over the same period, estimates of net C sequestration for the Land Use, Land-Use Change, and Forestry sector increased by 96.4 MMT CO₂ Eq. (12.7 percent).

Figure 2-4: U.S. Greenhouse Gas Emissions and Sinks by Chapter/IPCC Sector

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.




Table 2-3: Recent Trends in U.S. Greenhouse Gas Emissions and Sinks by Chapter/IPCC Sector (MMT CO₂ Eq.)

Chapter/IPCC Sector	1990	2005	2009	2010	2011	2012	2013
Energy	5,290.5	6,273.6	5,682.1	5,854.6	5,702.6	5,482.2	5,636.6
Fossil Fuel Combustion	4,740.7	5,747.7	5,197.1	5,367.1	5,231.3	5,026.0	5,157.7
Natural Gas Systems	216.8	206.3	200.2	191.9	194.8	189.2	195.2
Non-Energy Use of Fuels	117.7	138.9	106.0	114.6	108.4	104.9	119.8
Coal Mining	96.5	64.1	79.9	82.3	71.2	66.5	64.6
Petroleum Systems	36.0	28.4	26.2	25.5	26.4	28.3	31.2
Stationary Combustion	20.4	27.6	27.8	29.3	28.4	28.0	30.8
Mobile Combustion	46.9	41.1	26.9	26.0	24.8	22.4	20.6
Incineration of Waste	8.4	12.8	11.6	11.4	10.9	10.7	10.4
Abandoned Underground Coal Mines	7.2	6.6	6.4	6.6	6.4	6.2	6.2
Industrial Processes and Product Use	342.1	367.4	314.9	353.6	371.0	361.2	359.1
Substitution of Ozone Depleting Substances	0.3	111.1	136.0	144.4	148.4	153.5	158.6
Iron and Steel Production & Metallurgical Coke Production	100.9	67.5	43.5	56.4	60.7	55.1	53.0

Cement Production	33.3	45.9	29.4	31.3	32.0	35.1	36.1
Petrochemical Production	21.9	28.3	23.8	27.4	26.4	26.5	26.6
Lime Production	11.7	14.6	11.4	13.4	14.0	13.7	14.1
Nitric Acid Production	12.1	11.3	9.6	11.5	10.9	10.5	10.7
Ammonia Production	13.0	9.2	8.5	9.2	9.3	9.4	10.2
Aluminum Production	28.3	7.6	4.9	4.6	6.8	6.4	6.2
Electrical Transmission and Distribution	25.4	10.6	7.3	7.0	6.8	5.7	5.1
Urea Consumption for Non-Agricultural Purposes	3.8	3.7	3.4	4.7	4.0	4.4	4.7
Other Process Uses of Carbonates	4.9	6.3	7.6	9.6	9.3	8.0	4.4
N ₂ O from Product Uses	4.2	4.2	4.2	4.2	4.2	4.2	4.2
Semiconductor Manufacture	3.6	4.7	3.1	3.8	4.9	4.5	4.2
HCFC-22 Production	46.1	20.0	6.8	8.0	8.8	5.5	4.1
Adipic Acid Production	15.2	7.1	2.7	4.2	10.2	5.5	4.0
Soda Ash Production and Consumption	2.7	2.9	2.5	2.6	2.6	2.7	2.7
Ferroalloy Production	2.2	1.4	1.5	1.7	1.7	1.9	1.8
Titanium Dioxide Production	1.2	1.8	1.6	1.8	1.7	1.5	1.6
Magnesium Production and Processing	5.2	2.7	1.6	2.1	2.8	1.7	1.5
Zinc Production	0.6	1.0	0.9	1.2	1.3	1.5	1.4
Phosphoric Acid Production	1.6	1.4	1.0	1.1	1.2	1.1	1.2
Glass Production	1.5	1.9	1.0	1.5	1.3	1.2	1.2
Carbon Dioxide Consumption	1.5	1.4	1.8	1.2	0.8	0.8	0.9
Lead Production	0.5	0.6	0.5	0.5	0.5	0.5	0.5
Silicon Carbide Production and Consumption	0.4	0.2	0.2	0.2	0.2	0.2	0.2
Agriculture	448.7	494.5	523.3	524.8	522.1	523.0	515.7
Agricultural Soil Management	224.0	243.6	264.1	264.3	265.8	266.0	263.7
Enteric Fermentation	164.2	168.9	172.7	171.1	168.7	166.3	164.5
Manure Management	51.0	72.8	76.7	78.0	78.7	81.0	78.7
Rice Cultivation	9.2	8.9	9.4	11.1	8.5	9.3	8.3
Field Burning of Agricultural Residues	0.4	0.3	0.4	0.4	0.4	0.4	0.4
Land Use, Land-Use Change, and Forestry	13.8	25.5	20.6	20.3	36.1	39.8	23.3
Forest Fires	4.2	13.8	9.7	7.9	24.2	26.0	9.7
Liming of Agricultural Soils	4.7	4.3	3.7	4.8	3.9	5.8	5.9
Urea Fertilization	2.4	3.5	3.6	3.8	4.1	4.2	4.0
Settlement Soils	1.4	2.3	2.2	2.4	2.5	2.5	2.4
Peatlands Remaining Peatlands	1.1	1.1	1.0	1.0	0.9	0.8	0.8
Forest Soils	0.1	0.5	0.5	0.5	0.5	0.5	0.5
Waste	206.0	189.2	181.8	145.5	144.9	138.9	138.3
Landfills	186.2	165.5	158.1	121.8	121.3	115.3	114.6
Wastewater Treatment	19.0	20.2	20.2	20.2	20.1	20.0	20.0
Composting	0.7	3.5	3.6	3.5	3.5	3.7	3.7
Total Emissions	6,301.1	7,350.2	6,722.7	6,898.8	6,776.6	6,545.1	6,673.0
Total Sinks^a	(775.8)	(911.9)	(870.9)	(871.6)	(881.0)	(880.4)	(881.7)
Net Emissions (Sources and Sinks)	5,525.2	6,438.3	5,851.9	6,027.2	5,895.6	5,664.7	5,791.2

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.

^a Sinks (i.e., CO₂ removals) are only included in the Net Emissions total. Refer to Table 2-8 for a breakout of emissions and removals for Land Use, Land-Use Change, and Forestry by gas and source category.

Note: Totals may not sum due to independent rounding. Parentheses indicate negative values or sequestration.

Energy

Energy-related activities, primarily fossil fuel combustion, accounted for the vast majority of U.S. CO₂ emissions for the period of 1990 through 2013. In 2013, approximately 82 percent of the energy consumed in the United States (on a Btu basis) was produced through the combustion of fossil fuels. The remaining 18 percent came from other energy sources such as hydropower, biomass, nuclear, wind, and solar energy (see Figure 2-5 and Figure 2-6). A discussion of specific trends related to CO₂ as well as other greenhouse gas emissions from energy consumption is presented in the Energy chapter. Energy-related activities are also responsible for CH₄ and N₂O emissions (41 percent and 12 percent of total U.S. emissions of each gas, respectively). Table 2-4 presents greenhouse gas emissions from the Energy chapter, by source and gas.

Figure 2-5: 2013 Energy Chapter Greenhouse Gas Sources

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.




Figure 2-6: 2013 U.S. Fossil Carbon Flows (MMT CO₂ Eq.)

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.




Table 2-4: Emissions from Energy (MMT CO₂ Eq.)

Gas/Source	1990	2005	2009	2010	2011	2012	2013
CO₂	4,908.4	5,933.9	5,351.2	5,529.2	5,390.3	5,181.1	5,331.5
Fossil Fuel Combustion	4,740.7	5,747.7	5,197.1	5,367.1	5,231.3	5,026.0	5,157.7
Electricity Generation	1,820.8	2,400.9	2,145.7	2,258.4	2,157.7	2,022.2	2,039.8
Transportation	1,493.8	1,887.8	1,720.3	1,732.0	1,711.5	1,700.8	1,718.4
Industrial	842.5	827.8	727.7	775.7	774.1	784.2	817.3
Residential	338.3	357.8	336.4	334.7	327.2	283.1	329.6
Commercial	217.4	223.5	223.5	220.2	221.0	197.1	220.7
U.S. Territories	27.9	49.9	43.5	46.2	39.8	38.6	32.0
Non-Energy Use of Fuels	117.7	138.9	106.0	114.6	108.4	104.9	119.8
Natural Gas Systems	37.6	30.0	32.2	32.3	35.6	34.8	37.8
Incineration of Waste	8.0	12.5	11.3	11.0	10.5	10.4	10.1
Petroleum Systems	4.4	4.9	4.7	4.2	4.5	5.1	6.0
Biomass - Wood ^a	215.2	206.9	188.2	192.5	195.2	194.9	208.6
International Bunker Fuels ^b	103.5	113.1	106.4	117.0	111.7	105.8	99.8
Biomass - Ethanol ^a	4.2	22.9	62.3	72.6	72.9	72.8	74.7
CH₄	328.5	280.9	285.5	279.2	268.2	259.2	263.5
Natural Gas Systems	179.1	176.3	168.0	159.6	159.3	154.4	157.4
Coal Mining	96.5	64.1	79.9	82.3	71.2	66.5	64.6
Petroleum Systems	31.5	23.5	21.5	21.3	22.0	23.3	25.2
Stationary Combustion	8.5	7.4	7.4	7.1	7.1	6.6	8.0
Abandoned Underground Coal							
Mines	7.2	6.6	6.4	6.6	6.4	6.2	6.2
Mobile Combustion	5.6	3.0	2.3	2.3	2.3	2.2	2.1
Incineration of Waste	+	+	+	+	+	+	+
International Bunker Fuels ^b	0.2	0.1	0.1	0.1	0.1	0.1	0.1
N₂O	53.6	58.7	45.3	46.2	44.1	41.9	41.6
Stationary Combustion	11.9	20.2	20.4	22.2	21.3	21.4	22.9
Mobile Combustion	41.2	38.1	24.6	23.7	22.5	20.2	18.4

Incineration of Waste	0.5	0.4	0.3	0.3	0.3	0.3	0.3
<i>International Bunker Fuels^b</i>	0.9	1.0	0.9	1.0	1.0	0.9	0.9
Total	5,290.5	6,273.6	5,682.1	5,854.6	5,702.6	5,482.2	5,636.6

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.

Note: Totals may not sum due to independent rounding.

+ Does not exceed 0.05 MMT CO₂ Eq.

^a Emissions from Wood Biomass and Ethanol Consumption are not included specifically in summing energy sector totals. Net carbon fluxes from changes in biogenic carbon reservoirs are accounted for in the estimates for Land Use, Land-Use Change, and Forestry.

^b Emissions from International Bunker Fuels are not included in totals.

Carbon dioxide emissions from fossil fuel combustion are presented in Table 2-5 based on the underlying U.S. energy consumer data collected by EIA. Estimates of CO₂ emissions from fossil fuel combustion are calculated from these EIA “end-use sectors” based on total consumption and appropriate fuel properties (any additional analysis and refinement of the EIA data is further explained in the Energy chapter of this report). EIA’s fuel consumption data for the electric power sector comprises electricity-only and combined-heat-and-power (CHP) plants within the NAICS 22 category whose primary business is to sell electricity, or electricity and heat, to the public (nonutility power producers can be included in this sector as long as they meet the electric power sector definition). EIA statistics for the industrial sector include fossil fuel consumption that occurs in the fields of manufacturing, agriculture, mining, and construction. EIA’s fuel consumption data for the transportation sector consists of all vehicles whose primary purpose is transporting people and/or goods from one physical location to another. EIA’s fuel consumption data for the industrial sector consists of all facilities and equipment used for producing, processing, or assembling goods (EIA includes generators that produce electricity and/or useful thermal output primarily to support on-site industrial activities in this sector). EIA’s fuel consumption data for the residential sector consists of living quarters for private households. EIA’s fuel consumption data for the commercial sector consists of service-providing facilities and equipment from private and public organizations and businesses (EIA includes generators that produce electricity and/or useful thermal output primarily to support the activities at commercial establishments in this sector). Table 2-5 and Figure 2-7 summarize CO₂ emissions from fossil fuel combustion by end-use sector. Figure 2-8 further describes the total emissions from fossil fuel combustion, separated by end-use sector, including CH₄ and N₂O in addition to CO₂.

Table 2-5: CO₂ Emissions from Fossil Fuel Combustion by End-Use Sector (MMT CO₂ Eq.)

End-Use Sector	1990	2005	2009	2010	2011	2012	2013
Transportation	1,496.8	1,892.5	1,724.8	1,736.5	1,715.8	1,704.6	1,722.4
Combustion	1,493.8	1,887.8	1,720.3	1,732.0	1,711.5	1,700.8	1,718.4
Electricity	3.0	4.7	4.5	4.5	4.3	3.9	4.0
Industrial	1,529.2	1,564.4	1,329.5	1,416.5	1,398.8	1,377.0	1,399.8
Combustion	842.5	827.8	727.7	775.7	774.1	784.2	817.3
Electricity	686.7	736.6	601.8	640.8	624.7	592.8	582.5
Residential	931.4	1,214.1	1,122.6	1,174.8	1,117.9	1,008.4	1,070.2
Combustion	338.3	357.8	336.4	334.7	327.2	283.1	329.6
Electricity	593.0	856.3	786.2	840.1	790.7	725.3	740.6
Commercial	755.4	1,026.7	976.7	993.2	959.1	897.4	933.3
Combustion	217.4	223.5	223.5	220.2	221.0	197.1	220.7
Electricity	538.0	803.3	753.2	773.0	738.0	700.3	712.6
U.S. Territories^a	27.9	49.9	43.5	46.2	39.8	38.6	32.0
Total	4,740.7	5,747.7	5,197.1	5,367.1	5,231.3	5,026.0	5,157.7
Electricity Generation	1,820.8	2,400.9	2,145.7	2,258.4	2,157.7	2,022.2	2,039.8

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.

Note: Totals may not sum due to independent rounding. Combustion-related emissions from electricity generation are allocated based on aggregate national electricity consumption by each end-use sector.

^a Fuel consumption by U.S. Territories (i.e., American Samoa, Guam, Puerto Rico, U.S. Virgin Islands, Wake Island, and other U.S. Pacific Islands) is included in this report.

Figure 2-7: 2013 CO₂ Emissions from Fossil Fuel Combustion by Sector and Fuel Type

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.





Figure 2-8: 2013 End-Use Sector Emissions of CO₂ from Fossil Fuel Combustion

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.



The main driver of emissions in the Energy sector is CO₂ from fossil fuel combustion. Electricity generation is the largest emitter of CO₂, and electricity generators consumed 34 percent of U.S. energy from fossil fuels and emitted 40 percent of the CO₂ from fossil fuel combustion in 2013. Electricity generation emissions can also be allocated to the end-use sectors that are consuming that electricity, as presented in Table 2-5. The transportation end-use sector accounted for 1,722.4 MMT CO₂ Eq. in 2013 or approximately 33 percent of total CO₂ emissions from fossil fuel combustion. The industrial end-use sector accounted for 27 percent of CO₂ emissions from fossil fuel combustion. The residential and commercial end-use sectors accounted for 21 and 18 percent, respectively, of CO₂ emissions from fossil fuel combustion. Both of these end-use sectors were heavily reliant on electricity for meeting energy needs, with electricity consumption for lighting, heating, air conditioning, and operating appliances contributing 69 and 76 percent of emissions from the residential and commercial end-use sectors, respectively. Significant trends in emissions from energy source categories over the twenty four-year period from 1990 through 2013 included the following:

- Total CO₂ emissions from fossil fuel combustion increased from 4,740.7 MMT CO₂ Eq. in 1990 to 5,157.7 MMT CO₂ Eq. in 2013 – an 8.8 percent total increase over the twenty four-year period. From 2012 to 2013, these emissions increased by 131.7 MMT CO₂ Eq. (2.6 percent).
- CH₄ emissions from natural gas systems were the second largest anthropogenic source of CH₄ emissions in the United States with 157.4 MMT CO₂ Eq. emitted into the atmosphere in 2013; emissions have decreased by 21.8 MMT CO₂ Eq. (12.2 percent) since 1990.

- CO₂ emissions from non-energy use of fossil fuels increased by 2.2 MMT CO₂ Eq. (1.9 percent) from 1990 through 2013. Emissions from non-energy uses of fossil fuels were 119.8 MMT CO₂ Eq. in 2013, which constituted 2.2 percent of total national CO₂ emissions.
- N₂O emissions from stationary combustion increased by 11.0 MMT CO₂ Eq. (91.9 percent) from 1990 through 2013. N₂O emissions from this source increased primarily as a result of an increase in the number of coal fluidized bed boilers in the electric power sector.
- CO₂ emissions from incineration of waste (10.1 MMT CO₂ Eq. in 2013) increased by 2.2 MMT CO₂ Eq. (27.2 percent) from 1990 through 2013, as the volume of plastics and other fossil carbon-containing materials in municipal solid waste grew.

The increase in CO₂ emissions from fossil fuel combustion in 2013 was a result of multiple factors including: (1) the increase in the price of natural gas led to an increase of coal-fired generation in the electric power sector; (2) much colder winter conditions resulted in an increased demand for heating fuel in the residential and commercial sectors; (3) an increase in industrial production across multiple sectors which resulted in increases in industrial sector emissions,¹ and (4) an increase in transportation emissions resulting from a small increase in vehicle miles traveled (VMT) and fuel use across on-road transportation modes.

Industrial Processes and Product Use

The Industrial Processes and Product Use (IPPU) chapter includes greenhouse gas emissions occurring from industrial processes and from the use of greenhouse gases in products. This section includes sources of emissions formerly represented in the “Industrial Processes” and “Solvent and Other Product Use” sectors in prior versions of this report.

Greenhouse gas emissions are produced as the by-products of many non-energy-related industrial activities. For example, industrial processes can chemically transform raw materials, which often release waste gases such as CO₂, CH₄, and N₂O. These processes include iron and steel production and metallurgical coke production, cement production, ammonia production, urea consumption, lime production, other process uses of carbonates (e.g., flux stone, flue gas desulfurization, and glass manufacturing), soda ash production and consumption, titanium dioxide production, phosphoric acid production, ferroalloy production, CO₂ consumption, silicon carbide production and consumption, aluminum production, petrochemical production, nitric acid production, adipic acid production, lead production, zinc production, and N₂O from product uses (see Figure 2-9). Industrial processes also release HFCs, PFCs, SF₆, and NF₃. In addition to their use as ODS substitutes, HFCs, PFCs, SF₆, NF₃, and other fluorinated compounds are employed and emitted by a number of other industrial sources in the United States. These industries include aluminum production, HCFC-22 production, semiconductor manufacture, electric power transmission and distribution, and magnesium metal production and processing. Table 2-6 presents greenhouse gas emissions from industrial processes by source category.

¹ Further details on industrial sector combustion emissions are provided by EPA’s GHGRP (<http://ghgdata.epa.gov/ghgp/main.do>).

Figure 2-9: 2013 Industrial Processes and Product Use Chapter Greenhouse Gas Sources

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.




Table 2-6: Emissions from Industrial Processes and Product Use (MMT CO₂ Eq.)

Gas/Source	1990	2005	2009	2010	2011	2012	2013
CO₂	207.2	191.1	141.1	165.7	169.7	166.4	163.0
Iron and Steel Production & Metallurgical Coke Production	99.8	66.7	43.0	55.7	60.0	54.3	52.3
<i>Iron and Steel Production</i>	97.3	64.6	42.1	53.7	58.6	53.8	50.5
<i>Metallurgical Coke Production</i>	2.5	2.0	1.0	2.1	1.4	0.5	1.8
Cement Production	33.3	45.9	29.4	31.3	32.0	35.1	36.1
Petrochemical Production	21.6	28.1	23.7	27.4	26.4	26.5	26.5
Lime Production	11.7	14.6	11.4	13.4	14.0	13.7	14.1
Ammonia Production	13.0	9.2	8.5	9.2	9.3	9.4	10.2
Urea Consumption for Non-Agricultural Purposes	3.8	3.7	3.4	4.7	4.0	4.4	4.7
Other Process Uses of Carbonates	4.9	6.3	7.6	9.6	9.3	8.0	4.4
Aluminum Production	6.8	4.1	3.0	2.7	3.3	3.4	3.3
Soda Ash Production and Consumption	2.7	2.9	2.5	2.6	2.6	2.7	2.7
Ferroalloy Production	2.2	1.4	1.5	1.7	1.7	1.9	1.8
Titanium Dioxide Production	1.2	1.8	1.6	1.8	1.7	1.5	1.6
Zinc Production	0.6	1.0	0.9	1.2	1.3	1.5	1.4
Phosphoric Acid Production	1.6	1.4	1.0	1.1	1.2	1.1	1.2
Glass Production	1.5	1.9	1.0	1.5	1.3	1.2	1.2
Carbon Dioxide Consumption	1.5	1.4	1.8	1.2	0.8	0.8	0.9
Lead Production	0.5	0.6	0.5	0.5	0.5	0.5	0.5
Silicon Carbide Production and Consumption	0.4	0.2	0.1	0.2	0.2	0.2	0.2
Magnesium Production and Processing	+	+	+	+	+	+	+
CH₄	1.4	1.0	0.5	0.7	0.8	0.8	0.8

Iron and Steel Production & Metallurgical Coke							
Production	1.1	0.9	0.4	0.6	0.7	0.7	0.7
<i>Iron and Steel Production</i>	<i>1.1</i>	<i>0.9</i>	<i>0.4</i>	<i>0.6</i>	<i>0.7</i>	<i>0.7</i>	<i>0.7</i>
<i>Metallurgical Coke Production</i>	<i>+</i>						
Petrochemical Production	0.2	0.1	+	0.1	+	0.1	0.1
Ferroalloy Production	+	+	+	+	+	+	+
Silicon Carbide Production and Consumption	+	+	+	+	+	+	+
N₂O	31.6	22.8	16.7	20.1	25.5	20.4	19.1
Nitric Acid Production	12.1	11.3	9.6	11.5	10.9	10.5	10.7
N ₂ O from Product Uses	4.2	4.2	4.2	4.2	4.2	4.2	4.2
Adipic Acid Production	15.2	7.1	2.7	4.2	10.2	5.5	4.0
Semiconductor Manufacturing	+	0.1	0.1	0.1	0.2	0.2	0.2
HFCs	46.6	131.4	142.9	152.6	157.4	159.2	163.0
Substitution of Ozone Depleting Substances ^a	0.3	111.1	136.0	144.4	148.4	153.5	158.6
HCFC-22 Production	46.1	20.0	6.8	8.0	8.8	5.5	4.1
Semiconductor Manufacturing	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Magnesium Production and Processing	0	+	+	+	+	+	0.1
PFCs	24.3	6.6	3.9	4.4	6.9	6.0	5.8
Aluminum Production	21.5	3.4	1.9	1.9	3.5	2.9	3.0
Semiconductor Manufacturing	2.8	3.2	2.0	2.6	3.4	3.0	2.9
SF₆	31.1	14.0	9.3	9.5	10.0	7.7	6.9
Electrical Transmission and Distribution	25.4	10.6	7.3	7.0	6.8	5.7	5.1
Magnesium Production and Processing	5.2	2.7	1.6	2.1	2.8	1.6	1.4
Semiconductor Manufacturing	0.5	0.7	0.3	0.4	0.4	0.4	0.4
NF₃	+	0.5	0.4	0.5	0.7	0.6	0.6
Semiconductor Manufacturing	+	0.5	0.4	0.5	0.7	0.6	0.6
Total	342.1	367.4	314.9	353.6	371.0	361.2	359.1

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.

+ Does not exceed 0.05 MMT CO₂ Eq.

^a Small amounts of PFC emissions also result from this source.

Note: Totals may not sum due to independent rounding.

Overall, emissions from the IPPU sector increased by 5.0 percent from 1990 to 2013. Significant trends in emissions from IPPU source categories over the twenty four-year period from 1990 through 2013 included the following:

- HFC emissions from ODS substitutes have been increasing from small amounts in 1990 to 158.6 MMT CO₂ Eq. in 2013. This increase was in large part the result of efforts to phase out CFCs and other ODSs in the United States. In the short term, this trend is expected to continue, and will likely continue over the next decade as HCFCs, which are interim substitutes in many applications, are themselves phased-out under the provisions of the Copenhagen Amendments to the *Montreal Protocol*.
- Combined CO₂ and CH₄ emissions from iron and steel production and metallurgical coke production decreased by 3.8 percent to 53.0 MMT CO₂ Eq. from 2012 to 2013, and have declined overall by 47.9 MMT CO₂ Eq. (47.5 percent) from 1990 through 2013, due to restructuring of the industry, technological improvements, and increased scrap steel utilization.
- CO₂ emissions from ammonia production (10.2 MMT CO₂ Eq. in 2013) decreased by 2.9 MMT CO₂ Eq. (22.2 percent) since 1990. Ammonia production relies on natural gas as both a feedstock and a fuel, and as such, market fluctuations and volatility in natural gas prices affect the production of ammonia.
- Urea consumption for non-agricultural purposes (4.7 MMT CO₂ Eq. in 2013) increased by 0.9 MMT CO₂ Eq. (23.2 percent) since 1990.
- In 2013, N₂O emissions from product uses constituted 1.2 percent of U.S. N₂O emissions. From 1990 to 2013, emissions from this source category decreased by 0.4 percent, though slight increases occurred in intermediate years.

- N₂O emissions from adipic acid production were 4.0 MMT CO₂ Eq. in 2013, and have decreased significantly since 1990 due to both the widespread installation of pollution control measures in the late 1990s and plant idling in the late 2000s. Emissions from adipic acid production have decreased by 73.8 percent since 1990 and by 76.4 percent since a peak in 1995.
- PFC emissions from aluminum production decreased by 86.2 percent (18.5 MMT CO₂ Eq.) from 1990 to 2013, due to both industry emission reduction efforts and lower domestic aluminum production.

Agriculture

Agricultural activities contribute directly to emissions of greenhouse gases through a variety of processes, including the following source categories: enteric fermentation in domestic livestock, livestock manure management, rice cultivation, agricultural soil management, and field burning of agricultural residues.

In 2013, agricultural activities were responsible for emissions of 515.7 MMT CO₂ Eq., or 7.7 percent of total U.S. greenhouse gas emissions. CH₄ and N₂O were the primary greenhouse gases emitted by agricultural activities. CH₄ emissions from enteric fermentation and manure management represented about 25.9 percent and 9.6 percent of total CH₄ emissions from anthropogenic activities, respectively, in 2013. Agricultural soil management activities, such as fertilizer use and other cropping practices, were the largest source of U.S. N₂O emissions in 2013, accounting for 74.2 percent.

Figure 2-10: 2013 Agriculture Chapter Greenhouse Gas Sources

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.




Table 2-7: Emissions from Agriculture (MMT CO₂ Eq.)

Gas/Source	1990	2005	2009	2010	2011	2012	2013
CH₄	210.8	234.4	242.1	243.4	238.9	239.6	234.5
Enteric Fermentation	164.2	168.9	172.7	171.1	168.7	166.3	164.5
Manure Management	37.2	56.3	59.7	60.9	61.4	63.7	61.4
Rice Cultivation	9.2	8.9	9.4	11.1	8.5	9.3	8.3
Field Burning of Agricultural Residues	0.3	0.2	0.3	0.3	0.3	0.3	0.3
N₂O	237.9	260.1	281.2	281.4	283.2	283.4	281.1
Agricultural Soil Management	224.0	243.6	264.1	264.3	265.8	266.0	263.7
Manure Management	13.8	16.4	17.0	17.1	17.3	17.3	17.3
Field Burning of Agricultural Residues	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Total	448.7	494.5	523.3	524.8	522.1	523.0	515.7

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.

Note: Totals may not sum due to independent rounding.

Some significant trends in U.S. emissions from Agriculture source categories include the following:

- Agricultural soils produced approximately 74.2 percent of N₂O emissions in the United States in 2013. Estimated emissions from this source in 2013 were 263.7 MMT CO₂ Eq. Annual N₂O emissions from agricultural soils fluctuated between 1990 and 2013, although overall emissions were 17.7 percent higher in 2013 than in 1990. Year-to-year fluctuations are largely a reflection of annual variation in weather patterns, synthetic fertilizer use, and crop production.
- Enteric fermentation is the largest anthropogenic source of CH₄ emissions in the United States. In 2013, enteric fermentation CH₄ emissions were 164.5 MMT CO₂ Eq. (25.9 percent of total CH₄ emissions), which represents an increase of 0.4 MMT CO₂ Eq. (0.2 percent) since 1990. This increase in emissions from 1990 to 2013 in enteric generally follows the increasing trends in cattle populations. From 1990 to 1995 emissions increased and then generally decreased from 1996 to 2004, mainly due to fluctuations in beef cattle populations and increased digestibility of feed for feedlot cattle. Emissions increased from 2005 to 2007, as both dairy and beef populations underwent increases and the literature for dairy cow diets indicated a trend toward a decrease in feed digestibility for those years. Emissions decreased again from 2008 to 2013 as beef cattle populations again decreased.
- Overall, emissions from manure management increased 54.4 percent between 1990 and 2013. This encompassed an increase of 65.2 percent for CH₄, from 37.2 MMT CO₂ Eq. in 1990 to 61.4 MMT CO₂ Eq. in 2013; and an increase of 25.4 percent for N₂O, from 13.8 MMT CO₂ Eq. in 1990 to 17.3 MMT CO₂ Eq. in 2013. The majority of the increase observed in CH₄ resulted from swine and dairy cow manure, where emissions increased 48 and 115 percent, respectively, from 1990 to 2013. From 2012 to 2013, there was a 3.6 percent decrease in total CH₄ emissions from manure management, mainly due to minor shifts in the animal populations and the resultant effects on manure management system allocations.

Land Use, Land-Use Change, and Forestry

When humans alter the terrestrial biosphere through land use, changes in land use, and land management practices, they also alter the background carbon fluxes between biomass, soils, and the atmosphere. Forest management practices, tree planting in urban areas, the management of agricultural soils, and the landfilling of yard trimmings and food scraps have resulted in a net removal of CO₂ (sequestration of C) in the United States. Forests (including vegetation, soils, and harvested wood) accounted for approximately 88 percent of total 2013 CO₂ removals, urban trees accounted for 10 percent, mineral and organic soil carbon stock changes accounted for less than 0.5 percent, and landfilled yard trimmings and food scraps accounted for 1.4 percent of total CO₂ removals in 2013. The net forest sequestration is a result of net forest growth, increasing forest area, and a net accumulation of carbon stocks in harvested wood pools. The net sequestration in urban forests is a result of net tree growth and increased urban forest size. In agricultural soils, mineral and organic soils sequester approximately 2.4 times as much C as is emitted from these soils through liming and urea fertilization. The mineral soil C sequestration is largely due to the conversion of cropland to hay production fields, the limited use of bare-summer fallow areas in semi-arid areas, and an increase in the adoption of conservation tillage practices. The landfilled yard trimmings and food scraps net sequestration is due to the long-term accumulation of yard trimming and food scraps carbon in landfills.

Land use, land-use change, and forestry activities in 2013 resulted in a C sequestration (i.e., total sinks) of 881.7 MMT CO₂ Eq. (Table 2-3).² This represents an offset of approximately 13.2 percent of total (i.e., gross) greenhouse gas emissions in 2013. Emissions from land use, land-use change and forestry activities in 2013 represent 0.3 percent of total greenhouse gas emissions.³ Between 1990 and 2013, total land use, land-use change, and forestry C sequestration increased by 13.6 percent, primarily due to an increase in the rate of net C accumulation in forest C stocks, particularly in aboveground and belowground tree biomass, and harvested wood pools.

² The total sinks value includes the positive C sequestration reported for *Forest Land Remaining Forest Land*, *Cropland Remaining Cropland*, *Land Converted to Grassland*, *Settlements Remaining Settlements*, and *Other Land* plus the loss in C sequestration reported for *Land Converted to Cropland* and *Grassland Remaining Grassland*.

³ The emissions value includes the CO₂, CH₄, and N₂O emissions reported for *Forest Fires*, *Forest Soils*, *Liming of Agricultural Soils*, *Urea Fertilization*, *Settlement Soils*, and *Peatlands Remaining Peatlands*.

CO₂ removals are presented in Table 2-8 along with CO₂, CH₄, and N₂O emissions for Land Use, Land-Use Change, and Forestry source categories. Liming of agricultural soils and urea fertilization resulted in CO₂ emissions of 9.9 MMT CO₂ Eq. in 2013, an increase of about 40.3 percent relative to 1990. Lands undergoing peat extraction (i.e., *Peatlands Remaining Peatlands*) resulted in CO₂ emissions of 0.8 MMT CO₂ Eq. and CH₄ and N₂O emissions of less than 0.05 MMT CO₂ Eq. each. N₂O emissions from the application of synthetic fertilizers to forest soils have increased from 0.1 MMT CO₂ Eq. in 1990 to 0.5 MMT CO₂ Eq. in 2013. Settlement soils in 2013 resulted in N₂O emissions of 2.4 MMT CO₂ Eq., a 76.7 percent increase relative to 1990. Emissions from forest fires in 2013 resulted in CH₄ emissions of 5.8 MMT CO₂ and in N₂O emissions of 3.8 MMT CO₂ (see Table 2-8).

Table 2-8: Emissions and Removals (Flux) from Land Use, Land-Use Change, and Forestry (MMT CO₂ Eq.)

Gas/Land-Use Category	1990	2005	2009	2010	2011	2012	2013
CO₂	(767.7)	(903.0)	(862.6)	(862.0)	(872.1)	(869.6)	(871.0)
Forest Land Remaining Forest Land:							
Changes in Forest Carbon Stock ^a	(639.4)	(807.1)	(764.9)	(765.4)	(773.8)	(773.1)	(775.7)
Cropland Remaining Cropland:							
Changes in Agricultural Soil Carbon Stock	(65.2)	(28.0)	(27.5)	(25.9)	(25.8)	(25.0)	(23.4)
Cropland Remaining Cropland:							
Liming of Agricultural Soils	4.7	4.3	3.7	4.8	3.9	5.8	5.9
Cropland Remaining Cropland:							
Urea Fertilization	2.4	3.5	3.6	3.8	4.1	4.2	4.0
Land Converted to Cropland	24.5	19.8	16.2	16.2	16.2	16.1	16.1
Grassland Remaining Grassland	(1.9)	4.2	11.7	11.7	11.7	11.5	12.1
Land Converted to Grassland	(7.4)	(9.0)	(8.9)	(8.9)	(8.9)	(8.8)	(8.8)
Settlements Remaining Settlements:							
Changes in Urban Tree Carbon Stock ^b	(60.4)	(80.5)	(85.0)	(86.1)	(87.3)	(88.4)	(89.5)
Wetlands Remaining Wetlands:							
Peatlands Remaining Peatlands	1.0	1.1	1.0	1.0	0.9	0.8	0.8
Other:							
Landfilled Yard Trimmings and Food Scraps	(24.2)	(12.0)	(12.9)	(13.6)	(13.5)	(13.0)	(12.8)
CH₄	2.5	8.3	5.8	4.8	14.6	15.7	5.8
Forest Land Remaining Forest Land:							
Forest Fires	2.5	8.3	5.8	4.7	14.6	15.7	5.8
Wetlands Remaining Wetlands:							
Peatlands Remaining Peatlands	+	+	+	+	+	+	+
N₂O	3.1	8.3	6.5	6.0	12.6	13.3	6.7
Forest Land Remaining Forest Land:							
Forest Fires	1.7	5.5	3.8	3.1	9.6	10.3	3.8
Forest Land Remaining Forest Land:							
Forest Soils ^c	0.1	0.5	0.5	0.5	0.5	0.5	0.5
Settlements Remaining Settlements:							
Settlement Soils ^d	1.4	2.3	2.2	2.4	2.5	2.5	2.4
Wetlands Remaining Wetlands:							
Peatlands Remaining Peatlands	+	+	+	+	+	+	+
Total Flux^e	(762.1)	(886.4)	(850.2)	(851.3)	(844.9)	(840.6)	(858.5)

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.

+ Less than 0.05 MMT CO₂ Eq.

^a Estimates include C stock changes on both *Forest Land Remaining Forest Land* and *Land Converted to Forest Land*.

^b Estimates include C stock changes on both *Settlements Remaining Settlements* and *Land Converted to Settlements*.

^c Estimates include emissions from N fertilizer additions on both *Forest Land Remaining Forest Land*, and *Land Converted to Forest Land*, but not from land-use conversion.

^d Estimates include emissions from N fertilizer additions on both *Settlements Remaining Settlements*, and *Land Converted to Settlements*, but not from land-use conversion.

^e “Total Flux” is defined as the sum of positive emissions (i.e., sources) of greenhouse gases to the atmosphere plus removals of CO₂ (i.e., sinks or negative emissions) from the atmosphere.

Note: Totals may not sum due to independent rounding. Parentheses indicate net sequestration.

Other significant trends from 1990 to 2013 in emissions from land use, land-use change, and forestry source categories include:


- Annual C sequestration by forest land (i.e., annual carbon stock accumulation in the five carbon pools) has increased by approximately 21 percent. This is primarily due to increased forest management and the effects of previous reforestation. The increase in intensive forest management resulted in higher growth rates and higher biomass density. The tree planting and conservation efforts of the 1970s and 1980s continue to have a significant impact on sequestration rates. Finally, the forested area in the United States increased over the past twenty four-years, although only at an average rate of 0.1 percent per year.
- Annual C sequestration by urban trees has increased by 48.1 percent over the period from 1990 to 2013. This is primarily due to an increase in urbanized land area in the United States.
- Annual C sequestration in landfilled yard trimmings and food scraps has decreased by 51.6 percent since 1990. Food scrap generation has grown by 53 percent since 1990, and though the proportion of food scraps discarded in landfills has decreased slightly from 82 percent in 1990 to 78 percent in 2013, the tonnage disposed in landfills has increased considerably (by 46 percent). Overall, the decrease in the landfill disposal rate of yard trimmings has more than compensated for the increase in food scrap disposal in landfills.

Waste

Waste management and treatment activities are sources of greenhouse gas emissions (see Figure 2-11). In 2013, landfills were the third largest source of U.S. anthropogenic CH₄ emissions, accounting for 18.0 percent of total U.S. CH₄ emissions.⁴ Additionally, wastewater treatment accounts for 14.4 percent of Waste emissions, 2.4 percent of U.S. CH₄ emissions, and 1.4 percent of N₂O emissions. Emissions of CH₄ and N₂O from composting grew from 1990 to 2013, and resulted in emissions of 3.7 MMT CO₂ Eq. in 2013. A summary of greenhouse gas emissions from the Waste chapter is presented in Table 2-9.

Figure 2-11: 2013 Waste Chapter Greenhouse Gas Sources

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.



⁴ Landfills also store carbon, due to incomplete degradation of organic materials such as wood products and yard trimmings, as described in the Land Use, Land-Use Change, and Forestry chapter.

Overall, in 2013, waste activities generated emissions of 138.3 MMT CO₂ Eq., or 2.1 percent of total U.S. greenhouse gas emissions.

Table 2-9: Emissions from Waste (MMT CO₂ Eq.)

Gas/Source	1990	2005	2009	2010	2011	2012	2013
CH₄	202.3	183.2	175.5	139.1	138.4	132.4	131.6
Landfills	186.2	165.5	158.1	121.8	121.3	115.3	114.6
Wastewater Treatment	15.7	15.9	15.6	15.5	15.3	15.2	15.0
Composting	0.4	1.9	1.9	1.8	1.9	1.9	2.0
N₂O	3.7	6.0	6.3	6.4	6.5	6.6	6.7
Wastewater Treatment	3.4	4.3	4.6	4.7	4.8	4.9	4.9
Composting	0.3	1.7	1.7	1.6	1.7	1.7	1.8
Total	206.0	189.2	181.8	145.5	144.9	138.9	138.3

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.

Totals may not sum due to independent rounding.

Some significant trends in U.S. emissions from waste source categories include the following:

- From 1990 to 2013, net CH₄ emissions from landfills decreased by 71.6 MMT CO₂ Eq. (38.4 percent), with small increases occurring in interim years. This downward trend in overall emissions is the result of increases in the amount of landfill gas collected and combusted as well as reduction in the amount of decomposable materials (i.e., paper and paperboard, food scraps, and yard trimmings) discarded in MSW landfills over the time series,⁵ which has more than offset the additional CH₄ emissions resulting from an increase in the amount of municipal solid waste landfilled.
- Combined CH₄ and N₂O emissions from composting have generally increased since 1990, from 0.7 MMT CO₂ Eq. to 3.7 MMT CO₂ Eq. in 2013, which represents slightly more than a five-fold increase over the time series. The growth in composting since the 1990s is attributable to primarily two factors: (1) steady growth in population and residential housing, and (2) the enactment of legislation by state and local governments that discouraged the disposal of yard trimmings in landfills.
- From 1990 to 2013, CH₄ and N₂O emissions from wastewater treatment decreased by 0.6 MMT CO₂ Eq. (4.0 percent) and increased by 1.6 MMT CO₂ Eq. (46.5 percent), respectively. Methane emissions from domestic wastewater treatment have decreased since 1999 due to decreasing percentages of wastewater being treated in anaerobic systems, including reduced use of on-site septic systems and central anaerobic treatment systems. Nitrous oxide emissions from wastewater treatment processes gradually increased across the time series as a result of increasing U.S. population and protein consumption.

2.1 Emissions by Economic Sector

Throughout this report, emission estimates are grouped into six sectors (i.e., chapters) defined by the IPCC and detailed above: Energy; Industrial Processes; Solvent and Other Product Use; Agriculture; Land Use, Land-Use Change, and Forestry; and Waste. While it is important to use this characterization for consistency with UNFCCC reporting guidelines, it is also useful to allocate emissions into more commonly used sectoral categories. This section reports emissions by the following U.S. economic sectors: residential, commercial, industry, transportation, electricity generation, and agriculture, as well as U.S. Territories.

Using this categorization, emissions from electricity generation accounted for the largest portion (31 percent) of U.S. greenhouse gas emissions in 2013. Transportation activities, in aggregate, accounted for the second largest portion (27 percent). Emissions from industry accounted for about 21 percent of U.S. greenhouse gas emissions in

⁵ The CO₂ produced from combusted landfill CH₄ at landfills is not counted in national inventories as it is considered part of the natural C cycle of decomposition.

2013. In contrast to electricity generation and transportation, emissions from industry have in general declined over the past decade. The long-term decline in these emissions has been due to structural changes in the U.S. economy (i.e., shifts from a manufacturing-based to a service-based economy), fuel switching, and efficiency improvements. The remaining 21 percent of U.S. greenhouse gas emissions were contributed by the residential, agriculture, and commercial sectors, plus emissions from U.S. Territories. The residential sector accounted for 6 percent, and primarily consisted of CO₂ emissions from fossil fuel combustion. Activities related to agriculture accounted for roughly 9 percent of U.S. emissions; unlike other economic sectors, agricultural sector emissions were dominated by N₂O emissions from agricultural soil management and CH₄ emissions from enteric fermentation, rather than CO₂ from fossil fuel combustion. The commercial sector accounted for roughly 6 percent of emissions, while U.S. territories accounted for less than 1 percent. Carbon dioxide was also emitted and sequestered (in the form of C) by a variety of activities related to forest management practices, tree planting in urban areas, the management of agricultural soils, and landfilling of yard trimmings.

Table 2-10 presents a detailed breakdown of emissions from each of these economic sectors by source category, as they are defined in this report. Figure 2-12 shows the trend in emissions by sector from 1990 to 2013.

Figure 2-12: Emissions Allocated to Economic Sectors

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.




Table 2-10: U.S. Greenhouse Gas Emissions Allocated to Economic Sectors (MMT CO₂ Eq. and Percent of Total in 2013)

Sector/Source	1990	2005	2009	2010	2011	2012	2013	Percent ^a
Electric Power Industry	1,864.8	2,443.9	2,185.7	2,300.5	2,198.1	2,060.8	2,077.0	31.1%
CO ₂ from Fossil Fuel Combustion	1,820.8	2,400.9	2,145.7	2,258.4	2,157.7	2,022.2	2,039.8	30.6%
Stationary Combustion	7.7	16.5	17.2	18.9	18.0	18.2	19.5	0.3%
Incineration of Waste	8.4	12.8	11.6	11.4	10.9	10.7	10.4	0.2%
Electrical Transmission and Distribution	25.4	10.6	7.3	7.0	6.8	5.7	5.1	0.1%
Other Process Uses of Carbonates	2.5	3.2	3.8	4.8	4.7	4.0	2.2	+
Transportation	1,551.3	2,017.7	1,835.3	1,843.5	1,815.4	1,795.9	1,806.2	27.1%
CO ₂ from Fossil Fuel Combustion	1,493.8	1,887.8	1,720.3	1,732.0	1,711.5	1,700.8	1,718.4	25.8%
Substitution of Ozone Depleting Substances	+	80.4	81.4	77.9	72.0	66.3	60.5	0.9%
Mobile Combustion	45.7	39.4	25.1	24.1	22.9	20.5	18.6	0.3%
Non-Energy Use of Fuels	11.8	10.2	8.5	9.5	9.0	8.3	8.8	0.1%
Industry	1,587.7	1,462.8	1,272.5	1,353.3	1,353.0	1,338.9	1,392.1	20.9%
CO ₂ from Fossil Fuel Combustion	811.4	781.0	681.2	728.2	724.9	733.5	767.6	11.5%
Natural Gas Systems	216.8	206.3	200.2	191.9	194.8	189.2	195.2	2.9%
Non-Energy Use of Fuels	100.1	120.6	93.5	100.9	95.8	93.3	108.4	1.6%

Coal Mining	96.5	64.1	79.9	82.3	71.2	66.5	64.6	1.0%
Iron and Steel Production	100.9	67.5	43.5	56.4	60.7	55.1	53.0	0.8%
Cement Production	33.3	45.9	29.4	31.3	32.0	35.1	36.1	0.5%
Petroleum Systems	36.0	28.4	26.2	25.5	26.4	28.3	31.2	0.5%
Petrochemical Production	21.9	28.3	23.8	27.4	26.4	26.5	26.6	0.4%
Substitution of Ozone Depleting Substances	+	7.3	12.4	15.3	17.0	18.7	20.4	0.3%
Lime Production	11.7	14.6	11.4	13.4	14.0	13.7	14.1	0.2%
Nitric Acid Production	12.1	11.3	9.6	11.5	10.9	10.5	10.7	0.2%
Ammonia Production	13.0	9.2	8.5	9.2	9.3	9.4	10.2	0.2%
Abandoned Underground Coal Mines	7.2	6.6	6.4	6.6	6.4	6.2	6.2	0.1%
Aluminum Production	28.3	7.6	4.9	4.6	6.8	6.4	6.2	0.1%
Urea Consumption for Non-Agricultural Purposes	3.8	3.7	3.4	4.7	4.0	4.4	4.7	0.1%
N ₂ O from Product Uses	4.2	4.2	4.2	4.2	4.2	4.2	4.2	0.1%
Semiconductor Manufacture	3.6	4.7	3.1	3.8	4.9	4.5	4.2	0.1%
HCFC-22 Production	46.1	20.0	6.8	8.0	8.8	5.5	4.1	0.1%
Adipic Acid Production	15.2	7.1	2.7	4.2	10.2	5.5	4.0	0.1%
Stationary Combustion	4.9	4.6	3.6	3.9	3.9	3.9	3.9	0.1%
Soda Ash Production and Consumption	2.7	2.9	2.5	2.6	2.6	2.7	2.7	+
Other Process Uses of Carbonates	2.5	3.2	3.8	4.8	4.7	4.0	2.2	+
Ferroalloy Production	2.2	1.4	1.5	1.7	1.7	1.9	1.8	+
Titanium Dioxide Production	1.2	1.8	1.6	1.8	1.7	1.5	1.6	+
Magnesium Production and Processing	5.2	2.7	1.6	2.1	2.8	1.7	1.5	+
Mobile Combustion	0.9	1.3	1.3	1.4	1.4	1.4	1.5	+
Zinc Production	0.6	1.0	0.9	1.2	1.3	1.5	1.4	+
Phosphoric Acid Production	1.6	1.4	1.0	1.1	1.2	1.1	1.2	+
Glass Production	1.5	1.9	1.0	1.5	1.3	1.2	1.2	+
Carbon Dioxide Consumption	1.5	1.4	1.8	1.2	0.8	0.8	0.9	+
Lead Production	0.5	0.6	0.5	0.5	0.5	0.5	0.5	+
Silicon Carbide Production and Consumption	0.4	0.2	0.2	0.2	0.2	0.2	0.2	+
Agriculture	492.5	565.0	588.8	590.8	605.5	611.6	586.8	8.8%
N ₂ O from Agricultural Soil Management	224.0	243.6	264.1	264.3	265.8	266.0	263.7	4.0%
Enteric Fermentation	164.2	168.9	172.7	171.1	168.7	166.3	164.5	2.5%
Manure Management	51.0	72.8	76.7	78.0	78.7	81.0	78.7	1.2%
CO ₂ from Fossil Fuel Combustion	31.0	46.8	46.5	47.5	49.2	50.7	49.7	0.7%
CH ₄ and N ₂ O from Forest Fires	4.2	13.8	9.7	7.9	24.2	26.0	9.7	0.1%
Rice Cultivation	9.2	8.9	9.4	11.1	8.5	9.3	8.3	0.1%
Liming of Agricultural Soils	4.7	4.3	3.7	4.8	3.9	5.8	5.9	0.1%
Urea Fertilization	2.4	3.5	3.6	3.8	4.1	4.2	4.0	0.1%
CO ₂ , CH ₄ and N ₂ O from Managed Peatlands	1.1	1.1	1.0	1.0	0.9	0.8	0.8	+
Mobile Combustion	0.3	0.5	0.5	0.5	0.5	0.6	0.5	+
Field Burning of Agricultural Residues	0.4	0.3	0.4	0.4	0.4	0.4	0.4	+
N ₂ O from Forest Soils	0.1	0.5	0.5	0.5	0.5	0.5	0.5	+
Stationary Combustion	+	+	+	+	+	+	+	+
Commercial	424.8	429.8	431.9	396.4	400.7	374.3	401.1	6.0%
CO ₂ from Fossil Fuel Combustion	217.4	223.5	223.5	220.2	221.0	197.1	220.7	3.3%
Landfills	186.2	165.5	158.1	121.8	121.3	115.3	114.6	1.7%
Substitution of Ozone Depleting Substances	+	15.7	25.2	29.3	33.4	37.1	40.8	0.6%
Wastewater Treatment	15.7	15.9	15.6	15.5	15.3	15.2	15.0	0.2%
Human Sewage	3.4	4.3	4.6	4.7	4.8	4.9	4.9	0.1%
Composting	0.7	3.5	3.6	3.5	3.5	3.7	3.7	0.1%
Stationary Combustion	1.4	1.4	1.4	1.4	1.4	1.2	1.3	+
Residential	346.3	372.8	360.9	363.7	360.5	321.5	375.0	5.6%
CO ₂ from Fossil Fuel Combustion	338.3	357.8	336.4	334.7	327.2	283.1	329.6	4.9%

Substitution of Ozone Depleting Substances	0.3	7.7	17.0	21.8	25.9	31.4	37.0	0.6%
Stationary Combustion	6.3	4.9	5.3	4.8	4.9	4.5	5.9	0.1%
Settlement Soil Fertilization	1.4	2.3	2.2	2.4	2.5	2.5	2.4	+
U.S. Territories	33.7	58.2	47.6	50.6	43.5	42.1	34.8	0.5%
CO ₂ from Fossil Fuel Combustion	27.9	49.9	43.5	46.2	39.8	38.6	32.0	0.5%
Non-Energy Use of Fuels	5.7	8.1	3.9	4.2	3.6	3.3	2.7	+
Stationary Combustion	0.1	0.2	0.2	0.2	0.2	0.2	0.1	+
Total Emissions	6,301.1	7,350.2	6,722.7	6,898.8	6,776.6	6,545.1	6,673.0	100.0%
Sinks	(775.8)	(911.9)	(870.9)	(871.6)	(881.0)	(880.4)	(881.7)	-13.2%
CO ₂ Flux from Forests ^b	(639.4)	(807.1)	(764.9)	(765.4)	(773.8)	(773.1)	(775.7)	-11.6%
Urban Trees	(60.4)	(80.5)	(85.0)	(86.1)	(87.3)	(88.4)	(89.5)	-1.3%
Landfilled Yard Trimmings and Food Scraps	(26.0)	(11.4)	(12.5)	(13.2)	(13.2)	(12.8)	(12.6)	-0.2%
CO ₂ Flux from Agricultural Soil Carbon Stocks	(50.0)	(13.0)	(8.5)	(6.9)	(6.7)	(6.1)	(4.0)	-0.1%
Net Emissions	5,525.2	6,438.3	5,851.9	6,027.2	5,895.6	5,664.7	5,791.2	86.8%

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.

Note: Includes all emissions of CO₂, CH₄, N₂O, HFCs, PFCs, SF₆, and NF₃. Parentheses indicate negative values or sequestration. Totals may not sum due to independent rounding.

+ Does not exceed 0.05 MMT CO₂ Eq. or 0.05 percent.

^a Percent of total emissions for year 2013.

^b Includes the effects of net additions to stocks of carbon stored in harvested wood products.

Emissions with Electricity Distributed to Economic Sectors

It can also be useful to view greenhouse gas emissions from economic sectors with emissions related to electricity generation distributed into end-use categories (i.e., emissions from electricity generation are allocated to the economic sectors in which the electricity is consumed). The generation, transmission, and distribution of electricity, which is the largest economic sector in the United States, accounted for 31 percent of total U.S. greenhouse gas emissions in 2013. Emissions increased by 11 percent since 1990, as electricity demand grew and fossil fuels remained the dominant energy source for generation. Electricity generation-related emissions increased from 2012 to 2013 by 0.8 percent, primarily due to increased CO₂ emissions from fossil fuel combustion. Electricity sales to the residential and commercial end-use sectors in 2013 increased approximately 1.2 percent and 0.9 percent, respectively. The trend in the residential and commercial sectors can largely be attributed to colder more energy-intensive winter conditions compared to 2012. Electricity sales to the industrial sector in 2013 decreased by approximately 3.1 percent. Overall, in 2013, the amount of electricity generated (in kWh) decreased by 0.1 percent from the previous year. Despite the decrease in generation, CO₂ emissions from the electric power sector increased by 0.8 percent as the consumption of CO₂ intensive coal and petroleum for electricity generation increased by 4.2 percent and 18.8 percent, respectively, in 2013 and the consumption of natural gas for electricity generation, decreased by 10.2 percent. Table 2-11 provides a detailed summary of emissions from electricity generation-related activities.

Table 2-11: Electricity Generation-Related Greenhouse Gas Emissions (MMT CO₂ Eq.)

Gas/Fuel Type or Source	1990	2005	2009	2010	2011	2012	2013
CO₂	1,831.2	2,416.5	2,160.7	2,274.2	2,172.9	2,036.6	2,052.1
Fossil Fuel Combustion	1,820.8	2,400.9	2,145.7	2,258.4	2,157.7	2,022.2	2,039.8
<i>Coal</i>	<i>1,547.6</i>	<i>1,983.8</i>	<i>1,740.9</i>	<i>1,827.6</i>	<i>1,722.7</i>	<i>1,511.2</i>	<i>1,575.0</i>
<i>Natural Gas</i>	<i>175.3</i>	<i>318.8</i>	<i>372.2</i>	<i>399.0</i>	<i>408.8</i>	<i>492.2</i>	<i>441.9</i>
<i>Petroleum</i>	<i>97.5</i>	<i>97.9</i>	<i>32.2</i>	<i>31.4</i>	<i>25.8</i>	<i>18.3</i>	<i>22.4</i>
<i>Geothermal</i>	<i>0.4</i>						
Incineration of Waste	8.0	12.5	11.3	11.0	10.5	10.4	10.1
Other Process Uses of Carbonates	2.5	3.2	3.8	4.8	4.7	4.0	2.2
CH₄	0.3	0.5	0.4	0.5	0.4	0.4	0.4

Stationary Combustion ^a	0.3	0.5	0.4	0.5	0.4	0.4	0.4
Incineration of Waste	+	+	+	+	+	+	+
N₂O	7.8	16.4	17.1	18.8	17.9	18.1	19.4
Stationary Combustion ^a	7.4	16.0	16.8	18.5	17.6	17.8	19.1
Incineration of Waste	0.5	0.4	0.3	0.3	0.3	0.3	0.3
SF₆	25.4	10.6	7.3	7.0	6.8	5.7	5.1
Electrical Transmission and Distribution	25.4	10.6	7.3	7.0	6.8	5.7	5.1
Total	1,864.8	2,443.9	2,185.7	2,300.5	2,198.1	2,060.8	2,077.0

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.

Note: Totals may not sum due to independent rounding.

^a Includes only stationary combustion emissions related to the generation of electricity.

+ Does not exceed 0.05 MMT CO₂ Eq.


To distribute electricity emissions among economic end-use sectors, emissions from the source categories assigned to the electricity generation sector were allocated to the residential, commercial, industry, transportation, and agriculture economic sectors according to each economic sector's share of retail sales of electricity consumption (EIA 2015, Duffield 2006). These source categories include CO₂ from Fossil Fuel Combustion, CH₄ and N₂O from Stationary Combustion, Incineration of Waste, Other Process Uses of Carbonates, and SF₆ from Electrical Transmission and Distribution Systems. Note that only 50 percent of the Other Process Uses of Carbonates emissions were associated with electricity generation and distributed as described; the remainder of Other Process Uses of Carbonates emissions were attributed to the industrial processes economic end-use sector.⁶

When emissions from electricity are distributed among these sectors, industrial activities account for the largest share of total U.S. greenhouse gas emissions (28.8 percent), followed closely by emissions from transportation (27.1 percent). Emissions from the residential and commercial sectors also increase substantially when emissions from electricity are included. In all sectors except agriculture, CO₂ accounts for more than 80 percent of greenhouse gas emissions, primarily from the combustion of fossil fuels.

Table 2-12 presents a detailed breakdown of emissions from each of these economic sectors, with emissions from electricity generation distributed to them. Figure 2-13 shows the trend in these emissions by sector from 1990 to 2013.

Figure 2-13: Emissions with Electricity Distributed to Economic Sectors

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.



⁶ Emissions were not distributed to U.S. Territories, since the electricity generation sector only includes emissions related to the generation of electricity in the 50 states and the District of Columbia.

Table 2-12: U.S. Greenhouse Gas Emissions by Economic Sector and Gas with Electricity-Related Emissions Distributed (MMT CO₂ Eq.) and Percent of Total in 2013

Sector/Gas	1990	2005	2009	2010	2011	2012	2013	Percent ^a
Industry	2,229.7	2,148.5	1,817.7	1,937.7	1,923.9	1,880.9	1,922.6	28.8%
Direct Emissions	1,587.7	1,462.8	1,272.5	1,353.3	1,353.0	1,338.9	1,392.1	20.9%
CO ₂	1,158.3	1,124.5	948.9	1,026.5	1,025.7	1,029.0	1,080.6	16.2%
CH ₄	317.7	273.4	277.8	272.2	261.3	252.9	255.9	3.8%
N ₂ O	35.3	26.6	19.9	23.6	28.9	23.8	22.5	0.3%
HFCs, PFCs, SF ₆ , and NF ₃	76.3	38.3	25.8	31.1	37.0	33.3	33.1	0.5%
Electricity-Related	642.0	685.7	545.2	584.4	571.0	542.0	530.5	8.0%
CO ₂	630.4	678.0	539.0	577.7	564.4	535.6	524.2	7.9%
CH ₄	0.1	0.1	0.1	0.1	0.1	0.1	0.1	+
N ₂ O	2.7	4.6	4.3	4.8	4.7	4.8	5.0	0.1%
SF ₆	8.7	3.0	1.8	1.8	1.8	1.5	1.3	+
Transportation	1,554.4	2,022.5	1,839.9	1,848.1	1,819.7	1,799.8	1,810.3	27.1%
Direct Emissions	1,551.3	2,017.7	1,835.3	1,843.5	1,815.4	1,795.9	1,806.2	27.1%
CO ₂	1,505.6	1,898.0	1,728.9	1,741.5	1,720.5	1,709.1	1,727.2	25.9%
CH ₄	5.4	2.7	2.0	1.9	1.9	1.8	1.7	+
N ₂ O	40.26	36.70	23.12	22.20	20.98	18.68	16.84	0.3%
HFCs ^b	+	80.4	81.4	77.9	72.0	66.3	60.5	0.9%
Electricity-Related	3.1	4.8	4.6	4.6	4.3	3.9	4.1	0.1%
CO ₂	3.1	4.8	4.5	4.5	4.3	3.9	4.0	0.1%
CH ₄	+	+	+	+	+	+	+	+
N ₂ O	+	+	+	+	+	+	+	+
SF ₆	+	+	+	+	+	+	+	+
Commercial	975.8	1,247.5	1,199.2	1,183.8	1,152.6	1,088.0	1,126.7	16.9%
Direct Emissions	424.8	429.8	431.9	396.4	400.7	374.3	401.1	6.0%
CO ₂	217.4	223.5	223.5	220.2	221.0	197.1	220.7	3.3%
CH ₄	203.3	184.3	176.6	140.2	139.4	133.3	132.7	2.0%
N ₂ O	4.1	6.3	6.6	6.7	6.8	6.8	7.0	0.1%
HFCs	+	15.7	25.2	29.3	33.4	37.1	40.8	0.6%
Electricity-Related	551.0	817.7	767.2	787.4	751.9	713.6	725.6	10.9%
CO ₂	541.1	808.5	758.5	778.4	743.3	705.2	716.9	10.7%
CH ₄	0.1	0.2	0.2	0.2	0.1	0.1	0.1	+
N ₂ O	2.3	5.5	6.0	6.4	6.1	6.3	6.8	0.1%
SF ₆	7.5	3.5	2.6	2.4	2.3	2.0	1.8	+
Residential	953.6	1,244.4	1,161.8	1,219.5	1,166.0	1,060.6	1,129.1	16.9%
Direct Emissions	346.3	372.8	360.9	363.7	360.5	321.5	375.0	5.6%
CO ₂	338.3	357.8	336.4	334.7	327.2	283.1	329.6	4.9%
CH ₄	5.2	4.1	4.4	4.0	4.0	3.7	5.0	0.1%
N ₂ O	2.4	3.2	3.2	3.2	3.3	3.3	3.4	0.1%
HFCs	0.3	7.7	17.0	21.8	25.9	31.4	37.0	0.6%
Electricity-Related	607.3	871.6	800.9	855.8	805.5	739.1	754.2	11.3%
CO ₂	596.4	861.9	791.7	846.0	796.3	730.4	745.1	11.2%
CH ₄	0.1	0.2	0.2	0.2	0.2	0.2	0.2	+
N ₂ O	2.5	5.8	6.3	7.0	6.6	6.5	7.1	0.1%
SF ₆	8.3	3.8	2.7	2.6	2.5	2.0	1.9	+
Agriculture	553.9	629.1	656.6	659.2	670.9	673.7	649.4	9.7%
Direct Emissions	492.5	565.0	588.8	590.8	605.5	611.6	586.8	8.8%
CO ₂	39.2	55.7	54.8	57.1	58.1	61.5	60.4	0.9%
CH ₄	213.4	242.9	248.2	248.4	253.7	255.5	240.6	3.6%
N ₂ O	239.9	266.4	285.9	285.4	293.6	294.6	285.8	4.3%
Electricity-Related	61.3	64.1	67.8	68.4	65.4	62.1	62.6	0.9%
CO ₂	60.2	63.4	67.0	67.6	64.7	61.4	61.9	0.9%
CH ₄	+	+	+	+	+	+	+	+
N ₂ O	0.3	0.4	0.5	0.6	0.5	0.5	0.6	+
SF ₆	0.8	0.3	0.2	0.2	0.2	0.2	0.2	+

U.S. Territories	33.7	58.2	47.6	50.6	43.5	42.1	34.8	0.5%
Total	6,301.1	7,350.2	6,722.7	6,898.8	6,776.6	6,545.1	6,673.0	100.0%

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.

Note: Emissions from electricity generation are allocated based on aggregate electricity consumption in each end-use sector.

Totals may not sum due to independent rounding.

+ Does not exceed 0.05 MMT CO₂ Eq. or 0.05 percent.

^a Percent of total emissions for year 2013.

^b Includes primarily HFC-134a.

Industry

The industrial end-use sector includes CO₂ emissions from fossil fuel combustion from all manufacturing facilities, in aggregate. This sector also includes emissions that are produced as a byproduct of the non-energy-related industrial process activities. The variety of activities producing these non-energy-related emissions includes methane emissions from petroleum and natural gas systems, fugitive CH₄ emissions from coal mining, by-product CO₂ emissions from cement manufacture, and HFC, PFC, SF₆, and NF₃ byproduct emissions from semiconductor manufacture, to name a few. Since 1990, industrial sector emissions have declined. The decline has occurred both in direct emissions and indirect emissions associated with electricity use. In theory, emissions from the industrial end-use sector should be highly correlated with economic growth and industrial output, but heating of industrial buildings and agricultural energy consumption are also affected by weather conditions. In addition, structural changes within the U.S. economy that lead to shifts in industrial output away from energy-intensive manufacturing products to less energy-intensive products (e.g., from steel to computer equipment) also have a significant effect on industrial emissions.

Transportation

When electricity-related emissions are distributed to economic end-use sectors, transportation activities accounted for 27 percent of U.S. greenhouse gas emissions in 2013. The largest sources of transportation greenhouse gases in 2013 were passenger cars (42.2 percent), freight trucks (22.5 percent), light duty trucks, which include sport utility vehicles, pickup trucks, and minivans (17.9 percent), commercial aircraft (6.4 percent), rail (2.6 percent), pipelines (2.6 percent), and ships and boats (2.2 percent). These figures include direct CO₂, CH₄, and N₂O emissions from fossil fuel combustion used in transportation and emissions from non-energy use (i.e., lubricants) used in transportation, as well as HFC emissions from mobile air conditioners and refrigerated transport allocated to these vehicle types.

In terms of the overall trend, from 1990 to 2013, total transportation emissions rose by 16.5 percent due, in large part, to increased demand for travel as fleetwide light-duty vehicle fuel economy was relatively stable (average new vehicle fuel economy declined slowly from 1990 through 2004 and then increased more rapidly from 2005 through 2013). The number of vehicle miles traveled by light-duty motor vehicles (passenger cars and light-duty trucks) increased 35 percent from 1990 to 2013, as a result of a confluence of factors including population growth, economic growth, urban sprawl, and low fuel prices during the beginning of this period. The decline in new light-duty vehicle fuel economy between 1990 and 2004 reflected the increasing market share of light-duty trucks, which grew from about 30 percent of new vehicle sales in 1990 to 48 percent in 2004. Starting in 2005, the rate of VMT growth slowed considerably (and declined rapidly in 2008) while average new vehicle fuel economy began to increase. Average new vehicle fuel economy has improved almost every year since 2005, and the truck share has decreased to about 37 percent of new vehicles in MY 2013 (EPA 2014). Between 2012 and 2013, VMT increased by only 0.6 percent. Table 2-13 provides a detailed summary of greenhouse gas emissions from transportation-related activities with electricity-related emissions included in the totals.

From 2008 to 2009, CO₂ emissions from the transportation end-use sector declined 4.2 percent. The decrease in emissions could largely be attributed to decreased economic activity in 2009 and an associated decline in the demand for transportation. Modes such as medium- and heavy-duty trucks were significantly impacted by the decline in freight transport. From 2009 to 2013, CO₂ emissions from the transportation end-use sector stabilized even as economic activity rebounded slightly.

Almost all of the energy consumed for transportation was supplied by petroleum-based products, with more than half being related to gasoline consumption in automobiles and other highway vehicles. Other fuel uses, especially diesel fuel for freight trucks and jet fuel for aircraft, accounted for the remainder. The primary driver of transportation-related emissions was CO₂ from fossil fuel combustion, which increased by 15 percent from 1990 to 2013. This rise in CO₂ emissions, combined with an increase in HFCs from close to zero emissions in 1990 to 60.5 MMT CO₂ Eq. in 2013, led to an increase in overall emissions from transportation activities of 16 percent.

Table 2-13: Transportation-Related Greenhouse Gas Emissions (MMT CO₂ Eq.)

Gas/Vehicle	1990	2005	2009	2010	2011	2012	2013
Passenger Cars	656.7	711.2	792.9	783.6	774.3	768.0	763.3
CO ₂	629.3	660.1	748.0	742.0	736.9	735.6	735.5
CH ₄	3.2	1.4	1.2	1.2	1.2	1.1	1.1
N ₂ O	24.1	18.0	13.8	12.9	12.3	10.7	9.4
HFCs	+	31.7	29.9	27.5	23.9	20.6	17.3
Light-Duty Trucks	335.6	553.3	351.6	349.0	332.1	326.2	323.4
CO ₂	321.1	504.3	310.2	308.9	295.0	292.0	292.4
CH ₄	1.7	0.9	0.4	0.4	0.4	0.4	0.3
N ₂ O	12.8	14.8	5.8	5.5	5.1	4.4	3.9
HFCs	+	33.3	35.2	34.2	31.7	29.3	26.7
Medium- and Heavy-Duty Trucks	231.1	409.8	389.6	403.0	401.3	401.4	407.7
CO ₂	230.1	395.9	375.1	388.4	386.8	386.8	393.2
CH ₄	0.3	0.1	0.1	0.1	0.1	0.1	0.1
N ₂ O	0.7	1.1	1.2	1.2	1.1	1.1	1.1
HFCs	+	12.7	13.2	13.2	13.3	13.3	13.3
Buses	8.4	12.1	16.2	15.9	16.9	18.0	18.3
CO ₂	8.4	11.8	15.6	15.4	16.4	17.4	17.7
CH ₄	+	+	+	+	+	+	+
N ₂ O	+	+	0.1	0.1	0.1	0.1	0.1
HFCs	+	0.3	0.4	0.4	0.4	0.4	0.4
Motorcycles	1.8	1.7	4.2	3.7	3.6	4.2	4.0
CO ₂	1.7	1.6	4.1	3.6	3.6	4.1	3.9
CH ₄	+	+	+	+	+	+	+
N ₂ O	+	+	+	+	+	+	+
Commercial Aircraft^a	110.9	133.9	120.6	114.3	115.6	114.3	115.4
CO ₂	109.9	132.7	119.5	113.3	114.6	113.3	114.3
CH ₄	+	+	+	+	+	+	+
N ₂ O	1.0	1.2	1.1	1.0	1.1	1.0	1.1
Other Aircraft^b	78.3	59.6	36.8	40.4	34.2	32.1	34.7
CO ₂	77.5	59.1	36.4	40.1	33.9	31.8	34.4
CH ₄	0.1	0.1	+	+	+	+	+
N ₂ O	0.7	0.5	0.3	0.4	0.3	0.3	0.3
Ships and Boats^c	44.9	45.2	38.9	45.0	46.7	40.4	39.6
CO ₂	44.3	44.5	38.4	44.2	45.8	39.6	38.9
CH ₄	+	+	+	+	+	+	+
N ₂ O	0.6	0.6	0.5	0.8	0.8	0.7	0.7
HFCs	+	+	+	+	+	+	+
Rail	39.0	53.3	43.7	46.5	48.1	46.8	47.5
CO ₂	38.5	50.3	40.7	43.4	45.0	43.7	44.4
CH ₄	0.1	0.1	0.1	0.1	0.1	0.1	0.1
N ₂ O	0.3	0.4	0.3	0.3	0.3	0.3	0.3
HFCs	+	2.5	2.6	2.6	2.6	2.6	2.6
Other Emissions from Electricity Generation ^d	0.1	0.1	+	+	+	+	+
Pipelines^e	36.0	32.2	36.7	37.1	37.8	40.3	47.7
CO ₂	36.0	32.2	36.7	37.1	37.8	40.3	47.7
Lubricants	11.8	10.2	8.5	9.5	9.0	8.3	8.8

CO ₂	11.8	10.2	8.5	9.5	9.0	8.3	8.8
Total Transportation	1,554.4	2,022.5	1,839.9	1,848.1	1,819.7	1,799.8	1,810.3
<i>International Bunker Fuels^f</i>	<i>104.5</i>	<i>114.2</i>	<i>107.5</i>	<i>118.1</i>	<i>112.8</i>	<i>106.8</i>	<i>100.7</i>

Note: Emissions values are presented in CO₂ equivalent mass units using IPCC AR4 GWP values.

Note: Totals may not sum due to independent rounding. Passenger cars and light-duty trucks include vehicles typically used for personal travel and less than 8,500 lbs; medium- and heavy-duty trucks include vehicles larger than 8,500 lbs. HFC emissions primarily reflect HFC-134a.

+ Does not exceed 0.05 MMT CO₂ Eq.

^a Consists of emissions from jet fuel consumed by domestic operations of commercial aircraft (no bunkers).

^b Consists of emissions from jet fuel and aviation gasoline consumption by general aviation and military aircraft.

^c Fluctuations in emission estimates are associated with fluctuations in reported fuel consumption, and may reflect issues with data sources.

^d Other emissions from electricity generation are a result of waste incineration (as the majority of municipal solid waste is combusted in “trash-to-steam” electricity generation plants), electrical transmission and distribution, and a portion of Other Process Uses of Carbonates (from pollution control equipment installed in electricity generation plants).

^e CO₂ estimates reflect natural gas used to power pipelines, but not electricity. While the operation of pipelines produces CH₄ and N₂O, these emissions are not directly attributed to pipelines in the U.S. Inventory.

^f Emissions from International Bunker Fuels include emissions from both civilian and military activities; these emissions are not included in the transportation totals.

Commercial

The commercial sector is heavily reliant on electricity for meeting energy needs, with electricity consumption for lighting, heating, air conditioning, and operating appliances. The remaining emissions were largely due to the direct consumption of natural gas and petroleum products, primarily for heating and cooking needs. Energy-related emissions from the residential and commercial sectors have generally been increasing since 1990, and are often correlated with short-term fluctuations in energy consumption caused by weather conditions, rather than prevailing economic conditions. Landfills and wastewater treatment are included in this sector, with landfill emissions decreasing since 1990 and wastewater treatment emissions decreasing slightly.

Residential

The residential sector is heavily reliant on electricity for meeting energy needs, with electricity consumption for lighting, heating, air conditioning, and operating appliances. The remaining emissions were largely due to the direct consumption of natural gas and petroleum products, primarily for heating and cooking needs. Emissions from the residential sectors have generally been increasing since 1990, and are often correlated with short-term fluctuations in energy consumption caused by weather conditions, rather than prevailing economic conditions. In the long-term, this sector is also affected by population growth, regional migration trends, and changes in housing and building attributes (e.g., size and insulation).

Agriculture

The agriculture sector includes a variety of processes, including enteric fermentation in domestic livestock, livestock manure management, and agricultural soil management. In 2013, agricultural soil management was the largest source of N₂O emissions, and enteric fermentation was the largest source of CH₄ emissions in the United States. This sector also includes small amounts of CO₂ emissions from fossil fuel combustion by motorized farm equipment like tractors. The agriculture sector is less reliant on electricity than the other sectors.

Box 2-1: Methodology for Aggregating Emissions by Economic Sector

In presenting the Economic Sectors in the annual Inventory of U.S. Greenhouse Gas Emissions and Sinks, the Inventory expands upon the standard IPCC sectors common for UNFCCC reporting. Discussing greenhouse gas emissions relevant to U.S.-specific sectors improves communication of the report's findings.

In the Electricity Generation economic sector, CO₂ emissions from the combustion of fossil fuels included in the EIA electric utility fuel consuming sector are apportioned to this economic sector. Stationary combustion emissions of CH₄ and N₂O are also based on the EIA electric utility sector. Additional sources include CO₂, CH₄, and N₂O from waste incineration, as the majority of municipal solid waste is combusted in "trash-to-steam" electricity generation plants. The Electricity Generation economic sector also includes SF₆ from Electrical Transmission and Distribution, and a portion of CO₂ from Other Process Uses of Carbonates (from pollution control equipment installed in electricity generation plants).

In the Transportation economic sector, the CO₂ emissions from the combustion of fossil fuels included in the EIA transportation fuel consuming sector are apportioned to this economic sector (additional analyses and refinement of the EIA data is further explained in the Energy chapter of this report). Additional emissions are apportioned from the CH₄ and N₂O from Mobile Combustion, based on the EIA transportation sector. Substitutes of Ozone Depleting Substances are apportioned based on their specific end-uses within the source category, with emissions from transportation refrigeration/air-conditioning systems to this economic sector. Finally, CO₂ emissions from Non-Energy Uses of Fossil Fuels identified as lubricants for transportation vehicles are included in the Transportation economic sector.

For the Industry economic sector, the CO₂ emissions from the combustion of fossil fuels included in the EIA industrial fuel consuming sector, minus the agricultural use of fuel explained below, are apportioned to this economic sector. Stationary and mobile combustion emissions of CH₄ and N₂O are also based on the EIA industrial sector, minus emissions apportioned to the Agriculture economic sector described below. Substitutes of Ozone Depleting Substances are apportioned based on their specific end-uses within the source category, with most emissions falling within the Industry economic sector (minus emissions from the other economic sectors). Additionally, all process-related emissions from sources with methods considered within the IPCC Industrial Process guidance have been apportioned to this economic sector. This includes the process-related emissions (i.e., emissions from the actual process to make the material, not from fuels to power the plant) from such activities as Cement Production, Iron and Steel Production and Metallurgical Coke Production, and Ammonia Production. Additionally, fugitive emissions from energy production sources, such as Natural Gas Systems, Coal Mining, and Petroleum Systems are included in the Industry economic sector. A portion of CO₂ from Other Process Uses of Carbonates (from pollution control equipment installed in large industrial facilities) are also included in the Industry economic sector. Finally, all remaining CO₂ emissions from Non-Energy Uses of Fossil Fuels are assumed to be industrial in nature (besides the lubricants for transportation vehicles specified above), and are attributed to the Industry economic sector.

As agriculture equipment is included in EIA's industrial fuel consuming sector surveys, additional data is used to extract the fuel used by agricultural equipment, to allow for accurate reporting in the Agriculture economic sector from all sources of emissions, such as motorized farming equipment. Energy consumption estimates are obtained from Department of Agriculture survey data, in combination with separate EIA fuel sales reports. This supplementary data is used to apportion CO₂ emissions from fossil fuel combustion, and CH₄ and N₂O emissions from stationary and mobile combustion (all data is removed from the Industrial economic sector, to avoid double-counting). The other emission sources included in this economic sector are intuitive for the agriculture sectors, such as N₂O emissions from Agricultural Soils, CH₄ from Enteric Fermentation (i.e., exhalation from the digestive tracts of domesticated animals), CH₄ and N₂O from Manure Management, CH₄ from Rice Cultivation, CO₂ emissions from Liming of Agricultural Soils and Urea Application, and CH₄ and N₂O from Forest Fires. N₂O emissions from the Application of Fertilizers to tree plantations (termed "forest land" by the IPCC) are also included in the Agriculture economic sector.

The Residential economic sector includes the CO₂ emissions from the combustion of fossil fuels reported for the EIA residential sector. Stationary combustion emissions of CH₄ and N₂O are also based on the EIA residential fuel consuming sector. Substitutes of Ozone Depleting Substances are apportioned based on their specific end-uses within the source category, with emissions from residential air-conditioning systems to this economic sector. N₂O

emissions from the Application of Fertilizers to developed land (termed “settlements” by the IPCC) are also included in the Residential economic sector.

The Commercial economic sector includes the CO₂ emissions from the combustion of fossil fuels reported in the EIA commercial fuel consuming sector data. Stationary combustion emissions of CH₄ and N₂O are also based on the EIA commercial sector. Substitutes of Ozone Depleting Substances are apportioned based on their specific end-uses within the source category, with emissions from commercial refrigeration/air-conditioning systems to this economic sector. Public works sources including direct CH₄ from Landfills and CH₄ and N₂O from Wastewater Treatment and Composting are included in this economic sector.

Box 2-2: Recent Trends in Various U.S. Greenhouse Gas Emissions-Related Data

Total emissions can be compared to other economic and social indices to highlight changes over time. These comparisons include: (1) emissions per unit of aggregate energy consumption, because energy-related activities are the largest sources of emissions; (2) emissions per unit of fossil fuel consumption, because almost all energy-related emissions involve the combustion of fossil fuels; (3) emissions per unit of electricity consumption, because the electric power industry—utilities and non-utilities combined—was the largest source of U.S. greenhouse gas emissions in 2013; (4) emissions per unit of total gross domestic product as a measure of national economic activity; or (5) emissions per capita.

Table 2-14 provides data on various statistics related to U.S. greenhouse gas emissions normalized to 1990 as a baseline year. Greenhouse gas emissions in the United States have grown at an average annual rate of 0.3 percent since 1990. Since 1990, this rate is slightly slower than that for total energy and for fossil fuel consumption, and much slower than that for electricity consumption, overall gross domestic product and national population (see Table 2-14).

Table 2-14: Recent Trends in Various U.S. Data (Index 1990 = 100)

Chapter/IPCC Sector	1990	2005	2009	2010	2011	2012	2013	Growth ^a
Greenhouse Gas Emissions ^b	100	117	107	109	108	104	106	0.3%
Energy Consumption ^c	100	118	112	116	115	112	115	0.6%
Fossil Fuel Consumption ^c	100	119	108	112	110	107	110	0.4%
Electricity Consumption ^c	100	134	131	137	137	135	135	1.3%
GDP ^d	100	159	161	165	168	172	175	2.5%
Population ^e	100	118	123	124	125	125	126	1.0%

^a Average annual growth rate


^b GWP-weighted values

^c Energy-content-weighted values (EIA 2015)

^d Gross Domestic Product in chained 2009 dollars (BEA 2014)

^e U.S. Census Bureau (2014)

Figure 2-14: U.S. Greenhouse Gas Emissions Per Capita and Per Dollar of Gross Domestic Product



Source: BEA (2014), U.S. Census Bureau (2014), and emission estimates in this report.

2.2 Indirect Greenhouse Gas Emissions (CO, NO_x, NMVOCs, and SO₂)

The reporting requirements of the UNFCCC⁷ request that information be provided on indirect greenhouse gases, which include CO, NO_x, NMVOCs, and SO₂. These gases do not have a direct global warming effect, but indirectly affect terrestrial radiation absorption by influencing the formation and destruction of tropospheric and stratospheric ozone, or, in the case of SO₂, by affecting the absorptive characteristics of the atmosphere. Additionally, some of these gases may react with other chemical compounds in the atmosphere to form compounds that are greenhouse gases. Carbon monoxide is produced when carbon-containing fuels are combusted incompletely. Nitrogen oxides (i.e., NO and NO₂) are created by lightning, fires, fossil fuel combustion, and in the stratosphere from N₂O. Non-CH₄ volatile organic compounds—which include hundreds of organic compounds that participate in atmospheric chemical reactions (i.e., propane, butane, xylene, toluene, ethane, and many others)—are emitted primarily from transportation, industrial processes, and non-industrial consumption of organic solvents. In the United States, SO₂ is primarily emitted from coal combustion for electric power generation and the metals industry. Sulfur-containing compounds emitted into the atmosphere tend to exert a negative radiative forcing (i.e., cooling) and therefore are discussed separately.

One important indirect climate change effect of NMVOCs and NO_x is their role as precursors for tropospheric ozone formation. They can also alter the atmospheric lifetimes of other greenhouse gases. Another example of indirect greenhouse gas formation into greenhouse gases is CO's interaction with the hydroxyl radical—the major atmospheric sink for CH₄ emissions—to form CO₂. Therefore, increased atmospheric concentrations of CO limit the number of hydroxyl molecules (OH) available to destroy CH₄.

⁷ See < <http://unfccc.int/resource/docs/2013/cop19/eng/10a03.pdf#page=2>>.

Since 1970, the United States has published estimates of emissions of CO, NO_x, NMVOCs, and SO₂ (EPA 2015),⁸ which are regulated under the Clean Air Act. Table 2-15 shows that fuel combustion accounts for the majority of emissions of these indirect greenhouse gases. Industrial processes—such as the manufacture of chemical and allied products, metals processing, and industrial uses of solvents—are also significant sources of CO, NO_x, and NMVOCs.

Table 2-15: Emissions of NO_x, CO, NMVOCs, and SO₂ (kt)

Gas/Activity	1990	2005	2009	2010	2011	2012	2013
NO_x	21,771	17,394	13,450	12,607	12,630	11,912	11,167
Mobile Fossil Fuel Combustion	10,862	10,295	7,797	7,290	7,294	6,788	6,283
Stationary Fossil Fuel Combustion	10,023	5,858	4,452	4,092	3,807	3,567	3,579
Oil and Gas Activities	139	321	468	545	622	622	622
Industrial Processes and Product Use	592	572	493	472	452	452	452
Forest Fires	64	212	149	121	373	400	149
Waste Combustion	82	128	81	77	73	73	73
Agricultural Burning	8	6	8	8	8	8	8
Waste	+	2	1	1	1	1	1
CO	132,337	74,283	51,716	50,996	58,868	58,022	47,265
Mobile Fossil Fuel Combustion	119,360	58,615	39,256	39,475	38,305	36,491	34,676
Forest Fires	2,300	7,550	5,313	4,323	13,291	14,262	5,310
Stationary Fossil Fuel Combustion	5,000	4,648	4,036	4,103	4,170	4,170	4,170
Industrial Processes and Product Use	4,129	1,557	1,331	1,280	1,229	1,229	1,229
Waste Combustion	978	1,403	1,164	1,084	1,003	1,003	1,003
Oil and Gas Activities	302	318	363	487	610	610	610
Agricultural Burning	268	184	247	241	255	253	262
Waste	1	7	5	5	5	5	5
NMVOCs	20,930	13,154	11,586	11,641	11,726	11,416	11,107
Mobile Fossil Fuel Combustion	10,932	5,724	4,650	4,591	4,562	4,252	3,942
Industrial Processes and Product Use	7,638	5,849	4,337	4,133	3,929	3,929	3,929
Oil and Gas Activities	554	510	1,894	2,205	2,517	2,517	2,517
Stationary Fossil Fuel Combustion	912	716	553	576	599	599	599
Waste Combustion	222	241	103	92	81	81	81
Waste	673	114	49	44	38	38	38
Agricultural Burning	NA	NA	NA	NA	NA	NA	NA
SO₂	20,935	13,196	8,246	7,015	5,877	4,711	4,625
Stationary Fossil Fuel Combustion	18,407	11,541	7,228	6,120	5,008	3,859	3,790
Industrial Processes and Product Use	1,307	831	654	618	605	605	605
Oil and Gas Activities	390	180	126	117	108	108	108
Mobile Fossil Fuel Combustion	793	619	220	144	142	125	108
Waste Combustion	38	25	17	16	15	15	15
Waste	+	1	1	+	+	+	+
Agricultural Burning	NA	NA	NA	NA	NA	NA	NA

Source: (EPA 2015) except for estimates from Field Burning of Agricultural Residues.

NA (Not Available)

Note: Totals may not sum due to independent rounding.

+ Does not exceed 0.5 kt.

Box 2-3: Sources and Effects of Sulfur Dioxide

Sulfur dioxide (SO₂) emitted into the atmosphere through natural and anthropogenic processes affects the earth's radiative budget through its photochemical transformation into sulfate aerosols that can (1) scatter radiation from the sun back to space, thereby reducing the radiation reaching the earth's surface; (2) affect cloud formation; and (3)

⁸ NO_x and CO emission estimates from Field Burning of Agricultural Residues were estimated separately, and therefore not taken from EPA (2015).

affect atmospheric chemical composition (e.g., by providing surfaces for heterogeneous chemical reactions). The indirect effect of sulfur-derived aerosols on radiative forcing can be considered in two parts. The first indirect effect is the aerosols' tendency to decrease water droplet size and increase water droplet concentration in the atmosphere. The second indirect effect is the tendency of the reduction in cloud droplet size to affect precipitation by increasing cloud lifetime and thickness. Although still highly uncertain, the radiative forcing estimates from both the first and the second indirect effect are believed to be negative, as is the combined radiative forcing of the two (IPCC 2001). However, because SO₂ is short-lived and unevenly distributed in the atmosphere, its radiative forcing impacts are highly uncertain.

Sulfur dioxide is also a major contributor to the formation of regional haze, which can cause significant increases in acute and chronic respiratory diseases. Once SO₂ is emitted, it is chemically transformed in the atmosphere and returns to the earth as the primary source of acid rain. Because of these harmful effects, the United States has regulated SO₂ emissions in the Clean Air Act.

Electricity generation is the largest anthropogenic source of SO₂ emissions in the United States, accounting for 64.4 percent in 2013. Coal combustion contributes nearly all of those emissions (approximately 92 percent). Sulfur dioxide emissions have decreased in recent years, primarily as a result of electric power generators switching from high-sulfur to low-sulfur coal and installing flue gas desulfurization equipment.
