


## 5. Agriculture

Agricultural activities contribute directly to emissions of greenhouse gases through a variety of processes. This chapter provides an assessment of methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) emissions from enteric fermentation in domestic livestock, livestock manure management, rice cultivation, agricultural soil management, and field burning of agricultural residues; as well as carbon dioxide (CO<sub>2</sub>) emissions from liming and urea fertilization (see Figure 5-1). Additional CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O fluxes from agriculture-related land-use and land-use conversion activities, such as cultivation of cropland, grassland fires and conversion of forest land to cropland, are presented in the Land Use, Land-Use Change, and Forestry (LULUCF) chapter. Carbon dioxide emissions from on-farm energy use are reported in the Energy chapter.

**Figure 5-1: 2017 Agriculture Chapter Greenhouse Gas Emission Sources (MMT CO<sub>2</sub> Eq.)**



In 2017, the Agriculture sector was responsible for emissions of 542.1 MMT CO<sub>2</sub> Eq.,<sup>1</sup> or 8.4 percent of total U.S. greenhouse gas emissions.<sup>2</sup> Methane emissions from enteric fermentation and manure management represent 26.4 percent and 9.3 percent of total CH<sub>4</sub> emissions from anthropogenic activities, respectively. Of all domestic animal types, beef and dairy cattle were by far the largest emitters of CH<sub>4</sub>. Rice cultivation and field burning of agricultural residues were minor sources of CH<sub>4</sub>. Emissions of N<sub>2</sub>O by agricultural soil management through activities such as fertilizer application and other agricultural practices that increased nitrogen availability in the soil was the largest source of U.S. N<sub>2</sub>O emissions, accounting for 73.9 percent. Manure management and field burning of agricultural residues were also small sources of N<sub>2</sub>O emissions. Urea fertilization and liming each accounted for 0.1 percent of total CO<sub>2</sub> emissions from anthropogenic activities.

<sup>1</sup> Following the current reporting requirements under the United Nations Framework Convention on Climate Change (UNFCCC), this Inventory report presents CO<sub>2</sub> equivalent values based on the *IPCC Fourth Assessment Report (AR4)* GWP values. See the Introduction chapter for more information.

<sup>2</sup> Emissions reported in the Agriculture chapter include those from all states, including Hawaii and Alaska; however, U.S. Territories are not included.

Table 5-1 and Table 5-2 present emission estimates for the Agriculture sector. Between 1990 and 2017, CO<sub>2</sub> and CH<sub>4</sub> emissions from agricultural activities increased by 16.2 percent and 14.4 percent, respectively, while N<sub>2</sub>O emissions from agricultural activities fluctuated from year to year, but increased by 7.3 percent overall.

**Table 5-1: Emissions from Agriculture (MMT CO<sub>2</sub> Eq.)**

Gas/Source	1990	2005	2013	2014	2015	2016	2017
<b>CO<sub>2</sub></b>	<b>7.1</b>	<b>7.9</b>	<b>8.4</b>	<b>8.1</b>	<b>8.5</b>	<b>8.1</b>	<b>8.2</b>
Urea Fertilization	2.4	3.5	4.4	4.5	4.7	4.9	5.1
Liming	4.7	4.3	3.9	3.6	3.7	3.2	3.2
<b>CH<sub>4</sub></b>	<b>217.4</b>	<b>239.5</b>	<b>235.3</b>	<b>234.9</b>	<b>239.9</b>	<b>247.3</b>	<b>248.7</b>
Enteric Fermentation	164.2	168.9	165.5	164.2	166.5	171.9	175.4
Manure Management	37.1	53.7	58.1	57.8	60.9	61.5	61.7
Rice Cultivation	16.0	16.7	11.5	12.7	12.3	13.7	11.3
Field Burning of Agricultural Residues	0.1	0.2	0.2	0.2	0.2	0.2	0.2
<b>N<sub>2</sub>O</b>	<b>265.7</b>	<b>271.1</b>	<b>282.7</b>	<b>279.7</b>	<b>295.4</b>	<b>285.8</b>	<b>285.2</b>
Agricultural Soil Management	251.7	254.5	265.2	262.3	277.8	267.6	266.4
Manure Management	14.0	16.5	17.4	17.4	17.6	18.2	18.7
Field Burning of Agricultural Residues	+	0.1	0.1	0.1	0.1	0.1	0.1
<b>Total</b>	<b>490.2</b>	<b>518.4</b>	<b>526.3</b>	<b>522.8</b>	<b>543.8</b>	<b>541.2</b>	<b>542.1</b>

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

Note: Totals may not sum due to independent rounding.

**Table 5-2: Emissions from Agriculture (kt)**

Gas/Source	1990	2005	2013	2014	2015	2016	2017
<b>CO<sub>2</sub></b>	<b>7,084</b>	<b>7,854</b>	<b>8,350</b>	<b>8,124</b>	<b>8,464</b>	<b>8,083</b>	<b>8,234</b>
Urea Fertilization	2,417	3,504	4,443	4,515	4,728	4,877	5,051
Liming	4,667	4,349	3,907	3,609	3,737	3,206	3,182
<b>CH<sub>4</sub></b>	<b>8,697</b>	<b>9,579</b>	<b>9,412</b>	<b>9,397</b>	<b>9,597</b>	<b>9,892</b>	<b>9,946</b>
Enteric Fermentation	6,566	6,755	6,620	6,568	6,661	6,875	7,018
Manure Management	1,486	2,150	2,322	2,311	2,435	2,461	2,467
Rice Cultivation	641	667	462	510	493	549	454
Field Burning of Agricultural Residues	4	7	8	8	8	8	8
<b>N<sub>2</sub>O</b>	<b>892</b>	<b>910</b>	<b>949</b>	<b>939</b>	<b>991</b>	<b>959</b>	<b>957</b>
Agricultural Soil Management	845	854	890	880	932	898	894
Manure Management	47	55	58	58	59	61	63
Field Burning of Agricultural Residues	+	+	+	+	+	+	+

+ Does not exceed 0.5 kt.

Note: Totals may not sum due to independent rounding.

#### Box 5-1: Methodological Approach for Estimating and Reporting U.S. Emissions and Removals

In following the United Nations Framework Convention on Climate Change (UNFCCC) requirement under Article 4.1 to develop and submit national greenhouse gas emission inventories, the emissions and removals presented in this report and this chapter, are organized by source and sink categories and calculated using internationally-accepted methods provided by the Intergovernmental Panel on Climate Change (IPCC) in the *2006 IPCC Guidelines for National Greenhouse Gas Inventories (2006 IPCC Guidelines)*. Additionally, the calculated emissions and removals in a given year for the United States are presented in a common manner in line with the UNFCCC reporting guidelines for the reporting of inventories under this international agreement. The use of consistent methods to calculate emissions and removals by all nations providing their inventories to the UNFCCC ensures that these reports are comparable. The presentation of emissions and removals provided in this Inventory do not preclude alternative examinations, but rather, this Inventory presents emissions and removals in a common format consistent with how countries are to report Inventories under the UNFCCC. The report itself, and this chapter, follows this

standardized format, and provides an explanation of the application of methods used to calculate emissions and removals.

## 5.1 Enteric Fermentation (CRF Source Category 3A)

Methane is produced as part of normal digestive processes in animals. During digestion, microbes resident in an animal's digestive system ferment food consumed by the animal. This microbial fermentation process, referred to as enteric fermentation, produces CH<sub>4</sub> as a byproduct, which can be exhaled or eructated by the animal. The amount of CH<sub>4</sub> produced and emitted by an individual animal depends primarily upon the animal's digestive system, and the amount and type of feed it consumes.

Ruminant animals (e.g., cattle, buffalo, sheep, goats, and camels) are the major emitters of CH<sub>4</sub> because of their unique digestive system. Ruminants possess a rumen, or large "fore-stomach," in which microbial fermentation breaks down the feed they consume into products that can be absorbed and metabolized. The microbial fermentation that occurs in the rumen enables them to digest coarse plant material that non-ruminant animals cannot. Ruminant animals, consequently, have the highest CH<sub>4</sub> emissions per unit of body mass among all animal types.

Non-ruminant animals (e.g., swine, horses, and mules and asses) also produce CH<sub>4</sub> emissions through enteric fermentation, although this microbial fermentation occurs in the large intestine. These non-ruminants emit significantly less CH<sub>4</sub> on a per-animal-mass basis than ruminants because the capacity of the large intestine to produce CH<sub>4</sub> is lower.

In addition to the type of digestive system, an animal's feed quality and feed intake also affect CH<sub>4</sub> emissions. In general, lower feed quality and/or higher feed intake leads to higher CH<sub>4</sub> emissions. Feed intake is positively correlated to animal size, growth rate, level of activity and production (e.g., milk production, wool growth, pregnancy, or work). Therefore, feed intake varies among animal types as well as among different management practices for individual animal types (e.g., animals in feedlots or grazing on pasture).

Methane emission estimates from enteric fermentation are provided in Table 5-3 and Table 5-4. Total livestock CH<sub>4</sub> emissions in 2017 were 175.4 MMT CO<sub>2</sub> Eq. (7,018 kt). Beef cattle remain the largest contributor of CH<sub>4</sub> emissions from enteric fermentation, accounting for 72 percent in 2017. Emissions from dairy cattle in 2017 accounted for 25 percent, and the remaining emissions were from horses, sheep, swine, goats, American bison, mules and asses.

**Table 5-3: CH<sub>4</sub> Emissions from Enteric Fermentation (MMT CO<sub>2</sub> Eq.)**

Livestock Type	1990	2005	2013	2014	2015	2016	2017
Beef Cattle	119.1	125.2	118.0	116.5	118.0	123.0	126.3
Dairy Cattle	39.4	37.6	41.6	42.0	42.6	43.0	43.3
Swine	2.0	2.3	2.5	2.4	2.6	2.6	2.7
Horses	1.0	1.7	1.6	1.6	1.5	1.5	1.4
Sheep	2.3	1.2	1.1	1.0	1.1	1.1	1.1
Goats	0.3	0.4	0.3	0.3	0.3	0.3	0.3
American Bison	0.1	0.4	0.3	0.3	0.3	0.3	0.3
Mules and Asses	0.0	0.1	0.1	0.1	0.1	0.1	0.1
<b>Total</b>	<b>164.2</b>	<b>168.9</b>	<b>165.5</b>	<b>164.2</b>	<b>166.5</b>	<b>171.9</b>	<b>175.4</b>

Note: Totals may not sum due to independent rounding.

**Table 5-4: CH<sub>4</sub> Emissions from Enteric Fermentation (kt)**

Livestock Type	1990	2005	2013	2014	2015	2016	2017
Beef Cattle	4,763	5,007	4,722	4,660	4,722	4,919	5,052

Dairy Cattle	1,574	1,503	1,664	1,679	1,706	1,722	1,730
Swine	81	92	98	96	102	105	108
Horses	40	70	64	62	61	59	58
Sheep	91	49	43	42	42	42	42
Goats	13	14	13	12	12	11	11
American Bison	4	17	13	13	13	13	13
Mules and Asses	1	2	3	3	3	3	3
<b>Total</b>	<b>6,566</b>	<b>6,755</b>	<b>6,620</b>	<b>6,568</b>	<b>6,661</b>	<b>6,875</b>	<b>7,018</b>

Note: Totals may not sum due to independent rounding.

From 1990 to 2017, emissions from enteric fermentation have increased by 6.9 percent. Emissions have also increased from 2016 to 2017 by 2.1 percent, largely driven by an increase in beef cattle populations. While emissions generally follow trends in cattle populations, over the long term there are exceptions. For example, beef cattle emissions increased 6.1 percent from 1990 to 2017, while the national total beef cattle population slightly decreased (by 0.02 percent) from 1990 to 2017. Furthermore, while dairy cattle emissions increased 9.9 percent over the entire time series, the population has declined by 3.2 percent, and milk production increased 44 percent (USDA 2018). This trend indicates that while emission factors per head are increasing, emission factors per unit of product are going down.

Generally, from 1990 to 1995 emissions from beef cattle increased and then decreased from 1996 to 2004. These trends were mainly due to fluctuations in beef cattle populations and increased digestibility of feed for feedlot cattle. Beef cattle emissions generally increased from 2004 to 2007, as beef cattle populations underwent increases and an extensive literature review indicated a trend toward a decrease in feed digestibility for those years. Beef cattle emissions decreased again from 2007 to 2014, as populations again decreased, but increased from 2015 to 2017, consistent with another increase in population over those same years. Emissions from dairy cattle generally trended downward from 1990 to 2004, along with an overall dairy cattle population decline during the same period. Similar to beef cattle, dairy cattle emissions rose from 2004 to 2007 due to population increases and a decrease in feed digestibility (based on an analysis of more than 350 dairy cow diets). Dairy cattle emissions have continued to trend upward since 2007, in line with dairy cattle population increases. Regarding trends in other animals, populations of sheep have steadily declined, with an overall decrease of 54 percent since 1990. Horse populations are 45 percent greater than they were in 1990, but their numbers have been declining by about 2 percent annually since 2007. Goat populations increased by about 20 percent through 2007, then steadily decreased through 2017. Swine populations have trended upward through most of the time series, increasing 34 percent from 1990 to 2017. The population of American bison more than tripled over the 1990 to 2017 time period, while the population of mules and asses increased by nearly 5 times.

## Methodology

Livestock enteric fermentation emission estimate methodologies fall into two categories: cattle and other domesticated animals. Cattle, due to their large population, large size, and particular digestive characteristics, account for the majority of enteric fermentation CH<sub>4</sub> emissions from livestock in the United States. A more detailed methodology (i.e., IPCC Tier 2) was therefore applied to estimate emissions for all cattle. Emission estimates for other domesticated animals (horses, sheep, swine, goats, American bison, and mules and asses) were handled using a less detailed approach (i.e., IPCC Tier 1).

While the large diversity of animal management practices cannot be precisely characterized and evaluated, significant scientific literature exists that provides the necessary data to estimate cattle emissions using the IPCC Tier 2 approach. The Cattle Enteric Fermentation Model (CEFM), developed by EPA and used to estimate cattle CH<sub>4</sub> emissions from enteric fermentation, incorporates this information and other analyses of livestock population, feeding practices, and production characteristics.

### *Inventory Methodology for Cattle*

National cattle population statistics were disaggregated into the following cattle sub-populations:

- Dairy Cattle

- Calves
- Heifer Replacements
- Cows
- Beef Cattle
  - Calves
  - Heifer Replacements
  - Heifer and Steer Stockers
  - Animals in Feedlots (Heifers and Steer)
  - Cows
  - Bulls

Calf birth rates, end-of-year population statistics, detailed feedlot placement information, and slaughter weight data were used to create a transition matrix that models cohorts of individual animal types and their specific emission profiles. The key variables tracked for each of the cattle population categories are described in Annex 3.10. These variables include performance factors such as pregnancy and lactation as well as average weights and weight gain. Annual cattle population data were obtained from the U.S. Department of Agriculture's (USDA) National Agricultural Statistics Service (NASS) *QuickStats* database (USDA 2016).

Diet characteristics were estimated by region for dairy, grazing beef, and feedlot beef cattle. These diet characteristics were used to calculate digestible energy (DE) values (expressed as the percent of gross energy intake digested by the animal) and CH<sub>4</sub> conversion rates (Y<sub>m</sub>) (expressed as the fraction of gross energy converted to CH<sub>4</sub>) for each regional population category. The IPCC recommends Y<sub>m</sub> ranges of 3.0±1.0 percent for feedlot cattle and 6.5±1.0 percent for other well-fed cattle consuming temperate-climate feed types (IPCC 2006). Given the availability of detailed diet information for different regions and animal types in the United States, DE and Y<sub>m</sub> values unique to the United States were developed. The diet characterizations and estimation of DE and Y<sub>m</sub> values were based on information from state agricultural extension specialists, a review of published forage quality studies and scientific literature, expert opinion, and modeling of animal physiology.

The diet characteristics for dairy cattle were based on Donovan (1999) and an extensive review of nearly 20 years of literature from 1990 through 2009. Estimates of DE were national averages based on the feed components of the diets observed in the literature for the following year groupings: 1990 through 1993, 1994 through 1998, 1999 through 2003, 2004 through 2006, 2007, and 2008 onward.<sup>3</sup> Base year Y<sub>m</sub> values by region were estimated using Donovan (1999). As described in ERG (2016), a ruminant digestion model (COWPOLL, as selected in Kebreab et al. 2008) was used to evaluate Y<sub>m</sub> for each diet evaluated from the literature, and a function was developed to adjust regional values over time based on the national trend. Dairy replacement heifer diet assumptions were based on the observed relationship in the literature between dairy cow and dairy heifer diet characteristics.

For feedlot animals, the DE and Y<sub>m</sub> values used for 1990 were recommended by Johnson (1999). Values for DE and Y<sub>m</sub> for 1991 through 1999 were linearly extrapolated based on the 1990 and 2000 data. DE and Y<sub>m</sub> values for 2000 onwards were based on survey data in Galyean and Gleghorn (2001) and Vasconcelos and Galyean (2007).

For grazing beef cattle, Y<sub>m</sub> values were based on Johnson (2002), DE values for 1990 through 2006 were based on specific diet components estimated from Donovan (1999), and DE values from 2007 onwards were developed from an analysis by Archibeque (2011), based on diet information in Preston (2010) and USDA-APHIS:VS (2010). Weight and weight gains for cattle were estimated from Holstein (2010), Doren et al. (1989), Enns (2008), Lippke et al. (2000), Pinchack et al. (2004), Platter et al. (2003), Skogerboe et al. (2000), and expert opinion. See Annex 3.10 for more details on the method used to characterize cattle diets and weights in the United States.

Calves younger than 4 months are not included in emission estimates because calves consume mainly milk and the IPCC recommends the use of a Y<sub>m</sub> of zero for all juveniles consuming only milk. Diets for calves aged 4 to 6 months are assumed to go through a gradual weaning from milk decreasing to 75 percent at 4 months, 50 percent at age 5 months, and 25 percent at age 6 months. The portion of the diet made up with milk still results in zero emissions. For the remainder of the diet, beef calf DE and Y<sub>m</sub> are set equivalent to those of beef replacement heifers, while dairy calf DE is set equal to that of dairy replacement heifers and dairy calf Y<sub>m</sub> is provided at 4 and 7 months

<sup>3</sup> Due to inconsistencies in the 2003 literature values, the 2002 values were used for 2003, as well.

of age by Soliva (2006). Estimates of  $Y_m$  for 5 and 6 month old dairy calves are linearly interpolated from the values provided for 4 and 7 months.

To estimate  $CH_4$  emissions, the population was divided into state, age, sub-type (i.e., dairy cows and replacements, beef cows and replacements, heifer and steer stockers, heifers and steers in feedlots, bulls, beef calves 4 to 6 months, and dairy calves 4 to 6 months), and production (i.e., pregnant, lactating) groupings to more fully capture differences in  $CH_4$  emissions from these animal types. The transition matrix was used to simulate the age and weight structure of each sub-type on a monthly basis in order to more accurately reflect the fluctuations that occur throughout the year. Cattle diet characteristics were then used in conjunction with Tier 2 equations from IPCC (2006) to produce  $CH_4$  emission factors for the following cattle types: dairy cows, beef cows, dairy replacements, beef replacements, steer stockers, heifer stockers, steer feedlot animals, heifer feedlot animals, bulls, and calves. To estimate emissions from cattle, monthly population data from the transition matrix were multiplied by the calculated emission factor for each cattle type. More details are provided in Annex 3.10.

### *Inventory Methodology for Non-Cattle Livestock*

Emission estimates for other animal types were based on average emission factors representative of entire populations of each animal type. Methane emissions from these animals accounted for a minor portion of total  $CH_4$  emissions from livestock in the United States from 1990 through 2017. Additionally, the variability in emission factors for each of these other animal types (e.g., variability by age, production system, and feeding practice within each animal type) is less than that for cattle.

Annual livestock population data for 1990 to 2017 for sheep; swine; goats; horses; mules and asses; and American bison were obtained for available years from USDA-NASS (USDA 2016). Horse, goat and mule and ass population data were available for 1987, 1992, 1997, 2002, 2007, and 2012 (USDA 1992, 1997, 2016); the remaining years between 1990 and 2017 were interpolated and extrapolated from the available estimates (with the exception of goat populations being held constant between 1990 and 1992). American bison population estimates were available from USDA for 2002, 2007, and 2012 (USDA 2016) and from the National Bison Association (1999) for 1990 through 1999. Additional years were based on observed trends from the National Bison Association (1999), interpolation between known data points, and extrapolation beyond 2012, as described in more detail in Annex 3.10.

Methane emissions from sheep, goats, swine, horses, American bison, and mules and asses were estimated by using emission factors utilized in Crutzen et al. (1986, cited in IPCC 2006). These emission factors are representative of typical animal sizes, feed intakes, and feed characteristics in developed countries. For American bison the emission factor for buffalo was used and adjusted based on the ratio of live weights to the 0.75 power. The methodology is the same as that recommended by IPCC (2006).

See Annex 3.10 for more detailed information on the methodology and data used to calculate  $CH_4$  emissions from enteric fermentation.

## **Uncertainty and Time-Series Consistency**

A quantitative uncertainty analysis for this source category was performed using the IPCC-recommended Approach 2 uncertainty estimation methodology based on a Monte Carlo Stochastic Simulation technique as described in ICF (2003). These uncertainty estimates were developed for the 1990 through 2001 Inventory (i.e., 2003 submission to the UNFCCC). There have been no significant changes to the methodology since that time; consequently, these uncertainty estimates were directly applied to the 2017 emission estimates in this Inventory.

A total of 185 primary input variables (177 for cattle and 8 for non-cattle) were identified as key input variables for the uncertainty analysis. A normal distribution was assumed for almost all activity- and emission factor-related input variables. Triangular distributions were assigned to three input variables (specifically, cow-birth ratios for the three most recent years included in the 2001 model run) to ensure only positive values would be simulated. For some key input variables, the uncertainty ranges around their estimates (used for inventory estimation) were collected from published documents and other public sources; others were based on expert opinion and best estimates. In addition, both endogenous and exogenous correlations between selected primary input variables were modeled. The exogenous correlation coefficients between the probability distributions of selected activity-related variables were developed through expert judgment.

The uncertainty ranges associated with the activity data-related input variables were plus or minus 10 percent or lower. However, for many emission factor-related input variables, the lower- and/or the upper-bound uncertainty estimates were over 20 percent. The results of the quantitative uncertainty analysis are summarized in Table 5-5. Based on this analysis, enteric fermentation CH<sub>4</sub> emissions in 2017 were estimated to be between 156.1 and 207.0 MMT CO<sub>2</sub> Eq. at a 95 percent confidence level, which indicates a range of 11 percent below to 18 percent above the 2017 emission estimate of 175.4 MMT CO<sub>2</sub> Eq. Among the individual cattle sub-source categories, beef cattle account for the largest amount of CH<sub>4</sub> emissions, as well as the largest degree of uncertainty in the emission estimates—due mainly to the difficulty in estimating the diet characteristics for grazing members of this animal group. Among non-cattle, horses represent the largest percent of uncertainty in the previous uncertainty analysis because the Food and Agricultural Organization of the United Nations (FAO) population estimates used for horses at that time had a higher degree of uncertainty than for the USDA population estimates used for swine, goats, and sheep. The horse populations are now from the same USDA source as the other animal types, and therefore the uncertainty range around horses is likely overestimated. Cattle calves, American bison, mules and asses were excluded from the initial uncertainty estimate because they were not included in emission estimates at that time.

**Table 5-5: Approach 2 Quantitative Uncertainty Estimates for CH<sub>4</sub> Emissions from Enteric Fermentation (MMT CO<sub>2</sub> Eq. and Percent)**

Source	Gas	2017 Emission Estimate (MMT CO <sub>2</sub> Eq.)	Uncertainty Range Relative to Emission Estimate <sup>a, b, c</sup>			
			(MMT CO <sub>2</sub> Eq.)		(%)	
			Lower Bound	Upper Bound	Lower Bound	Upper Bound
Enteric Fermentation	CH <sub>4</sub>	175.4	156.1	207.0	-11%	+18%

<sup>a</sup> Range of emissions estimates predicted by Monte Carlo Stochastic Simulation for a 95 percent confidence interval.

<sup>b</sup> Note that the relative uncertainty range was estimated with respect to the 2001 emission estimates from the 2003 submission and applied to the 2017 estimates.

<sup>c</sup> The overall uncertainty calculated in 2003, and applied to the 2017 emission estimate, did not include uncertainty estimates for calves, American bison, and mules and asses. Additionally, for bulls the emissions estimate was based on the Tier 1 methodology. Since bull emissions are now estimated using the Tier 2 method, the uncertainty surrounding their estimates is likely lower than indicated by the previous uncertainty analysis.

Details on the emission trends through time are described in more detail in the Methodology section.

## QA/QC and Verification

In order to ensure the quality of the emission estimates from enteric fermentation, the General (IPCC Tier 1) and category-specific (Tier 2) Quality Assurance/Quality Control (QA/QC) procedures were implemented consistent with the U.S. QA/QC plan (EPA 2002). Category-specific or Tier 2 QA procedures included independent review of emission estimate methodologies from previous inventories. Over the past few years, particular importance has been placed on harmonizing the data exchange between the enteric fermentation and manure management source categories. The current Inventory now utilizes the transition matrix from the CEFM for estimating cattle populations and weights for both source categories, and the CEFM is used to output volatile solids and nitrogen excretion estimates using the diet assumptions in the model in conjunction with the energy balance equations from the IPCC (2006). This approach facilitates the QA/QC process for both of these source categories.

## Recalculations Discussion

In the previous Inventory, 1990 to 2015 estimates were retained from the prior Inventory (i.e., 1990 through 2015 Inventory), and 2016 estimates were based on a simplified approach that used emission factors and extrapolated population estimates for all animals. For the current Inventory, the CEFM was used for cattle for all years, resulting in different estimates for 2016 than the prior Inventory. For non-cattle livestock in the current Inventory, updated Tier 1 estimates were used for 2016, yielding different results than the simplified approach used for 2016 in the prior Inventory.



1 There were also changes to emissions resulting from activity data changes, including:

- 2 • The USDA published minor revisions in several categories that impacted emissions estimated for cattle for  
3 2015, including the following:
  - 4 ○ Calf birth data were revised;
  - 5 ○ Dairy cow milk production values were revised for several states;
  - 6 ○ Slaughter values were revised for steers and heifers.
- 7 • The USDA also revised population estimates for some categories of non-cattle animals, which impacted  
8 emissions estimated for “other” livestock. Populations for market and breeding swine were changed for some  
9 states for 2015.
- 10 • American Bison populations from the 2012 Census were carried over for 2013 through 2017 values instead of  
11 using predictive estimates of the populations. This change yielded different emissions estimates for 2013  
12 through 2016 for American Bison as compared to the previous Inventory.

13 These recalculations had an insignificant impact on the overall emission estimates.

## 14 **Planned Improvements**

15 Continued research and regular updates are necessary to maintain an emissions inventory that reflects the current  
16 base of knowledge. Depending upon the outcome of ongoing investigations, future improvements for enteric  
17 fermentation could include some of the following options:

- 18 • Further research to improve the estimation of dry matter intake (as gross energy intake) using data from  
19 appropriate production systems;
- 20 • Updating input variables that are from older data sources, such as beef births by month and beef cow  
21 lactation rates;
- 22 • Investigation of the availability of annual data for the DE,  $Y_m$ , and crude protein values of specific diet and  
23 feed components for grazing and feedlot animals;
- 24 • Further investigation on additional sources or methodologies for estimating DE for dairy cattle, given the  
25 many challenges in characterizing dairy cattle diets;
- 26 • Further evaluation of the assumptions about weights and weight gains for beef cows, such that trends  
27 beyond 2007 are updated, rather than held constant;
- 28 • Further evaluation of the estimated weight for dairy cows (i.e., 1,500 lbs) that is based solely on Holstein  
29 cows as mature dairy cow weight is likely slightly overestimated, based on knowledge of the breeds of  
30 dairy cows in the United States;
- 31 • Potentially updating to a Tier 2 methodology for other animal types (i.e., sheep, swine, goats, horses);
- 32 • Investigation of methodologies and emission factors for including enteric fermentation emission estimates  
33 from poultry;
- 34 • Comparison of the current CEFM processing of animal population data to estimates developed using  
35 annual average populations to determine if the model could be simplified to use annual population data;
- 36 • Investigation of recent research implications suggesting that certain parameters in enteric models may be  
37 simplified without significantly diminishing model accuracy; and
- 38 • Recent changes that have been implemented to the CEFM warrant an assessment of the current uncertainty  
39 analysis; therefore, a revision of the quantitative uncertainty surrounding emission estimates from this  
40 source category will be initiated.

41 In future Inventory reports, the final 2019 Refinement to the IPCC Guidelines [currently in draft] will be reviewed  
42 and any changes will be incorporated, as applicable, to update the current Inventory estimation methodologies.



## 5.2 Manure Management (CRF Source Category 3B)

---

The treatment, storage, and transportation of livestock manure can produce anthropogenic CH<sub>4</sub> and N<sub>2</sub>O emissions. Methane is produced by the anaerobic decomposition of manure and nitrous oxide is produced from direct and indirect pathways through the processes of nitrification and denitrification; in addition, there are many underlying factors that can affect these resulting emissions from manure management, as described below.

When livestock or poultry manure are stored or treated in systems that promote anaerobic conditions (e.g., as a liquid/slurry in lagoons, ponds, tanks, or pits), the decomposition of the volatile solids component in the manure tends to produce CH<sub>4</sub>. When manure is handled as a solid (e.g., in stacks or drylots) or deposited on pasture, range, or paddock lands, it tends to decompose aerobically and produce CO<sub>2</sub> and little or no CH<sub>4</sub>. Ambient temperature, moisture, and manure storage or residency time affect the amount of CH<sub>4</sub> produced because they influence the growth of the bacteria responsible for CH<sub>4</sub> formation. For non-liquid-based manure systems, moist conditions (which are a function of rainfall and humidity) can promote CH<sub>4</sub> production. Manure composition, which varies by animal diet, growth rate, and animal type (particularly the different animal digestive systems), also affects the amount of CH<sub>4</sub> produced. In general, the greater the energy content of the feed, the greater the potential for CH<sub>4</sub> emissions. However, some higher-energy feeds also are more digestible than lower quality forages, which can result in less overall waste excreted from the animal.

As previously stated, nitrous oxide emissions are produced through both direct and indirect pathways. Direct N<sub>2</sub>O emissions are produced as part of the nitrogen (N) cycle through the nitrification and denitrification of the N in livestock dung and urine.<sup>4</sup> There are two pathways for indirect N<sub>2</sub>O emissions. The first is the result of the volatilization of N in manure (as NH<sub>3</sub> and NO<sub>x</sub>) and the subsequent deposition of these gases and their products (NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>) onto soils and the surface of lakes and other waters. The second pathway is the runoff and leaching of N from manure into the groundwater below, into riparian zones receiving drain or runoff water, or into the ditches, streams, rivers, and estuaries into which the land drainage water eventually flows.

The production of direct N<sub>2</sub>O emissions from livestock manure depends on the composition of the manure (manure includes both feces and urine), the type of bacteria involved in the process, and the amount of oxygen and liquid in the manure system. For direct N<sub>2</sub>O emissions to occur, the manure must first be handled aerobically where organic N is mineralized or decomposed to NH<sub>4</sub> which is then nitrified to NO<sub>3</sub> (producing some N<sub>2</sub>O as a byproduct) (nitrification). Next, the manure must be handled anaerobically where the nitrate is then denitrified to N<sub>2</sub>O and N<sub>2</sub> (denitrification). NO<sub>x</sub> can also be produced during denitrification. (Groffman et al. 2000; Robertson and Groffman 2015). These emissions are most likely to occur in dry manure handling systems that have aerobic conditions, but that also contain pockets of anaerobic conditions due to saturation. A very small portion of the total N excreted is expected to convert to N<sub>2</sub>O in the waste management system (WMS). Indirect N<sub>2</sub>O emissions are produced when nitrogen is lost from the system through volatilization (as NH<sub>3</sub> or NO<sub>x</sub>) or through runoff and leaching. The vast majority of volatilization losses from these operations are NH<sub>3</sub>. Although there are also some small losses of NO<sub>x</sub>, there are no quantified estimates available for use, so losses due to volatilization are only based on NH<sub>3</sub> loss factors. Runoff losses would be expected from operations that house animals or store manure in a manner that is exposed to weather. Runoff losses are also specific to the type of animal housed on the operation due to differences in manure characteristics. Little information is known about leaching from manure management systems as most research focuses on leaching from land application systems. Since leaching losses are expected to be minimal, leaching losses are coupled with runoff losses and the runoff/leaching estimate provided in this chapter does not account for any leaching losses.

Estimates of CH<sub>4</sub> emissions from manure management in 2017 were 61.7 MMT CO<sub>2</sub> Eq. (2,467 kt); in 1990, emissions were 37.1 MMT CO<sub>2</sub> Eq. (1,486 kt). This represents a 66 percent increase in emissions from 1990.

---

<sup>4</sup> Direct and indirect N<sub>2</sub>O emissions from dung and urine spread onto fields either directly as daily spread or after it is removed from manure management systems (i.e., lagoon, pit, etc.) and from livestock dung and urine deposited on pasture, range, or paddock lands are accounted for and discussed in the Agricultural Soil Management source category within the Agriculture sector.

Emissions increased on average by 1.0 MMT CO<sub>2</sub> Eq. (2.0 percent) annually over this period. The majority of this increase is due to swine and dairy cow manure, where emissions increased 29 and 134 percent, respectively. From 2016 to 2017, there was a 0.2 percent increase in total CH<sub>4</sub> emissions from manure management, due to an increase in animal populations.

Although a large quantity of managed manure in the United States is handled as a solid, producing little CH<sub>4</sub>, the general trend in manure management, particularly for dairy cattle and swine (which are both shifting towards larger facilities), is one of increasing use of liquid systems. Also, new regulations controlling the application of manure nutrients to land have shifted manure management practices at smaller dairies from daily spread systems to storage and management of the manure on site. In many cases, manure management systems with the most substantial methane emissions are those associated with confined animal management operations where manure is handled in liquid-based systems. Nitrous oxide emissions from manure management vary significantly between the types of management system used and can also result in indirect emissions due to other forms of nitrogen loss from the system (IPCC 2006).

While national dairy animal populations have decreased since 1990, some states have seen increases in their dairy cattle populations as the industry becomes more concentrated in certain areas of the country and the number of animals contained on each facility increases. These areas of concentration, such as California, New Mexico, and Idaho, tend to utilize more liquid-based systems to manage (flush or scrape) and store manure. Thus, the shift toward larger dairy cattle and swine facilities since 1990 has translated into an increasing use of liquid manure management systems, which have higher potential CH<sub>4</sub> emissions than dry systems. This significant shift in both the dairy cattle and swine industries was accounted for by incorporating state and WMS-specific CH<sub>4</sub> conversion factor (MCF) values in combination with the 1992, 1997, 2002, 2007 and 2012 farm-size distribution data reported in the U.S. Department of Agriculture (USDA) *Census of Agriculture* (USDA 2016c).

In 2017, total N<sub>2</sub>O emissions from manure management were estimated to be 18.7 MMT CO<sub>2</sub> Eq. (63 kt); in 1990, emissions were 14.0 MMT CO<sub>2</sub> Eq. (47 kt). These values include both direct and indirect N<sub>2</sub>O emissions from manure management. Nitrous oxide emissions have increased since 1990. Small changes in N<sub>2</sub>O emissions from individual animal groups exhibit the same trends as the animal group populations, with the overall net effect that N<sub>2</sub>O emissions showed a 34 percent increase from 1990 to 2017 and a 3 percent increase from 2016 through 2017. Overall shifts toward liquid systems have driven down the emissions per unit of nitrogen excreted as dry manure handling systems have greater aerobic conditions that promote N<sub>2</sub>O emissions.

Table 5-6 and Table 5-7 provide estimates of CH<sub>4</sub> and N<sub>2</sub>O emissions from manure management by animal category.

**Table 5-6: CH<sub>4</sub> and N<sub>2</sub>O Emissions from Manure Management (MMT CO<sub>2</sub> Eq.)**

Gas/Animal Type	1990	2005	2013	2014	2015	2016	2017
<b>CH<sub>4</sub><sup>a</sup></b>	<b>37.1</b>	<b>53.7</b>	<b>58.1</b>	<b>57.8</b>	<b>60.9</b>	<b>61.5</b>	<b>61.7</b>
Dairy Cattle	14.7	26.4	33.4	34.0	34.8	34.4	34.5
Beef Cattle	3.1	3.3	3.1	3.0	3.1	3.3	3.4
Swine	15.5	20.3	18.0	17.2	19.2	20.2	20.0
Sheep	0.2	0.1	0.1	0.1	0.1	0.1	0.1
Goats	+	+	+	+	+	+	+
Poultry	3.3	3.2	3.2	3.3	3.4	3.4	3.4
Horses	0.2	0.3	0.2	0.2	0.2	0.2	0.2
American Bison	+	+	+	+	+	+	+
Mules and Asses	+	+	+	+	+	+	+
<b>N<sub>2</sub>O<sup>b</sup></b>	<b>14.0</b>	<b>16.5</b>	<b>17.4</b>	<b>17.4</b>	<b>17.6</b>	<b>18.2</b>	<b>18.7</b>
Dairy Cattle	5.3	5.6	5.9	5.9	6.1	6.1	6.1
Beef Cattle	5.9	7.2	7.7	7.8	7.7	8.1	8.6
Swine	1.2	1.6	1.8	1.7	1.8	1.9	1.9
Sheep	0.1	0.3	0.3	0.3	0.3	0.3	0.3
Goats	+	+	+	+	+	+	+
Poultry	1.4	1.6	1.6	1.6	1.6	1.6	1.6
Horses	0.1	0.1	0.1	0.1	0.1	0.1	0.1
American Bison <sup>c</sup>	NA	NA	NA	NA	NA	NA	NA
Mules and Asses	+	+	+	+	+	+	+

<b>Total</b>	<b>51.1</b>	<b>70.2</b>	<b>75.5</b>	<b>75.2</b>	<b>78.5</b>	<b>79.7</b>	<b>80.4</b>
--------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

NA (Not Available)

<sup>a</sup> Accounts for CH<sub>4</sub> reductions due to capture and destruction of CH<sub>4</sub> at facilities using anaerobic digesters.

<sup>b</sup> Includes both direct and indirect N<sub>2</sub>O emissions.

<sup>c</sup> There are no American bison N<sub>2</sub>O emissions from managed systems; American bison are maintained entirely on pasture, range, and paddock.

Notes: Emissions from manure deposited on pasture are included in the Agricultural Soils Management sector. Totals may not sum due to independent rounding.

1 **Table 5-7: CH<sub>4</sub> and N<sub>2</sub>O Emissions from Manure Management (kt)**

<b>Gas/Animal Type</b>	<b>1990</b>	<b>2005</b>	<b>2013</b>	<b>2014</b>	<b>2015</b>	<b>2016</b>	<b>2017</b>
<b>CH<sub>4</sub><sup>a</sup></b>	<b>1,486</b>	<b>2,150</b>	<b>2,322</b>	<b>2,311</b>	<b>2,435</b>	<b>2,461</b>	<b>2,467</b>
Dairy Cattle	590	1,057	1,338	1,360	1,390	1,374	1,381
Beef Cattle	126	133	122	120	126	131	135
Swine	622	812	721	688	770	807	802
Sheep	7	3	3	3	3	3	3
Goats	1	1	1	1	1	1	1
Poultry	131	129	129	132	136	136	137
Horses	9	12	9	9	9	9	8
American Bison	+	+	+	+	+	+	+
Mules and Asses	+	+	+	+	+	+	+
<b>N<sub>2</sub>O<sup>b</sup></b>	<b>47</b>	<b>55</b>	<b>58</b>	<b>58</b>	<b>59</b>	<b>61</b>	<b>63</b>
Dairy Cattle	18	19	20	20	20	21	21
Beef Cattle	20	24	26	26	26	27	29
Swine	4	5	6	6	6	6	7
Sheep	+	1	1	1	1	1	1
Goats	+	+	+	+	+	+	+
Poultry	5	5	5	5	5	5	5
Horses	+	+	+	+	+	+	+
American Bison <sup>c</sup>	NA	NA	NA	NA	NA	NA	NA
Mules and Asses	+	+	+	+	+	+	+

+ Does not exceed 0.5 kt.

NA (Not available)

<sup>a</sup> Accounts for CH<sub>4</sub> reductions due to capture and destruction of CH<sub>4</sub> at facilities using anaerobic digesters.

<sup>b</sup> Includes both direct and indirect N<sub>2</sub>O emissions.

<sup>c</sup> There are no American bison N<sub>2</sub>O emissions from managed systems; American bison are maintained entirely on pasture, range, and paddock.

Notes: Emissions from manure deposited on pasture are included in the Agricultural Soils Management sector. Totals may not sum due to independent rounding.

## 2 Methodology

3 The methodologies presented in IPCC (2006) form the basis of the CH<sub>4</sub> and N<sub>2</sub>O emission estimates for each animal  
4 type. This section presents a summary of the methodologies used to estimate CH<sub>4</sub> and N<sub>2</sub>O emissions from manure  
5 management. See Annex 3.11 for more detailed information on the methodology and data used to calculate CH<sub>4</sub> and  
6 N<sub>2</sub>O emissions from manure management.

### 7 Methane Calculation Methods

8 The following inputs were used in the calculation of manure management CH<sub>4</sub> emissions for 1990 through 2017:

- 9
  - Animal population data (by animal type and state);

- Typical animal mass (TAM) data (by animal type);
- Portion of manure managed in each WMS, by state and animal type;
- Volatile solids (VS) production rate (by animal type and state or United States);
- Methane producing potential ( $B_0$ ) of the volatile solids (by animal type); and
- Methane conversion factors (MCF), the extent to which the  $CH_4$  producing potential is realized for each type of WMS (by state and manure management system, including the impacts of any biogas collection efforts).

Methane emissions were estimated by first determining activity data, including animal population, TAM, WMS usage, and waste characteristics. The activity data sources are described below:

- Annual animal population data for 1990 through 2017 for all livestock types, except goats, horses, mules and asses, and American bison were obtained from the USDA-NASS. For cattle, the USDA populations were utilized in conjunction with birth rates, detailed feedlot placement information, and slaughter weight data to create the transition matrix in the Cattle Enteric Fermentation Model (CEFM) that models cohorts of individual animal types and their specific emission profiles. The key variables tracked for each of the cattle population categories are described in Section 5.1 and in more detail in Annex 3.10. Goat population data for 1992, 1997, 2002, 2007, and 2012; horse and mule and ass population data for 1987, 1992, 1997, 2002, 2007, and 2012; and American bison population for 2002, 2007 and 2012 were obtained from the *Census of Agriculture* (USDA 2014a). American bison population data for 1990 through 1999 were obtained from the National Bison Association (1999).
- The TAM is an annual average weight that was obtained for animal types other than cattle from information in USDA's *Agricultural Waste Management Field Handbook* (USDA 1996), the American Society of Agricultural Engineers, Standard D384.1 (ASAE 1998) and others (Meagher 1986; EPA 1992; Safley 2000; ERG 2003b; IPCC 2006; ERG 2010a). For a description of the TAM used for cattle, see Annex 3.10.
- WMS usage was estimated for swine and dairy cattle for different farm size categories using state and regional data from USDA (USDA APHIS 1996; Bush 1998; Ott 2000; USDA 2016c) and EPA (ERG 2000a; EPA 2002a and 2002b; ERG 2018). For beef cattle and poultry, manure management system usage data were not tied to farm size but were based on other data sources (ERG 2000a; USDA APHIS 2000; UEP 1999). For other animal types, manure management system usage was based on previous estimates (EPA 1992). American bison WMS usage was assumed to be the same as not on feed (NOF) cattle, while mules and asses were assumed to be the same as horses.
- VS production rates for all cattle except for calves were calculated by head for each state and animal type in the CEFM. VS production rates by animal mass for all other animals were determined using data from USDA's *Agricultural Waste Management Field Handbook* (USDA 1996 and 2008; ERG 2010b and 2010c) and data that was not available in the most recent *Handbook* were obtained from the American Society of Agricultural Engineers, Standard D384.1 (ASAE 1998) or the *2006 IPCC Guidelines* (IPCC 2006). American bison VS production was assumed to be the same as NOF bulls.
- The maximum  $CH_4$ -producing capacity of the VS ( $B_0$ ) was determined for each animal type based on literature values (Morris 1976; Bryant et al. 1976; Hashimoto 1981; Hashimoto 1984; EPA 1992; Hill 1982; Hill 1984).
- MCFs for dry systems were set equal to default IPCC factors based on state climate for each year (IPCC 2006). MCFs for liquid/slurry, anaerobic lagoon, and deep pit systems were calculated based on the forecast performance of biological systems relative to temperature changes as predicted in the van't Hoff-Arrhenius equation which is consistent with IPCC (2006) Tier 2 methodology.
- Data from anaerobic digestion systems with  $CH_4$  capture and combustion were obtained from the EPA AgSTAR Program, including information presented in the *AgSTAR Digest* (EPA 2000, 2003, 2006) and the AgSTAR project database (EPA 2018). Anaerobic digester emissions were calculated based on estimated methane production and collection and destruction efficiency assumptions (ERG 2008).
- For all cattle except for calves, the estimated amount of VS (kg per animal-year) managed in each WMS for each animal type, state, and year were taken from the CEFM, assuming American bison VS production to be the same as NOF bulls. For animals other than cattle, the annual amount of VS (kg per year) from

manure excreted in each WMS was calculated for each animal type, state, and year. This calculation multiplied the animal population (head) by the VS excretion rate (kg VS per 1,000 kg animal mass per day), the TAM (kg animal mass per head) divided by 1,000, the WMS distribution (percent), and the number of days per year (365.25).

The estimated amount of VS managed in each WMS was used to estimate the CH<sub>4</sub> emissions (kg CH<sub>4</sub> per year) from each WMS. The amount of VS (kg per year) were multiplied by the maximum CH<sub>4</sub> producing capacity of the VS (B<sub>0</sub>) (m<sup>3</sup> CH<sub>4</sub> per kg VS), the MCF for that WMS (percent), and the density of CH<sub>4</sub> (kg CH<sub>4</sub> per m<sup>3</sup> CH<sub>4</sub>). The CH<sub>4</sub> emissions for each WMS, state, and animal type were summed to determine the total U.S. CH<sub>4</sub> emissions.

## Nitrous Oxide Calculation Methods

The following inputs were used in the calculation of direct and indirect manure management N<sub>2</sub>O emissions for 1990 through 2017:

- Animal population data (by animal type and state);
- TAM data (by animal type);
- Portion of manure managed in each WMS (by state and animal type);
- Total Kjeldahl N excretion rate (N<sub>ex</sub>);
- Direct N<sub>2</sub>O emission factor (EF<sub>WMS</sub>);
- Indirect N<sub>2</sub>O emission factor for volatilization (EF<sub>volatilization</sub>);
- Indirect N<sub>2</sub>O emission factor for runoff and leaching (EF<sub>runoff/leach</sub>);
- Fraction of N loss from volatilization of NH<sub>3</sub> and NO<sub>x</sub> (Frac<sub>gas</sub>); and
- Fraction of N loss from runoff and leaching (Frac<sub>runoff/leach</sub>).

Nitrous oxide emissions were estimated by first determining activity data, including animal population, TAM, WMS usage, and waste characteristics. The activity data sources (except for population, TAM, and WMS, which were described above) are described below:

- Nex rates for all cattle except for calves were calculated by head for each state and animal type in the CEFM. Nex rates by animal mass for all other animals were determined using data from USDA's *Agricultural Waste Management Field Handbook* (USDA 1996 and 2008; ERG 2010b and 2010c) and data from the American Society of Agricultural Engineers, Standard D384.1 (ASAE 1998) and IPCC (2006). American bison Nex rates were assumed to be the same as NOF bulls.<sup>5</sup>
- All N<sub>2</sub>O emission factors (direct and indirect) were taken from IPCC (2006). These data are appropriate because they were developed using U.S. data.
- Country-specific estimates for the fraction of N loss from volatilization (Frac<sub>gas</sub>) and runoff and leaching (Frac<sub>runoff/leach</sub>) were developed. Frac<sub>gas</sub> values were based on WMS-specific volatilization values as estimated from EPA's *National Emission Inventory - Ammonia Emissions from Animal Agriculture Operations* (EPA 2005). Frac<sub>runoff/leaching</sub> values were based on regional cattle runoff data from EPA's Office of Water (EPA 2002b; see Annex 3.11).

To estimate N<sub>2</sub>O emissions for cattle (except for calves), the estimated amount of N excreted (kg per animal-year) that is managed in each WMS for each animal type, state, and year were taken from the CEFM. For calves and other animals, the amount of N excreted (kg per year) in manure in each WMS for each animal type, state, and year was calculated. The population (head) for each state and animal was multiplied by TAM (kg animal mass per head) divided by 1,000, the nitrogen excretion rate (N<sub>ex</sub>, in kg N per 1,000 kg animal mass per day), WMS distribution (percent), and the number of days per year.

Direct N<sub>2</sub>O emissions were calculated by multiplying the amount of N excreted (kg per year) in each WMS by the N<sub>2</sub>O direct emission factor for that WMS (EF<sub>WMS</sub>, in kg N<sub>2</sub>O-N per kg N) and the conversion factor of N<sub>2</sub>O-N to

---

<sup>5</sup> The N<sub>2</sub>O emissions from N excreted (N<sub>ex</sub>) by American bison on grazing lands are accounted for and discussed in the Agricultural Soil Management source category and included under pasture, range and paddock (PRP) emissions. Because American bison are maintained entirely on unmanaged WMS and N<sub>2</sub>O emissions from unmanaged WMS are not included in the Manure Management source category, there are no N<sub>2</sub>O emissions from American bison included in the Manure Management source category.

N<sub>2</sub>O. These emissions were summed over state, animal, and WMS to determine the total direct N<sub>2</sub>O emissions (kg of N<sub>2</sub>O per year).

Next, indirect N<sub>2</sub>O emissions from volatilization (kg N<sub>2</sub>O per year) were calculated by multiplying the amount of N excreted (kg per year) in each WMS by the fraction of N lost through volatilization ( $\text{Frac}_{\text{tas}}$ ) divided by 100, the emission factor for volatilization ( $\text{EF}_{\text{volatilization}}$ , in kg N<sub>2</sub>O per kg N), and the conversion factor of N<sub>2</sub>O-N to N<sub>2</sub>O. Indirect N<sub>2</sub>O emissions from runoff and leaching (kg N<sub>2</sub>O per year) were then calculated by multiplying the amount of N excreted (kg per year) in each WMS by the fraction of N lost through runoff and leaching ( $\text{Frac}_{\text{runoff/leach}}$ ) divided by 100, and the emission factor for runoff and leaching ( $\text{EF}_{\text{runoff/leach}}$ , in kg N<sub>2</sub>O per kg N), and the conversion factor of N<sub>2</sub>O-N to N<sub>2</sub>O. The indirect N<sub>2</sub>O emissions from volatilization and runoff and leaching were summed to determine the total indirect N<sub>2</sub>O emissions.

Following these steps, direct and indirect N<sub>2</sub>O emissions were summed to determine total N<sub>2</sub>O emissions (kg N<sub>2</sub>O per year) for the years 1990 to 2017.

# Uncertainty and Time-Series Consistency

An analysis (ERG 2003a) was conducted for the manure management emission estimates presented in the 1990 through 2001 Inventory (i.e., 2003 submission to the UNFCCC) to determine the uncertainty associated with estimating CH<sub>4</sub> and N<sub>2</sub>O emissions from livestock manure management. The quantitative uncertainty analysis for this source category was performed in 2002 through the IPCC-recommended Approach 2 uncertainty estimation methodology, the Monte Carlo Stochastic Simulation technique. The uncertainty analysis was developed based on the methods used to estimate CH<sub>4</sub> and N<sub>2</sub>O emissions from manure management systems. A normal probability distribution was assumed for each source data category. The series of equations used were condensed into a single equation for each animal type and state. The equations for each animal group contained four to five variables around which the uncertainty analysis was performed for each state. These uncertainty estimates were directly applied to the 2017 emission estimates as there have not been significant changes in the methodology since that time.

The results of the Approach 2 quantitative uncertainty analysis are summarized in Table 5-8. Manure management CH<sub>4</sub> emissions in 2017 were estimated to be between 50.6 and 74.0 MMT CO<sub>2</sub> Eq. at a 95 percent confidence level, which indicates a range of 18 percent below to 20 percent above the actual 2017 emission estimate of 61.7 MMT CO<sub>2</sub> Eq. At the 95 percent confidence level, N<sub>2</sub>O emissions were estimated to be between 15.7 and 23.2 MMT CO<sub>2</sub> Eq. (or approximately 16 percent below and 24 percent above the actual 2017 emission estimate of 18.7 MMT CO<sub>2</sub> Eq.).

**Table 5-8: Approach 2 Quantitative Uncertainty Estimates for CH<sub>4</sub> and N<sub>2</sub>O (Direct and Indirect) Emissions from Manure Management (MMT CO<sub>2</sub> Eq. and Percent)**

Source	Gas	2017 Emission Estimate (MMT CO <sub>2</sub> Eq.)	Uncertainty Range Relative to Emission Estimate <sup>a</sup>			
			(MMT CO <sub>2</sub> Eq.)		(%)	
			Lower Bound	Upper Bound	Lower Bound	Upper Bound
Manure Management	CH <sub>4</sub>	61.7	50.6	74.0	-18%	20%
Manure Management	N <sub>2</sub> O	18.7	15.7	23.2	-16%	24%

<sup>a</sup> Range of emission estimates predicted by Monte Carlo Stochastic Simulation for a 95 percent confidence interval.

## QA/QC and Verification

Tier 1 and Tier 2 QA/QC activities were conducted consistent with the U.S. QA/QC plan. Tier 2 activities focused on comparing estimates for the previous and current Inventories for N<sub>2</sub>O emissions from managed systems and CH<sub>4</sub> emissions from livestock manure. All errors identified were corrected. Order of magnitude checks were also conducted, and corrections made where needed. In addition, manure N data were checked by comparing state-level data with bottom up estimates derived at the county level and summed to the state level. Similarly, a comparison was made by animal and WMS type for the full time series, between national level estimates for N excreted and the sum of county estimates for the full time series.

Time-series data, including population, are validated by experts to ensure they are representative of the best available U.S.-specific data. The U.S.-specific values for TAM, Nex, VS, B<sub>o</sub>, and MCF were also compared to the IPCC default values and validated by experts. Although significant differences exist in some instances, these differences are due to the use of U.S.-specific data and the differences in U.S. agriculture as compared to other countries. The U.S. manure management emission estimates use the most reliable country-specific data, which are more representative of U.S. animals and systems than the IPCC (2006) default values.

For additional verification of the 1990 to 2017 estimates, the implied CH<sub>4</sub> emission factors for manure management (kg of CH<sub>4</sub> per head per year) were compared against the default IPCC (2006) values. Table 5-9 presents the implied emission factors of kg of CH<sub>4</sub> per head per year used for the manure management emission estimates as well as the IPCC (2006) default emission factors. The U.S. implied emission factors fall within the range of the IPCC (2006) default values, except in the case of sheep, goats, and some years for horses and dairy cattle. The U.S. implied emission factors are greater than the IPCC (2006) default value for those animals due to the use of U.S.-specific data for typical animal mass and VS excretion. There is an increase in implied emission factors for dairy cattle and swine across the time series. This increase reflects the dairy cattle and swine industry trend towards larger farm sizes; large farms are more likely to manage manure as a liquid and therefore produce more CH<sub>4</sub> emissions.

**Table 5-9: IPCC (2006) Implied Emission Factor Default Values Compared with Calculated Values for CH<sub>4</sub> from Manure Management (kg/head/year)**

Animal Type	IPCC Default CH <sub>4</sub> Emission Factors (kg/head/year)	Implied CH <sub>4</sub> Emission Factors (kg/head/year)						
		1990	2005	2013	2014	2015	2016	2017
Dairy Cattle	48-112	30.2	59.4	72.3	73.4	73.9	72.9	73.1
Beef Cattle	1-2	1.5	1.6	1.6	1.6	1.7	1.7	1.7
Swine	10-45	11.5	13.3	11.0	10.7	11.3	11.5	11.1
Sheep	0.19-0.37	0.6	0.6	0.5	0.5	0.5	0.5	0.5
Goats	0.13-0.26	0.4	0.3	0.3	0.3	0.3	0.3	0.3
Poultry	0.02-1.4	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Horses	1.56-3.13	4.3	3.1	2.5	2.5	2.6	2.6	2.6
American Bison	NA	1.8	2.0	2.0	2.0	2.1	2.1	2.1
Mules and Asses	0.76-1.14	0.9	1.0	0.9	0.9	1.0	1.0	1.0
NA (Not Applicable)								

In addition, default IPCC (2006) emission factors for N<sub>2</sub>O were compared to the U.S. Inventory implied N<sub>2</sub>O emission factors. Default N<sub>2</sub>O emission factors from the *2006 IPCC Guidelines* were used to estimate N<sub>2</sub>O emission from each WMS in conjunction with U.S.-specific Nex values. The implied emission factors differed from the U.S. Inventory values due to the use of U.S.-specific Nex values and differences in populations present in each WMS throughout the time series.

## Recalculations Discussion

The manure management emission estimates include the following recalculations relative to the previous Inventory:

- The CEFM produces population, VS and Nex data for cattle (except calves), that are used in the manure management inventory. As a result, all changes to the CEFM described in Section 5.1 contributed to changes in the population, VS and Nex data used for calculating CH<sub>4</sub> and N<sub>2</sub>O cattle emissions from manure management.



- State animal populations were updated to reflect updated USDA NASS datasets, which resulted in population changes for:
  - Poultry in 2015,
  - Calves in 2015,
  - Beef OF heifers and steers in 2015,
  - Dairy heifers in 2015,
  - American bison in 2012-2015; and
  - Swine in 2015 (USDA 2018).
- WMS distribution data for swine were updated with data from the 2009 USDA Agricultural Resource Management Survey (ARMS) of swine producers (ERG 2018). Anaerobic digestion data were also updated for swine, using data from EPA’s AgSTAR Program (EPA 2018).
- Temperature data were updated by NOAA due to an update in their computer program which affected the precision of the output dataset (NOAA 2018). This resulted in minor temperature changes and subsequently, MCF changes for all animals across the time series.

These changes impacted total emission estimates for 1990 through 2016, overall decreasing annual estimations from less than 1 percent to 7.2 percent across the time series. The most significant changes were to the swine emissions estimates, resulting primarily from the swine WMS update. Total swine annual estimations decreased throughout the entire time series, but most significantly for 2013 through 2015 during which time they decreased by over 20 percent.

## Planned Improvements

During the Public Review period of the previous Inventory report (i.e., 1990 through 2016), EPA received comment on various aspects of the manure management inventory, including recommended improvements to clarify the scope of the manure management sector and better align terminology with those used within the industry (e.g., clarifying “managed” versus “unmanaged”), as well as comments to update data and methods which reiterated those improvements already identified by EPA and listed below. EPA notes that many of these improvements, identified below, are major updates and may take multiple years to implement in full, but will add clarity to improve the transparency of future inventories.

Potential improvements for future Inventory years include:

- Continuing to obtain and incorporate existing data sources (such as the 2016 USDA ARMS dairy data) to update WMS distributions.
- Revising the methodology for population distribution to states where USDA population data are withheld due to disclosure concerns.
- Revising the anaerobic digestion estimates to estimate CH<sub>4</sub> emissions reductions due to the use of anaerobic digesters (the Inventory currently estimates only emissions from anaerobic digestion systems).
- Investigating improved emissions estimate methodologies for swine pit systems with less than one month of storage (the new swine WMS data included this WMS category).
- Updating the B<sub>0</sub> data used in the Inventory, which are dated.
- Comparing CH<sub>4</sub> and N<sub>2</sub>O emission estimates with estimates from other models and more recent studies and compare the results to the Inventory, such as USDA’s Dairy Gas Emissions Model.
- Comparing manure management emission estimates with on-farm measurement data to identify opportunities for improved estimates.
- Improving collaboration with the Enteric Fermentation source category estimates. For future inventories, it may be beneficial to have the CEFM and Manure Management calculations in the same model, as they rely on much of the same activity data and they depend on each other’s outputs to properly calculation emissions.
- Implementing a methodology to calculate monthly emissions estimates to present data that show seasonal changes in emissions from each WMS.
- Revising the uncertainty analysis to address changes that have been implemented to the CH<sub>4</sub> and N<sub>2</sub>O estimates.

## 5.3 Rice Cultivation (CRF Source Category 3C)

---

Most of the world's rice is grown on flooded fields (Baich 2013), and flooding creates anaerobic conditions that foster CH<sub>4</sub> production through a process known as methanogenesis. Approximately 60 to 90 percent of the CH<sub>4</sub> produced by methanogenic bacteria is oxidized in the soil and converted to CO<sub>2</sub> by methanotrophic bacteria. The remainder is emitted to the atmosphere (Holzapfel-Pschorn et al. 1985; Sass et al. 1990) or transported as dissolved CH<sub>4</sub> into groundwater and waterways (Neue et al. 1997). Methane is transported to the atmosphere primarily through the rice plants, but some CH<sub>4</sub> also escapes via ebullition (i.e., bubbling through the water) and to a much lesser extent by diffusion through the water (van Bodegom et al. 2001).

Water management is arguably the most important factor affecting CH<sub>4</sub> emissions, and improved water management has the largest potential to mitigate emissions (Yan et al. 2009). Upland rice fields are not flooded, and therefore do not produce CH<sub>4</sub>, but large amounts of CH<sub>4</sub> can be emitted in continuously irrigated fields, which is the most common practices in the United States (USDA 2012). Single or multiple aeration events with drainage of a field during the growing season can significantly reduce these emissions (Wassmann et al. 2000a), but drainage may also increase N<sub>2</sub>O emissions. Deepwater rice fields (i.e., fields with flooding depths greater than one meter, such as natural wetlands) tend to have less living stems reaching the soil, thus reducing the amount of CH<sub>4</sub> transport to the atmosphere through the plant compared to shallow-flooded systems (Sass 2001).

Other management practices also influence CH<sub>4</sub> emissions from flooded rice fields including rice residue straw management and application of organic amendments, in addition to cultivar selection due to differences in the amount of root exudates<sup>6</sup> among rice varieties (Neue et al. 1997). These practices influence the amount of organic matter available for methanogenesis, and some practices, such as mulching rice straw or composting organic amendments, can reduce the amount of labile carbon and limit CH<sub>4</sub> emissions (Wassmann et al. 2000b). Fertilization practices also influence CH<sub>4</sub> emissions, particularly the use of fertilizers with sulfate (Wassmann et al. 2000b; Linquist et al. 2012), which can reduce CH<sub>4</sub> emissions. Other environmental variables also impact the methanogenesis process such as soil temperature and soil type. Soil temperature regulates the activity of methanogenic bacteria which in turn affects the rate of CH<sub>4</sub> production. Soil texture influences decomposition of soil organic matter, but is also thought to have an impact on oxidation of CH<sub>4</sub> in the soil (Sass et al. 1994).

Rice is currently cultivated in twelve states, including Arkansas, California, Florida, Illinois, Kentucky, Louisiana, Minnesota, Mississippi, Missouri, New York, South Carolina, Tennessee and Texas. Soil types, rice varieties, and cultivation practices vary across the United States, but most farmers apply fertilizers and do not harvest crop residues. In addition, a second, ratoon rice crop is sometimes grown in the Southeastern region of the country. Ratoon crops are produced from regrowth of the stubble remaining after the harvest of the first rice crop. Methane emissions from ratoon crops are higher than those from the primary crops due to the increased amount of labile organic matter available for anaerobic decomposition in the form of relatively fresh crop residue straw. Emissions tend to be higher in rice fields if the residues have been in the field for less than 30 days before planting the next rice crop (Lindau and Bollich 1993; IPCC 2006; Wang et al. 2013).

A combination of Tier 1 and 3 methods are used to estimate CH<sub>4</sub> emissions from rice cultivation across most of the time series, while a surrogate data method has been applied to estimate national emissions for 2013-2017 in this Inventory. National emission estimates based on surrogate data will be recalculated in the next (i.e., 1990 through 2018) Inventory submission using the Tier 1 and 3 methods.

Overall, rice cultivation is a minor source of CH<sub>4</sub> emissions in the United States relative to other source categories (see Table 5-10, Table 5-11, and Figure 5-2). The majority of emissions occur in Arkansas, California, Louisiana and Texas. In 2017, CH<sub>4</sub> emissions from rice cultivation were 11.3 MMT CO<sub>2</sub> Eq. (454 kt). Annual emissions fluctuate between 1990 and 2017, which is largely due to differences in the amount of rice harvested areas over time, which has been decreasing over the past two decades. Consequently, emissions in 2017 are 29 percent lower than emissions in 1990.

---

<sup>6</sup> The roots of rice plants add organic material to the soil through a process called "root exudation." Root exudation is thought to enhance decomposition of the soil organic matter and release nutrients that the plant can absorb and use to stimulate more production. The amount of root exudate produced by a rice plant over a growing season varies among rice varieties.

1 **Table 5-10: CH<sub>4</sub> Emissions from Rice Cultivation (MMT CO<sub>2</sub> Eq.)**

State	1990	2005	2012	2013	2014	2015	2016	2017
Arkansas	3.3	4.7	3.8	NE	NE	NE	NE	NE
California	2.0	2.1	2.0	NE	NE	NE	NE	NE
Florida	0.0	0.1	0.0	NE	NE	NE	NE	NE
Illinois	0.0	+	0.0	NE	NE	NE	NE	NE
Kentucky	0.0	+	0.0	NE	NE	NE	NE	NE
Louisiana	6.1	6.5	3.9	NE	NE	NE	NE	NE
Minnesota	+	+	+	NE	NE	NE	NE	NE
Mississippi	0.6	0.6	0.5	NE	NE	NE	NE	NE
Missouri	0.3	0.6	0.3	NE	NE	NE	NE	NE
New York	+	0.0	0.0	NE	NE	NE	NE	NE
South Carolina	0.0	0.0	0.0	NE	NE	NE	NE	NE
Tennessee	0.0	+	0.0	NE	NE	NE	NE	NE
Texas	3.7	2.1	0.9	NE	NE	NE	NE	NE
<b>Total</b>	<b>16.0</b>	<b>16.7</b>	<b>11.3</b>	<b>11.5</b>	<b>12.7</b>	<b>12.3</b>	<b>13.7</b>	<b>11.3</b>

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

NE (Not Estimated). State-level emissions are not estimated for 2013 through 2017 Inventory, and national emissions are determined using a surrogate data method.

Note: Totals may not sum due to independent rounding.

2 **Table 5-11: CH<sub>4</sub> Emissions from Rice Cultivation (kt)**


State	1990	2005	2012	2013	2014	2015	2016	2017
Arkansas	132	188	151	NE	NE	NE	NE	NE
California	81	82	81	NE	NE	NE	NE	NE
Florida	0	3	0	NE	NE	NE	NE	NE
Illinois	0	+	0	NE	NE	NE	NE	NE
Kentucky	0	+	0	NE	NE	NE	NE	NE
Louisiana	246	261	156	NE	NE	NE	NE	NE
Minnesota	1	2	1	NE	NE	NE	NE	NE
Mississippi	23	23	19	NE	NE	NE	NE	NE
Missouri	12	22	12	NE	NE	NE	NE	NE
New York	+	0	0	NE	NE	NE	NE	NE
South Carolina	0	0	0	NE	NE	NE	NE	NE
Tennessee	0	+	0	NE	NE	NE	NE	NE
Texas	146	86	34	NE	NE	NE	NE	NE
<b>Total</b>	<b>641</b>	<b>667</b>	<b>453</b>	<b>462</b>	<b>510</b>	<b>493</b>	<b>549</b>	<b>454</b>

+ Does not exceed 0.5 kt.

NE (Not Estimated). State-level emissions are not estimated for 2013 through 2017 Inventory, and national emissions are determined using a surrogate data method.

Note: Totals may not sum due to independent rounding.

**Figure 5-2: Annual CH<sub>4</sub> Emissions from Rice Cultivation, 2012 (MMT CO<sub>2</sub> Eq./Year)**



Note: Only national-scale emissions are estimated for 2013 through 2017 in this Inventory using the surrogate data method described in the Methodology section; therefore the fine-scale emission patterns in this map are based on the estimates for 2012.

## Methodology

The methodology used to estimate CH<sub>4</sub> emissions from rice cultivation is based on a combination of IPCC Tier 1 and 3 approaches. The Tier 3 method utilizes a process-based model (DAYCENT) to estimate CH<sub>4</sub> emissions from rice cultivation (Cheng et al. 2013), and has been tested in the United States (see Annex 3.12) and Asia (Cheng et al. 2013, 2014). The model simulates hydrological conditions and thermal regimes, organic matter decomposition, root exudation, rice plant growth and its influence on oxidation of CH<sub>4</sub>, as well as CH<sub>4</sub> transport through the plant and via ebullition (Cheng et al. 2013). The method simulates the influence of organic amendments, rice straw management on methanogenesis in the flooded soils, and ratooning of rice crops with a second harvest during the growing season. In addition to CH<sub>4</sub> emissions, DAYCENT simulates soil C stock changes and N<sub>2</sub>O emissions (Parton et al. 1987 and 1998; Del Grosso et al. 2010), and allows for a seamless set of simulations for crop rotations that include both rice and non-rice crops.

The Tier 1 method is applied to estimate CH<sub>4</sub> emissions from rice when grown in rotation with crops that are not simulated by DAYCENT, such as vegetable crops. The Tier 1 method is also used for areas converted between agriculture (i.e., cropland and grassland) and other land uses, such as forest land, wetland, and settlements. In addition, the Tier 1 method is used to estimate CH<sub>4</sub> emissions from organic soils (i.e., Histosols) and from areas with very gravelly, cobbly, or shaley soils (greater than 35 percent by volume). The Tier 3 method using DAYCENT has not been fully tested for estimating emissions associated with these crops and rotations, land uses, as well as organic soils or cobbly, gravelly, and shaley mineral soils.

The Tier 1 method for estimating CH<sub>4</sub> emissions from rice production utilizes a default base emission rate and scaling factors (IPCC 2006). The base emission factor represents emissions for continuously flooded fields with no organic amendments. Scaling factors are used to adjust for water management and organic amendments that differ from continuous flooding with no organic amendments. The method accounts for pre-season and growing season flooding; types and amounts of organic amendments; and the number of rice production seasons within a single year.

(i.e., single cropping, ratooning, etc.). The Tier 1 analysis is implemented in the Agriculture and Land Use National Greenhouse Gas Inventory (ALU) software (Ogle et al. 2016).<sup>7</sup>

Rice cultivation areas are based on cropping and land use histories recorded in the USDA National Resources Inventory (NRI) survey (USDA-NRCS 2015). The NRI is a statistically-based sample of all non-federal land, and includes 380,956 survey points of which 1,588 are in locations with rice cultivation at the end of the NRI time series. The Tier 3 method is used to estimate CH<sub>4</sub> emissions from 1,393 of the NRI survey locations, and the remaining 195 survey locations are estimated with the Tier 1 method. Each NRI survey point is associated with an “expansion factor” that allows scaling of CH<sub>4</sub> emissions from NRI points to the entire country (i.e., each expansion factor represents the amount of area with the same land-use/management history as the sample point). Land-use and some management information in the NRI (e.g., crop type, soil attributes, and irrigation) were collected on a 5-year cycle beginning in 1982, along with cropping rotation data in 4 out of 5 years for each 5-year time period (i.e., 1979 to 1982, 1984 to 1987, 1989 to 1992, and 1994 to 1997). The NRI program began collecting annual data in 1998, with data currently available through 2012 (USDA-NRCS 2015). The current Inventory only uses NRI data through 2012 because newer data are not available, but will be incorporated when additional years of data are released by USDA-NRCS. The harvested rice areas in each state are presented in Table 5-12.

**Table 5-12: Rice Area Harvested (1,000 Hectares)**

State/Crop	1990	2005	2012	2013	2014	2015	2016	2017
Arkansas	599	796	613	NE	NE	NE	NE	NE
California	248	247	244	NE	NE	NE	NE	NE
Florida	0	11	0	NE	NE	NE	NE	NE
Illinois	0	0	0	NE	NE	NE	NE	NE
Kentucky	0	0	0	NE	NE	NE	NE	NE
Louisiana	380	402	226	NE	NE	NE	NE	NE
Minnesota	4	10	6	NE	NE	NE	NE	NE
Mississippi	119	115	92	NE	NE	NE	NE	NE
Missouri	47	93	46	NE	NE	NE	NE	NE
New York	1	0	0	NE	NE	NE	NE	NE
South Carolina	0	0	0	NE	NE	NE	NE	NE
Tennessee	0	1	0	NE	NE	NE	NE	NE
Texas	300	150	66	NE	NE	NE	NE	NE
<b>Total</b>	<b>1,698</b>	<b>1,826</b>	<b>1,292</b>	<b>NE</b>	<b>NE</b>	<b>NE</b>	<b>NE</b>	<b>NE</b>

NE (Not Estimated).

Notes: Totals may not sum due to independent rounding. States are included if NRI reports an area of rice production in the state at any time between 1990 and 2012. Rice harvested area data have not been compiled for 2013 to 2017.

The Southeastern states have sufficient growing periods for a ratoon crop in some years. For example, in Arkansas, the length of growing season is occasionally sufficient for ratoon crops on an average of 1 percent of the rice fields. No data are available about ratoon crops in Missouri or Mississippi, and the average amount of ratooning in Arkansas was assigned to these states. Ratoon cropping occurs much more frequently in Louisiana (LSU 2015 for years 2000 through 2013, 2015) and Texas (TAMU 2015 for years 1993 through 2014), averaging 32 percent and 45 percent of rice acres planted, respectively. Florida also has a large fraction of area with a ratoon crop (49 percent). Ratoon rice crops are not grown in California. Ratooned crop area as a percent of primary crop area is presented in Table 5-13.

**Table 5-13: Average Ratooned Area as Percent of Primary Growth Area (Percent)**

State	1990-2012
Arkansas <sup>a</sup>	1%
California	0%
Florida <sup>b</sup>	49%
Louisiana <sup>c</sup>	32%
Mississippi <sup>a</sup>	1%

<sup>7</sup> See <<http://www.nrel.colostate.edu/projects/ALUsoftware/>>.

Missouri <sup>a</sup>	1%
Texas <sup>d</sup>	45%

<sup>a</sup> Arkansas: 1990–2000 (Slaton 1999 through 2001); 2001–2011 (Wilson 2002 through 2007, 2009 through 2012); 2012–2013 (Hardke 2013, 2014).

<sup>b</sup> Florida - Ratoon: 1990–2000 (Schueneman 1997, 1999 through 2001); 2001 (Deren 2002); 2002–2003 (Kirstein 2003 through 2004, 2006); 2004 (Cantens 2004 through 2005); 2005–2013 (Gonzalez 2007 through 2014).

<sup>c</sup> Louisiana: 1990–2013 (Linscombe 1999, 2001 through 2014).

<sup>d</sup> Texas: 1990–2002 (Klosterboer 1997, 1999 through 2003); 2003–2004 (Stansel 2004 through 2005); 2005 (Texas Agricultural Experiment Station 2006); 2006–2013 (Texas Agricultural Experiment Station 2007 through 2014).

While rice crop production in the United States includes a minor amount of land with mid-season drainage or alternate wet-dry periods, the majority of rice growers use continuously flooded water management systems (Hardke 2015; UCCE 2015; Hollier 1999; Way et al. 2014). Therefore, continuous flooding was assumed in the DAYCENT simulations and the Tier 1 method. Variation in flooding can be incorporated in future Inventories if water management data are collected.

Winter flooding is another key practice associated with water management in rice fields, and the impact of winter flooding on CH<sub>4</sub> emissions is addressed in the Tier 3 and Tier 1 analyses. Flooding is used to prepare fields for the next growing season, and to create waterfowl habitat (Young 2013; Miller et al. 2010; Fleskes et al. 2005). Fitzgerald et al. (2000) suggests that as much as 50 percent of the annual emissions may occur during the winter flood. Winter flooding is a common practice with an average of 34 percent of fields managed with winter flooding in California (Miller et al. 2010; Fleskes et al. 2005), and approximately 21 percent of the fields managed with winter flooding in Arkansas (Wilson and Branson 2005 and 2006; Wilson and Runsick 2007 and 2008; Wilson et al. 2009 and 2010; Hardke and Wilson 2013 and 2014; Hardke 2015). No data are available on winter flooding for Texas, Louisiana, Florida, Missouri, or Mississippi. For these states, the average amount of flooding is assumed to be similar to Arkansas. In addition, the amount of flooding is assumed to be relatively constant over the Inventory time period.

A surrogate data method is used to estimate emissions from 2013 to 2017 associated with the rice CH<sub>4</sub> emissions for Tier 1 and 3 methods. Specifically, a linear regression model with autoregressive moving-average (ARMA) errors was used to estimate the relationship between the surrogate data and the 1990 through 2012 emissions data that was derived using the Tier 1 and 3 methods (Brockwell and Davis 2016). Surrogate data for this model are based on rice commodity statistics from USDA-NASS.<sup>8</sup> See Box 5-2 for more information about the surrogate data method.

## Box 5-2: Surrogate Data Method

An approach to extend the time series is needed to estimate emissions from Rice cultivation because there are gaps in activity data at the end of the time series. This is mainly due to the fact that the National Resources Inventory (NRI) does not release data every year, and the NRI is a key data source for estimating greenhouse gas emissions.

A surrogate data method has been selected to impute missing emissions at the end of the time series. A linear regression model with autoregressive moving-average (ARMA) errors (Brockwell and Davis, 2016) is used to estimate the relationship between the surrogate data and the observed 1990 to 2012 emissions data that has been compiled using the inventory methods described in this section. The model to extend the time series is given by

$$Y = X\beta + \varepsilon,$$

where Y is the response variable (e.g., soil organic carbon), Xβ contains specific surrogate data depending on the response variable, and ε is the remaining unexplained error. EPA tested models with a variety of surrogate data, including commodity statistics, weather data, or other relevant information. Parameters are estimated from the observed data for 1990 to 2012 using standard statistical techniques, and these estimates are used to predict the missing emissions data for 2013 to 2017.

A critical issue in using splicing methods in general, is to adequately account for the additional uncertainty introduced by predicting emissions with related information without compiling the full inventory. For example, predicting CH<sub>4</sub> emissions will increase the total variation in the emission estimates for these specific years,

<sup>8</sup> See <<https://quickstats.nass.usda.gov/>>.

compared to those years in which the full inventory is compiled. This added uncertainty is quantified within the model framework using a Monte Carlo approach. The approach requires estimating parameters for results in each Monte Carlo simulation for the full inventory (i.e., the surrogate data model is refit with the emissions estimated in each Monte Carlo iteration from the full inventory analysis with data from 1990 to 2012).

## Uncertainty and Time-Series Consistency

Sources of uncertainty in the Tier 3 method include management practices, uncertainties in model structure (i.e., algorithms and parameterization), and variance associated with the NRI sample. Sources of uncertainty in the IPCC (2006) Tier 1 method include the emission factors, management practices, and variance associated with the NRI sample. A Monte Carlo analysis was used to propagate uncertainties in the Tier 1 and 3 methods. For 2013 to 2017, there is additional uncertainty propagated through the Monte Carlo Analysis associated with the surrogate data method. (See Box 5-2 for information about propagating uncertainty with the surrogate data method.) The uncertainties from the Tier 1 and 3 approaches are combined to produce the final CH<sub>4</sub> emissions estimate using simple error propagation (IPCC 2006). Additional details on the uncertainty methods are provided in Annex 3.12. Rice cultivation CH<sub>4</sub> emissions in 2017 were estimated to be between 8.6 and 16.9 MMT CO<sub>2</sub> Eq. at a 95 percent confidence level, which indicates a range of 25 percent below to 49 percent above the actual 2017 emission estimate of 11.3 MMT CO<sub>2</sub> Eq. (see Table 5-14).

**Table 5-14: Approach 2 Quantitative Uncertainty Estimates for CH<sub>4</sub> Emissions from Rice Cultivation (MMT CO<sub>2</sub> Eq. and Percent)**

Source	Inventory Method	Gas	2017 Emission Estimate (MMT CO <sub>2</sub> Eq.)	Uncertainty Range Relative to Emission Estimate <sup>a</sup>			
				(MMT CO <sub>2</sub> Eq.)		(%)	
				Lower Bound	Upper Bound	Lower Bound	Upper Bound
Rice Cultivation	Tier 3	CH <sub>4</sub>	9.6	6.9	12.2	-27%	+27%
Rice Cultivation	Tier 1	CH <sub>4</sub>	1.8	0.8	2.8	-55%	+55%
<b>Rice Cultivation</b>	<b>Total</b>	<b>CH<sub>4</sub></b>	<b>11.3</b>	<b>8.6</b>	<b>16.9</b>	<b>-25%</b>	<b>+49%</b>

<sup>a</sup> Range of emission estimates predicted by Monte Carlo Stochastic Simulation for a 95 percent confidence interval.

## QA/QC and Verification

Quality control measures include checking input data, model scripts, and results to ensure data are properly handled throughout the inventory process. Inventory reporting forms and text are reviewed and revised as needed to correct transcription errors. No errors were found in the reporting forms and text.

Model results are compared to field measurements to verify if results adequately represent CH<sub>4</sub> emissions. The comparisons included over 15 long-term experiments, representing about 80 combinations of management treatments across all of the sites. A statistical relationship was developed to assess uncertainties in the model structure, adjusting the estimates for model bias and assessing precision in the resulting estimates (methods are described in Ogle et al. 2007). See Annex 3.12 for more information.

## Planned Improvements

New land representation data and rice cultivation data were not compiled for the current Inventory. A surrogate data method has been applied to estimate emissions in the latter part of the time series, which introduces additional uncertainty in the emissions data. Therefore, a key improvement for a future Inventory will be to update the time series for CH<sub>4</sub> emissions from rice cultivation by compiling the latest land use data and related management statistics.

In addition, a major improvement is underway to update the time series of management data with information from the USDA-NRCS Conservation Effects Assessment Program (CEAP). This improvement will fill several gaps in the management data including more specific data on fertilizer rates, updated tillage practices, water management,



organic amendments and more information on planting and harvesting dates. This improvement is expected to be completed for the 1990 through 2018 Inventory (i.e., 2020 submission). However, the timeline may be extended if there are insufficient resources to fund this improvement.

## 5.4 Agricultural Soil Management (CRF Source Category 3D)

---

Nitrous oxide is naturally produced in soils through the microbial processes of nitrification and denitrification that is driven by the availability of mineral nitrogen (N) (Firestone and Davidson 1989).<sup>9</sup> Mineral N is made available in soils through decomposition of soil organic matter and plant litter, as well as asymbiotic fixation of N from the atmosphere.<sup>10</sup> A number of agricultural activities increase mineral N availability in soils that lead to direct N<sub>2</sub>O emissions from nitrification and denitrification at the site of a management activity (see Figure 5-3) (Mosier et al. 1998), including N fertilization; application of managed livestock manure and other organic materials such as biosolids (i.e., sewage sludge); deposition of manure on soils by domesticated animals in pastures, rangelands, and paddocks (PRP) (i.e., by grazing animals and other animals whose manure is not managed); production of N-fixing crops and forages; retention of crop residues; and drainage of organic soils (i.e., soils with a high organic matter content, otherwise known as Histosols<sup>11</sup>) (IPCC 2006). Additionally, agricultural soil management activities, including irrigation, drainage, tillage practices, and fallowing of land, can influence N mineralization from soil organic matter and levels of asymbiotic N fixation by impacting moisture and temperature regimes in soils. Indirect emissions of N<sub>2</sub>O occur when N is transported from a site and is subsequently converted to N<sub>2</sub>O; there are two pathways for indirect emissions: (1) volatilization and subsequent atmospheric deposition of applied/mineralized N, and (2) surface runoff and leaching of applied/mineralized N into groundwater and surface water.<sup>12</sup> Direct and indirect emissions from agricultural lands are included in this section (i.e., cropland and grassland as defined in Section 6.1 Representation of the U.S. Land Base; N<sub>2</sub>O emissions from Forest Land and Settlements soils are found in Sections 6.2 and 6.10, respectively).

---


<sup>9</sup> Nitrification and denitrification are driven by the activity of microorganisms in soils. Nitrification is the aerobic microbial oxidation of ammonium (NH<sub>4</sub><sup>+</sup>) to nitrate (NO<sub>3</sub><sup>-</sup>), and denitrification is the anaerobic microbial reduction of nitrate to N<sub>2</sub>. Nitrous oxide is a gaseous intermediate product in the reaction sequence of denitrification, which leaks from microbial cells into the soil and then into the atmosphere. Nitrous oxide is also produced during nitrification, although by a less well-understood mechanism (Nevison 2000).

<sup>10</sup> Asymbiotic N fixation is the fixation of atmospheric N<sub>2</sub> by bacteria living in soils that do not have a direct relationship with plants.

<sup>11</sup> Drainage of organic soils in former wetlands enhances mineralization of N-rich organic matter, thereby increasing N<sub>2</sub>O emissions from these soils.

<sup>12</sup> These processes entail volatilization of applied or mineralized N as NH<sub>3</sub> and NO<sub>x</sub>, transformation of these gases within the atmosphere (or upon deposition), and deposition of the N primarily in the form of particulate NH<sub>4</sub><sup>+</sup>, nitric acid (HNO<sub>3</sub>), and NO<sub>x</sub>, in addition to leaching and runoff of NO<sub>3</sub><sup>-</sup> that is converted to N<sub>2</sub>O in aquatic systems.

**Figure 5-3: Sources and Pathways of N that Result in N<sub>2</sub>O Emissions from Agricultural Soil Management**



Agricultural soils produce the majority of N<sub>2</sub>O emissions in the United States. Estimated emissions from this source in 2017 are 266.4 MMT CO<sub>2</sub> Eq. (894 kt) (see Table 5-15 and Table 5-16). Annual N<sub>2</sub>O emissions from agricultural soils are 6 percent greater in the 2017 compared to 1990, but emissions fluctuated between 1990 and 2017 due to inter-annual variability largely associated with weather patterns, synthetic fertilizer use, and crop production. From 1990 to 2017, on average, cropland accounted for 70 percent of total direct emissions, while grassland accounted for 30 percent. On average, 81 percent of indirect emissions are from croplands and 19 percent from grasslands. Estimated direct and indirect N<sub>2</sub>O emissions by sub-source category are shown in Table 5-17 and Table 5-18.

**Table 5-15: N<sub>2</sub>O Emissions from Agricultural Soils (MMT CO<sub>2</sub> Eq.)**

Activity	1990	2005	2013	2014	2015	2016	2017
<b>Direct</b>	<b>212.7</b>	<b>218.9</b>	<b>225.9</b>	<b>223.4</b>	<b>239.0</b>	<b>228.8</b>	<b>227.7</b>
Cropland	148.1	154.7	161.3	160.1	166.8	162.2	161.6
Grassland	64.6	64.2	64.5	63.4	72.2	66.6	66.1
<b>Indirect</b>	<b>39.0</b>	<b>35.7</b>	<b>39.3</b>	<b>38.8</b>	<b>38.8</b>	<b>38.8</b>	<b>38.8</b>
Cropland	31.4	28.7	32.2	31.7	31.6	31.6	31.6
Grassland	7.6	7.0	7.2	7.1	7.1	7.1	7.2
<b>Total</b>	<b>251.7</b>	<b>254.5</b>	<b>265.2</b>	<b>262.3</b>	<b>277.8</b>	<b>267.6</b>	<b>266.4</b>

Notes: Estimates after 2012 are based on a data splicing method (See Methodology section). Totals may not sum due to independent rounding.

**Table 5-16: N<sub>2</sub>O Emissions from Agricultural Soils (kt)**

Activity	1990	2005	2013	2014	2015	2016	2017
<b>Direct</b>	<b>714</b>	<b>735</b>	<b>758</b>	<b>750</b>	<b>802</b>	<b>768</b>	<b>764</b>
Cropland	497	519	541	537	560	544	542
Grassland	217	216	217	213	242	224	222
<b>Indirect</b>	<b>131</b>	<b>120</b>	<b>132</b>	<b>130</b>	<b>130</b>	<b>130</b>	<b>130</b>
Cropland	105	96	108	106	106	106	106
Grassland	26	23	24	24	24	24	24
<b>Total</b>	<b>845</b>	<b>854</b>	<b>890</b>	<b>880</b>	<b>932</b>	<b>898</b>	<b>894</b>

Notes: Estimates after 2012 are based on a data splicing method (See Methodology section). Totals may not sum due to independent rounding.

**Table 5-17: Direct N<sub>2</sub>O Emissions from Agricultural Soils by Land Use Type and N Input Type (MMT CO<sub>2</sub> Eq.)**

Activity	1990	2005	2013	2014	2015	2016	2017
<b>Cropland</b>	<b>148.1</b>	<b>154.6</b>	<b>161.3</b>	<b>160.0</b>	<b>166.7</b>	<b>162.2</b>	<b>161.5</b>
<b>Mineral Soils</b>	<b>144.7</b>	<b>151.3</b>	<b>158.0</b>	<b>156.8</b>	<b>163.5</b>	<b>159.0</b>	<b>158.3</b>
Synthetic Fertilizer	53.3	54.3	60.0	59.6	62.2	60.5	60.2
Organic Amendment <sup>a</sup>	11.5	12.6	13.4	13.3	13.5	13.4	13.3
Residue N <sup>b</sup>	21.7	22.4	24.4	24.2	25.3	24.6	24.5
Mineralization and Asymbiotic Fixation	58.3	62.0	60.2	59.7	62.5	60.6	60.3
<b>Drained Organic Soils</b>	<b>3.3</b>	<b>3.3</b>	<b>3.3</b>	<b>3.2</b>	<b>3.2</b>	<b>3.2</b>	<b>3.2</b>
<b>Grassland</b>	<b>64.6</b>	<b>64.2</b>	<b>64.5</b>	<b>63.4</b>	<b>72.2</b>	<b>66.6</b>	<b>66.1</b>
<b>Mineral Soils</b>	<b>61.7</b>	<b>61.5</b>	<b>61.9</b>	<b>60.8</b>	<b>69.6</b>	<b>64.0</b>	<b>63.4</b>
Synthetic Fertilizer	0.9	0.8	1.0	1.0	1.1	1.0	1.0
PRP Manure	16.3	14.0	12.5	12.4	13.4	12.8	12.7
Managed Manure <sup>c</sup>	0.1	0.1	0.1	0.1	0.2	0.1	0.1
Biosolids (i.e., Sewage Sludge)	0.2	0.5	0.6	0.6	0.6	0.6	0.6
Residue N <sup>d</sup>	15.9	16.6	18.8	18.4	21.4	19.5	19.3
Mineralization and Asymbiotic Fixation	28.2	29.5	29.0	28.3	32.9	30.0	29.7
<b>Drained Organic Soils</b>	<b>2.9</b>	<b>2.7</b>	<b>2.6</b>	<b>2.6</b>	<b>2.6</b>	<b>2.6</b>	<b>2.6</b>
<b>Total</b>	<b>212.7</b>	<b>218.9</b>	<b>225.9</b>	<b>223.4</b>	<b>239.0</b>	<b>228.8</b>	<b>227.7</b>

<sup>a</sup> Organic amendment inputs include managed manure, daily spread manure, and commercial organic fertilizers (i.e., dried blood, dried manure, tankage, compost, and other).

<sup>b</sup> Cropland residue N inputs include N in unharvested legumes as well as crop residue N.

<sup>c</sup> Managed manure inputs include managed manure and daily spread manure amendments that are applied to grassland soils.

<sup>d</sup> Grassland residue N inputs include N in ungrazed legumes as well as ungrazed grass residue N.

Notes: Estimates after 2012 are based on a data splicing method (See Methodology section). Totals may not sum due to independent rounding.

**Table 5-18: Indirect N<sub>2</sub>O Emissions from Agricultural Soils (MMT CO<sub>2</sub> Eq.)**


Activity	1990	2005	2013	2014	2015	2016	2017
<b>Cropland</b>	<b>31.4</b>	<b>28.7</b>	<b>32.2</b>	<b>31.7</b>	<b>31.6</b>	<b>31.6</b>	<b>31.6</b>
Volatilization & Atm.							
Deposition	6.2	7.0	6.8	6.8	6.7	6.7	6.7
Surface Leaching & Run-Off	25.3	21.7	25.3	24.9	24.9	24.9	24.9
<b>Grassland</b>	<b>7.6</b>	<b>7.0</b>	<b>7.2</b>	<b>7.1</b>	<b>7.1</b>	<b>7.1</b>	<b>7.2</b>
Volatilization & Atm.							
Deposition	4.3	4.5	4.2	4.2	4.2	4.2	4.2
Surface Leaching & Run-Off	3.2	2.5	2.9	2.9	2.9	2.9	2.9
<b>Total</b>	<b>39.0</b>	<b>35.7</b>	<b>39.3</b>	<b>38.8</b>	<b>38.8</b>	<b>38.8</b>	<b>38.8</b>

Notes: Estimates after 2012 are based on a data splicing method (See Methodology section). Totals may not sum due to independent rounding.

Figure 5-4 and Figure 5-5 show regional patterns for direct N<sub>2</sub>O emissions. Figure 5-6 and Figure 5-7 show indirect N<sub>2</sub>O emissions from volatilization, and Figure 5-8 and Figure 5-9 show the indirect N<sub>2</sub>O emissions from leaching and runoff in croplands and grasslands, respectively. Annual emissions in 2012<sup>13</sup> are shown for the Tier 3 Approach only.

<sup>13</sup> Only national-scale emissions are estimated for 2013 to 2017 in the current Inventory using the splicing method, and therefore the fine-scale emission patterns in these maps are based on Inventory data from 2012.

**Figure 5-4: Crops, 2012 Annual Direct N<sub>2</sub>O Emissions Estimated Using the Tier 3 DAYCENT Model (MMT CO<sub>2</sub> Eq./year)**




Note: Only national-scale emissions are estimated for 2013 to 2017 using a splicing method, and therefore the fine-scale emission patterns in this map are based on Inventory data from 2012.

Direct N<sub>2</sub>O emissions from croplands occur throughout all of the cropland regions but tend to be high in the Midwestern Corn Belt Region (Illinois, Iowa, Indiana, Ohio, southern Minnesota and Wisconsin, and eastern Nebraska), where a large portion of the land is used for growing highly fertilized corn and N-fixing soybean crops (see Figure 5-4). Emissions are also high in the Lower Mississippi River Basin from Missouri to Louisiana, and highly productive irrigated areas, such as Platte River, which flows from Colorado through Nebraska, Snake River Valley in Idaho and the Central Valley in California. Direct emissions are low in many parts of the eastern United States because only a small portion of land is cultivated, and also in many western states where rainfall and access to irrigation water are limited.

Direct emissions from grasslands are highest in the southeast, particularly Kentucky and Tennessee, in addition to areas in east Texas and Iowa, where there tends to be higher rates of manure amendments on a relatively small amount of pasture, compared to other regions of the United States. However, total emissions from grasslands tend to be higher in the Great Plains and western United States (see Figure 5-5) where a high proportion of the land is dominated by grasslands and used for cattle grazing.




**Figure 5-5: Grasslands, 2012 Annual Direct N<sub>2</sub>O Emissions Estimated Using the Tier 3 DAYCENT Model (MMT CO<sub>2</sub> Eq./year)**



Note: Only national-scale emissions are estimated for 2013 to 2017 using a splicing method, and therefore the fine-scale emission patterns in this map are based on Inventory data from 2012.

Indirect N<sub>2</sub>O emissions from volatilization in croplands have a similar pattern as the direct N<sub>2</sub>O emissions with high emissions in the Midwestern Corn Belt and Lower Mississippi River Basin. Indirect N<sub>2</sub>O emissions from volatilization in grasslands are higher in the Southeastern United States than in other regions. The higher emissions in this region are mainly due to productive pastures that support intensive grazing, which in turn, stimulates NH<sub>3</sub> volatilization. Indirect N<sub>2</sub>O emissions from surface runoff and leaching of applied/mineralized N is highest in the Eastern United States for both croplands and grasslands. This region has greater precipitation and higher levels of leaching and runoff compared to arid to semi-arid regions in the Western United States.


**Figure 5-6: Crops, 2012 Annual Indirect N<sub>2</sub>O Emissions from Volatilization Using the Tier 3 DAYCENT Model (MMT CO<sub>2</sub> Eq./year)**



Note: Only national-scale emissions are estimated for 2013 to 2017 using a splicing method, and therefore the fine-scale emission patterns in this map are based on Inventory data from 2012.




**Figure 5-7: Grasslands, 2012 Annual Indirect N<sub>2</sub>O Emissions from Volatilization Using the Tier 3 DAYCENT Model (MMT CO<sub>2</sub> Eq./year)**




Note: Only national-scale emissions are estimated for 2013 to 2017 using a splicing method, and therefore the fine-scale emission patterns in this map are based on Inventory data from 2012.

**Figure 5-8: Crops, 2012 Annual Indirect N<sub>2</sub>O Emissions from Leaching and Runoff Using the Tier 3 DAYCENT Model (MMT CO<sub>2</sub> Eq./year)**



Note: Only national-scale emissions are estimated for 2013 to 2017 using a splicing method, and therefore the fine-scale emission patterns in this map are based on Inventory data from 2012.

**Figure 5-9: Grasslands, 2012 Annual Indirect N<sub>2</sub>O Emissions from Leaching and Runoff Using the Tier 3 DAYCENT Model (MMT CO<sub>2</sub> Eq./year)**



Note: Only national-scale emissions are estimated for 2013 to 2017 using a splicing method, and therefore the fine-scale emission patterns in this map are based on Inventory data from 2012.

## Methodology

The 2006 IPCC Guidelines (IPCC 2006) divide emissions from the agricultural soil management source category into five components, including (1) direct emissions from N additions to cropland and grassland mineral soils from synthetic fertilizers, biosolids (i.e., sewage sludge) applications, crop residues, organic amendments, and biological N fixation associated with planting of legumes on cropland and grassland soils; (2) direct emissions from soil organic matter mineralization due to land use and management change; (3) direct emissions from drainage of organic soils in croplands and grasslands; (4) direct emissions from soils due to manure deposited by livestock on PRP grasslands; and (5) indirect emissions from soils and water from N additions and manure deposition to soils that lead to volatilization, leaching, or runoff of N and subsequent conversion to N<sub>2</sub>O.

In this source category, the United States reports on all croplands, as well as all “managed” grasslands, whereby anthropogenic greenhouse gas emissions are estimated consistent with the managed land concept (IPCC 2006), including direct and indirect N<sub>2</sub>O emissions from asymbiotic fixation<sup>14</sup> and mineralization of soil organic matter and litter. One recommendation from IPCC (2006) that has not been completely adopted is the estimation of emissions from grassland pasture renewal, which involves occasional plowing to improve forage production in pastures. Currently no data are available to address pasture renewal.

<sup>14</sup> N inputs from asymbiotic N fixation are not directly addressed in 2006 IPCC Guidelines, but are a component of the total emissions from managed lands and are included in the Tier 3 approach developed for this source.

## Direct N<sub>2</sub>O Emissions

The methodology used to estimate direct N<sub>2</sub>O emissions from agricultural soil management in the United States is based on a combination of IPCC Tier 1 and 3 approaches, along with application of a splicing method for latter years in the Inventory time series (IPCC 2006; Del Grosso et al. 2010). A Tier 3 process-based model (DAYCENT) is used to estimate direct emissions from a variety of crops that are grown on mineral (i.e., non-organic) soils, as well as the direct emissions from non-federal grasslands with the exception of biosolids (i.e., sewage sludge) amendments (Del Grosso et al. 2010). The Tier 3 approach has been specifically designed and tested to estimate N<sub>2</sub>O emissions in the United States, accounting for more of the environmental and management influences on soil N<sub>2</sub>O emissions than the IPCC Tier 1 method (see Box 5-3 for further elaboration). Moreover, the Tier 3 approach allows for the Inventory to address direct N<sub>2</sub>O emissions and soil C stock changes from mineral cropland soils in a single analysis. Carbon and N dynamics are linked in plant-soil systems through biogeochemical processes of microbial decomposition and plant production (McGill and Cole 1981). Coupling the two source categories (i.e., agricultural soil C and N<sub>2</sub>O) in a single inventory analysis ensures that there is consistent activity data and treatment of the processes, and interactions are taken into account between C and N cycling in soils.

The Tier 3 approach is based on the cropping and land use histories recorded in the USDA National Resources Inventory (NRI) (USDA-NRCS 2015). The NRI is a statistically-based sample of all non-federal land,<sup>15</sup> and includes 363,286 points on agricultural land for the conterminous United States that are included in the Tier 3 method. The Tier 1 approach is used to estimate the emissions from the remaining 205,487 in the NRI survey that are designated as cropland or grassland (discussed later in this section). Each point is associated with an “expansion factor” that allows scaling of N<sub>2</sub>O emissions from NRI points to the entire country (i.e., each expansion factor represents the amount of area with the same land-use/management history as the sample point). Each NRI point was sampled on a 5-year cycle from 1982 until 1997. For cropland, data were collected in 4 out of 5 years in the cycle (i.e., 1979 through 1982, 1984 through 1987, 1989 through 1992, and 1994 through 1997). In 1998, the NRI program began collecting annual data, and the annual data are currently available through 2012 (USDA-NRCS 2015).

### Box 5-3: Tier 1 vs. Tier 3 Approach for Estimating N<sub>2</sub>O Emissions

The IPCC (2006) Tier 1 approach is based on multiplying activity data on different N inputs (i.e., synthetic fertilizer, manure, N fixation, etc.) by the appropriate default IPCC emission factors to estimate N<sub>2</sub>O emissions on an input-by-input basis. The Tier 1 approach requires a minimal amount of activity data, readily available in most countries (e.g., total N applied to crops); calculations are simple; and the methodology is highly transparent. In contrast, the Tier 3 approach developed for this Inventory employs a process-based model (i.e., DAYCENT) that represents the interaction of N inputs, land use and management, as well as environmental conditions at specific locations. Consequently, the Tier 3 approach produces more accurate estimates; it accounts more comprehensively for land-use and management impacts and their interaction with environmental factors (i.e., weather patterns and soil characteristics), which will enhance or dampen anthropogenic influences. However, the Tier 3 approach requires more detailed activity data (e.g., crop-specific N amendment rates), additional data inputs (i.e., daily weather, soil types, etc.), and considerable computational resources and programming expertise. The Tier 3 methodology is less transparent, and thus it is critical to evaluate the output of Tier 3 methods against measured data in order to demonstrate that the method is an improvement over lower tier methods for estimating emissions (IPCC 2006). Another important difference between the Tier 1 and Tier 3 approaches relates to assumptions regarding N cycling. Tier 1 assumes that N added to a system is subject to N<sub>2</sub>O emissions only during that year and cannot be stored in soils and contribute to N<sub>2</sub>O emissions in subsequent years. This is a simplifying assumption that is likely to create bias in estimated N<sub>2</sub>O emissions for a specific year. In contrast, the process-based model used in the Tier 3 approach includes the legacy effect of N added to soils in previous years that is re-mineralized from soil organic matter and emitted as N<sub>2</sub>O during subsequent years.

---

<sup>15</sup> The NRI survey does include sample points on federal lands, but the program does not collect data from those sample locations.



DAYCENT is used to estimate N<sub>2</sub>O emissions associated with production of alfalfa hay, barley, corn, cotton, dry beans, grass hay, grass-clover hay, lentils, oats, onions, peanuts, peas, potatoes, rice, sorghum, soybeans, sugar beets, sunflowers, tobacco, tomatoes, and wheat, but is not applied to estimate N<sub>2</sub>O emissions from other crops or rotations with other crops,<sup>16</sup> such as sugarcane, some vegetables, tobacco, and perennial/horticultural crops. Areas that are converted between agriculture (i.e., cropland and grassland) and other land uses, such as forest land, wetland and settlements, are not simulated with DAYCENT. DAYCENT is also not used to estimate emissions from land areas with very gravelly, cobbly, or shaley soils in the topsoil (greater than 35 percent by volume in the top 30 cm of the soil profile), or to estimate emissions from drained organic soils (Histosols). The Tier 3 method has not been fully tested for estimating N<sub>2</sub>O emissions associated with these crops and rotations, land uses, as well as organic soils or cobbly, gravelly, and shaley mineral soils. In addition, federal grassland areas are not simulated with DAYCENT due to limited activity data on land use histories. For areas that are not included in the DAYCENT simulations, the Tier 1 IPCC (2006) methodology is used to estimate (1) direct emissions from crops on mineral soils that are not simulated by DAYCENT; (2) direct emissions from PRP on federal grasslands; and (3) direct emissions from drained organic soils in croplands and grasslands.

A splicing method is used to estimate soil N<sub>2</sub>O emissions from 2013 to 2017 at the national scale because new NRI activity data are not available for those years. Specifically, linear regression models with autoregressive moving-average (ARMA) errors (Brockwell and Davis 2016) are used to estimate the relationship between surrogate data and the 1990 to 2012 emissions that are derived using the Tier 3 method. Surrogate data for these regression models include corn and soybean yields from USDA-NASS statistics,<sup>17</sup> and weather data from the PRISM Climate Group (PRISM 2015). For the Tier 1 method, a linear-time series model is used to estimate emissions from 2013 to 2017 without surrogate data. See Box 5-4 for more information about the splicing method. Emission estimates for 2013 to 2017 will be recalculated in future Inventory reports when new NRI data are available.

#### Box 5-4: Surrogate Data Method

An approach to extend the time series is needed for Agricultural Soil Management because the Inventory is only fully re-compiled every two years for many categories in the AFOLU sector as part of the biennial update reporting process, and even in years that the Inventory is re-compiled fully with the Tier 1 and 3 methods, there are typically gaps at the end of the time series. This is mainly because the National Resources Inventory (NRI), which provides critical information for estimating greenhouse gas emissions and removals, does not release data every year.

Splicing methods have been used to impute missing data at the end of the emission time series for both the Tier 1 and 3 methods. Specifically, a linear regression model with autoregressive moving-average (ARMA) errors (Brockwell and Davis 2016) is used to estimate emissions based on the modeled 1990 to 2012 emissions data, which has been compiled using the inventory methods described in this section. The model to extend the time series is given by

$$Y = X\beta + \varepsilon,$$

where Y is the response variable (e.g., soil organic carbon), Xβ for the Tier 3 data contains specific surrogate data depending on the response variable, and ε is the remaining unexplained error. Models with a variety of surrogate data were tested, including commodity statistics, weather data, or other relevant information. Xβ for the Tier 1 data only contains year as a predictor of emission patterns over the time series, and therefore, is a linear time series model with no surrogate data. Parameters are estimated from the emissions data for 1990 to 2012 using standard statistical techniques, and these estimates are used in the model described above to predict the missing emissions data for 2013 to 2017.

A critical issue when applying splicing methods is to account for the additional uncertainty introduced by predicting emissions with related information without compiling the full inventory. Specifically, uncertainty will increase for years with imputed estimates based on the splicing methods, compared to those years in which the full inventory is compiled. This additional uncertainty is quantified within the model framework using a Monte Carlo approach. Consequently, the uncertainty from the original inventory data, which is produced with the Tier 1 and 3 methods, is

<sup>16</sup> A small proportion of the major commodity crop production, such as corn and wheat, is included in the Tier 1 analysis because these crops are rotated with other crops or land uses (e.g., forest lands) that are not simulated by DAYCENT.

<sup>17</sup> See <<https://quickstats.nass.usda.gov/>>.

combined with the uncertainty in the parameters from the data splicing model. The approach requires estimating parameters for results in each Monte Carlo simulation for the full inventory (i.e., the surrogate data model is refit with the emissions estimated in each Monte Carlo iteration from the full inventory analysis with data from 1990 to 2012). Therefore, the data splicing method generates emissions estimates from each surrogate data model, which are used to derive confidence intervals in the estimates for the missing emissions data from 2013 to 2017.

---

### *Tier 3 Approach for Mineral Cropland Soils*

The DAYCENT biogeochemical model (Parton et al. 1998; Del Grosso et al. 2001 and 2011) is used to estimate direct N<sub>2</sub>O emissions from mineral cropland soils that are managed for production of a wide variety of crops (see list in previous paragraph) based on the cropping histories in the 2012 NRI (USDA-NRCS 2015). Crops simulated by DAYCENT are grown on approximately 91 percent of total cropland area in the United States. For agricultural systems in the central region of the United States, crop production for key crops (i.e., corn, soybeans, sorghum, cotton, and wheat) is simulated in DAYCENT with a NASA-CASA production algorithm (Potter et al. 1993; Potter et al. 2007) using the Moderate Resolution Imaging Spectroradiometer (MODIS) Enhanced Vegetation Index (EVI) products, MOD13Q1 and MYD13Q1, with a pixel resolution of 250m.<sup>18</sup>

DAYCENT is used to estimate direct N<sub>2</sub>O emissions due to mineral N available from the following sources: (1) the application of synthetic fertilizers; (2) the application of livestock manure; (3) the retention of crop residues and subsequent mineralization of N during microbial decomposition (i.e., leaving residues in the field after harvest instead of burning or collecting residues); (4) mineralization of soil organic matter; and (5) asymbiotic fixation. Note that commercial organic fertilizers (TVA 1991 through 1994; AAPFCO 1995 through 2015) are addressed with the Tier 1 method because county-level application data would be needed to simulate applications in DAYCENT, and currently data are only available at the national scale. The third and fourth sources are generated internally by the DAYCENT model.

Synthetic fertilizer data are based on fertilizer use and rates by crop type for different regions of the United States, and are obtained primarily from the USDA Economic Research Service. The data collection program was known as the Cropping Practices Surveys through 1995 (USDA-ERS 1997), and then became the Agricultural Resource Management Surveys (ARMS) (USDA-ERS 2015). Additional data are compiled through other sources particularly the National Agricultural Statistics Service (NASS 1992, 1999, 2004). Frequency and rates of livestock manure application to cropland during 1997 are estimated from data compiled by the USDA Natural Resources Conservation Service (Edmonds et al. 2003), and then adjusted using county-level estimates of manure available for application in other years. The adjustments are based on county-scale ratios of manure available for application to soils in other years relative to 1997 (see Annex 3.12 for further details). Greater availability of managed manure N relative to 1997 is assumed to increase the area amended with manure, while reduced availability of manure N relative to 1997 is assumed to reduce the amended area. Data on the county-level N available for application is estimated for managed manure systems based on the total amount of N excreted in manure minus N losses during storage and transport, and including the addition of N from bedding materials. Nitrogen losses include direct N<sub>2</sub>O emissions, volatilization of ammonia and NO<sub>x</sub>, runoff and leaching, and poultry manure used as a feed supplement. For unmanaged manure systems, it is assumed that no N losses or additions occur prior to the application of manure to the soil. More information on livestock manure production is available in Section 5.2 Manure Management and Annex 3.11.

The IPCC approach considers crop residue N and N mineralized from soil organic matter as activity data. However, they are not treated as activity data in DAYCENT simulations because residue production, symbiotic N fixation (e.g., legumes), mineralization of N from soil organic matter, and asymbiotic N fixation are internally generated by the model as part of the simulation. In other words, DAYCENT accounts for the influence of symbiotic N fixation, mineralization of N from soil organic matter and crop residue retained in the field, and asymbiotic N fixation on N<sub>2</sub>O emissions, but these are not model inputs. The N<sub>2</sub>O emissions from crop residues are reduced by approximately 3 percent (the assumed average burned portion for crop residues in the United States) to avoid double-counting associated with non-CO<sub>2</sub> greenhouse gas emissions from agricultural residue burning. The estimate of residue

---

<sup>18</sup> See <[https://lpdaac.usgs.gov/products/modis\\_products\\_table](https://lpdaac.usgs.gov/products/modis_products_table)>.

burning is based on state inventory data (ILENR 1993; Oregon Department of Energy 1995; Noller 1996; Wisconsin Department of Natural Resources 1993; Cibrowski 1996).

Additional sources of data are used to supplement the mineral N (USDA-ERS 1997, 2011), livestock manure (Edmonds et al. 2003), and land-use information (USDA-NRCS 2015). The Conservation Technology Information Center (CTIC 2004) provides annual data on tillage activity with adjustments for long-term adoption of no-till agriculture (Towery 2001). Tillage has an influence on soil organic matter decomposition and subsequent soil N<sub>2</sub>O emissions. The time series of tillage data from CTIC began in 1989 and ended in 2004, so further changes in tillage practices since 2004 are not currently captured in the Inventory and practices used in 2004 are assumed to apply for subsequent years. Daily weather data are used as an input in the model simulations, based on gridded weather data at a 4 km scale from the PRISM Climate Group (PRISM 2015). Soil attributes are obtained from the Soil Survey Geographic Database (SSURGO) (Soil Survey Staff 2011).

Each NRI point is run 100 times as part of the uncertainty assessment, yielding a total of over 18 million simulations for the analysis. Soil N<sub>2</sub>O emission estimates from DAYCENT are adjusted using a structural uncertainty estimator to account for uncertainty in model algorithms and parameter values (Del Grosso et al. 2010). Soil N<sub>2</sub>O emissions and associated 95 percent confidence intervals are estimated for each year between 1990 and 2012, but emissions from 2013 to 2017 are estimated using a splicing method that accounts for uncertainty in the original inventory data and the splicing method (See Box 5-4). Annual data are currently available through 2012 (USDA-NRCS 2015), and the Inventory time series will be updated in the future when new NRI data are released.

Nitrous oxide emissions from managed agricultural lands are the result of interactions among anthropogenic activities (e.g., N fertilization, manure application, tillage) and other driving variables, such as weather and soil characteristics. These factors influence key processes associated with N dynamics in the soil profile, including immobilization of N by soil microbial organisms, decomposition of organic matter, plant uptake, leaching, runoff, and volatilization, as well as the processes leading to N<sub>2</sub>O production (nitrification and denitrification). It is not possible to partition N<sub>2</sub>O emissions into each anthropogenic activity directly from model outputs due to the complexity of the interactions (e.g., N<sub>2</sub>O emissions from synthetic fertilizer applications cannot be distinguished from those resulting from manure applications). To approximate emissions by activity, the amount of mineral N added to the soil, or made available through decomposition of soil organic matter and plant litter, as well as asymbiotic fixation of N from the atmosphere, is determined for each N source and then divided by the total amount of mineral N in the soil according to the DAYCENT model simulation. The percentages are then multiplied by the total of direct N<sub>2</sub>O emissions in order to approximate the portion attributed to N management practices. This approach is only an approximation because it assumes that all N made available in soil has an equal probability of being released as N<sub>2</sub>O, regardless of its source, which is unlikely to be the case (Delgado et al. 2009). However, this approach allows for further disaggregation of emissions by source of N, which is valuable for reporting purposes and is analogous to the reporting associated with the IPCC (2006) Tier 1 method, in that it associates portions of the total soil N<sub>2</sub>O emissions with individual sources of N.

### *Tier 1 Approach for Mineral Cropland Soils*

The IPCC (2006) Tier 1 methodology is used to estimate direct N<sub>2</sub>O emissions for mineral cropland soils that are not simulated by DAYCENT (e.g., DAYCENT has not been parametrized to simulate all crop types and some soil types such as *Histosols*). For the Tier 1 Approach, estimates of direct N<sub>2</sub>O emissions from N applications are based on mineral soil N that is made available from the following practices: (1) the application of synthetic commercial fertilizers; (2) application of managed manure and non-manure commercial organic fertilizers; and (3) decomposition and mineralization of nitrogen from above- and below-ground crop residues in agricultural fields (i.e., crop biomass that is not harvested). Non-manure commercial organic amendments are only included in the Tier 1 analysis because these data are not available at the county-level, which is necessary for the DAYCENT simulations.<sup>19</sup> Consequently, all commercial organic fertilizer, as well as manure that is not added to crops in the DAYCENT simulations, are included in the Tier 1 analysis. The following sources are used to derive activity data:

---

<sup>19</sup> Commercial organic fertilizers include dried blood, tankage, compost, and other, but the dried manure and biosolids (i.e., sewage sludge) is removed from the dataset in order to avoid double counting with other datasets that are used for manure N and biosolids.



- A process-of-elimination approach is used to estimate synthetic N fertilizer additions for crop areas not simulated by DAYCENT. The total amount of fertilizer used on farms has been estimated at the county-level by the USGS from sales records (Ruddy et al. 2006), and these data are aggregated to obtain state-level N additions to farms. For 2002 through 2012, state-level fertilizer for on-farm use is adjusted based on annual fluctuations in total U.S. fertilizer sales (AAPFCO 1995 through 2007, 2008 through 2012). After subtracting the portion of fertilizer applied to crops and grasslands simulated by DAYCENT (see Tier 3 Approach for Mineral Cropland Soils and Direct N<sub>2</sub>O Emissions from Grassland Soils sections for information on data sources), the remainder of the total fertilizer used on farms is assumed to be applied to crops that are not simulated by DAYCENT.
- Similarly, a process-of-elimination approach is used to estimate manure N additions for crops that are not simulated by DAYCENT. The amount of manure N applied in the Tier 3 approach to crops and grasslands is subtracted from total manure N available for land application (see Tier 3 Approach for Mineral Cropland Soils and Direct N<sub>2</sub>O Emissions from Grassland Soils sections for information on data sources), and this difference is assumed to be applied to crops that are not simulated by DAYCENT.
- Commercial organic fertilizer additions are based on organic fertilizer consumption statistics, which are converted to units of N using average organic fertilizer N content (TVA 1991 through 1994; AAPFCO 1995 through 2012). Commercial fertilizers do include some manure and biosolids (i.e., sewage sludge), but the amounts are removed from the commercial fertilizer data to avoid double counting with the manure N dataset described above and the biosolids (i.e., sewage sludge) amendment data discussed later in this section.
- Crop residue N is derived by combining amounts of above- and below-ground biomass, which are determined based on NRI crop area data (USDA-NRCS 2013), crop production yield statistics (USDA-NASS 2018), dry matter fractions (IPCC 2006), linear equations to estimate above-ground biomass given dry matter crop yields from harvest (IPCC 2006), ratios of below-to-above-ground biomass (IPCC 2006), and N contents of the residues (IPCC 2006). N inputs from residue were reduced by 3 percent to account for average residue burning portions in the United States.

The total increase in soil mineral N from applied fertilizers and crop residues is multiplied by the IPCC (2006) default emission factor to derive an estimate of direct N<sub>2</sub>O emissions using the Tier 1 Approach.

Tier 1 soil N<sub>2</sub>O emissions from 2013 to 2017 are estimated using a splicing method that is described in Box 5-4, with the exception of commercial fertilizer additions, which are estimated with a splicing method from 2015 to 2017. As with the Tier 3 method, the time series will be recalculated in future Inventory reports (see Planned Improvements section).

### *Tier 1 Approach for Drainage of Organic Soils in Croplands and Grasslands*

The IPCC (2006) Tier 1 methods are used to estimate direct N<sub>2</sub>O emissions due to drainage of organic soils in croplands and grasslands at a state scale. State-scale estimates of the total area of drained organic soils are obtained from the 2012 NRI (USDA-NRCS 2015) using soils data from the Soil Survey Geographic Database (SSURGO) (Soil Survey Staff 2011). Temperature data from Daly et al. (1994 and 1998) are used to subdivide areas into temperate and tropical climates using the climate classification from IPCC (2006). To estimate annual emissions, the total temperate area is multiplied by the IPCC default emission factor for temperate regions, and the total tropical area is multiplied by the IPCC default emission factor for tropical regions (IPCC 2006). Annual NRI data are only available between 1990 and 2012. Consequently, emissions from 2013 to 2017 are estimated using a linear time series model (see Box 5-4). Estimates for 2013 to 2017 will be recalculated in future Inventory reports when new NRI data are available.

### *Tier 1 and 3 Approaches for Direct N<sub>2</sub>O Emissions from Grassland Soils*

As with N<sub>2</sub>O from croplands, the Tier 3 process-based DAYCENT model and Tier 1 method described in IPCC (2006) are combined to estimate emissions from non-federal grasslands and PRP manure N additions for federal grasslands, respectively. Grassland includes pasture and rangeland that produce grass forage primarily for livestock grazing. Rangelands are typically extensive areas of native grassland that are not intensively managed, while pastures are typically seeded grassland (possibly following tree removal) that may also have additional management, such as irrigation, fertilization, or interseeding legumes. DAYCENT is used to simulate N<sub>2</sub>O emissions from NRI

survey locations (USDA-NRCS 2015) on non-federal grasslands resulting from manure deposited by livestock directly onto pastures and rangelands (i.e., PRP manure), N fixation from legume seeding, managed manure amendments (i.e., manure other than PRP manure such as Daily Spread), and synthetic fertilizer application. Other N inputs are simulated within the DAYCENT framework, including N input from mineralization due to decomposition of soil organic matter and N inputs from senesced grass litter, as well as asymbiotic fixation of N from the atmosphere. The simulations used the same weather, soil, and synthetic N fertilizer data as discussed under the Tier 3 Approach in the Mineral Cropland Soils section. Mineral N fertilization rates are based on Carbon Sequestration Rural Appraisals (CSRA) conducted by the USDA-NRCS (USDA-NRCS, unpublished data). The CSRA was a solicitation of expert knowledge from USDA-NRCS staff throughout the United States to support the Inventory. Managed manure N amendments to grasslands are estimated from Edmonds et al. (2003) and adjusted for annual variation using data on the availability of managed manure N for application to soils, according to methods described in the Manure Management section (Section 5.2) and Annex 3.11. Biological N fixation is simulated within DAYCENT, and therefore is not an input to the model.

Manure N deposition from grazing animals in PRP systems (i.e., PRP manure) is another key input of N to grasslands. The amounts of PRP manure N applied on non-federal grasslands for each NRI point are based on amount of N excreted by livestock in PRP systems. The total amount of N excreted in each county is divided by the grassland area to estimate the N input rate associated with PRP manure. The resulting input rates are used in the DAYCENT simulations. DAYCENT simulations of non-federal grasslands accounted for approximately 78 percent of total PRP manure N in aggregate across the country. The remainder of the PRP manure N in each state is assumed to be excreted on federal grasslands, and the N<sub>2</sub>O emissions are estimated using the IPCC (2006) Tier 1 method with IPCC default emission factors.

Biosolids (i.e., sewage sludge) are assumed to be applied on grasslands because of the heavy metal content and other pollutants in human waste that limit its use as an amendment to croplands. Biosolids application is estimated from data compiled by EPA (1993, 1999, 2003), McFarland (2001), and NEBRA (2007) (see Section 7.2 Wastewater Treatment for a detailed discussion of the methodology for estimating sewage sludge available for land application). Biosolids soil amendments are only available at the national scale, and it is not possible to associate application with specific soil conditions and weather at NRI survey locations. Therefore, DAYCENT could not be used to simulate the influence of biosolids amendments on N<sub>2</sub>O emissions from grassland soils, and consequently, emissions from biosolids are estimated using the IPCC (2006) Tier 1 method.

As previously mentioned, each NRI point is simulated 100 times as part of the uncertainty assessment, yielding a total of over 18 million simulation runs for the analysis. Soil N<sub>2</sub>O emission estimates from DAYCENT are adjusted using a structural uncertainty estimator accounting for uncertainty in model algorithms and parameter values (Del Grosso et al. 2010). N<sub>2</sub>O emissions for the PRP manure N deposited on federal grasslands and applied biosolids N are estimated using the Tier 1 method by multiplying the N input by the default emission factor. Emissions from manure N are estimated at the state level and aggregated to the entire country, but emissions from biosolids N are calculated exclusively at the national scale.

Soil N<sub>2</sub>O emissions and 95 percent confidence intervals are estimated for each year between 1990 and 2012 based on the Tier 1 and 3 methods, with the exception of biosolids (discussed below), and emissions from 2013 to 2017 are estimated using a splicing method as described in Box 5-4. As with croplands, estimates for 2013 to 2017 will be recalculated in future inventories when new NRI data are available. Biosolids application data are compiled through 2017 in this Inventory, and therefore soil N<sub>2</sub>O emissions and confidence intervals are estimated using the Tier 1 method for all years in the time series without application of the splicing method.

## **Total Direct N<sub>2</sub>O Emissions from Cropland and Grassland Soils**

Annual direct emissions from the Tier 1 and 3 approaches for mineral and drained organic soils occurring in both croplands and grasslands are summed to obtain the total direct N<sub>2</sub>O emissions from agricultural soil management (see Table 5-15 and Table 5-16).

## **Indirect N<sub>2</sub>O Emissions**

This section describes the methods used for estimating indirect soil N<sub>2</sub>O emissions from croplands and grasslands. Indirect N<sub>2</sub>O emissions occur when mineral N made available through anthropogenic activity is transported from the soil either in gaseous or aqueous forms and later converted into N<sub>2</sub>O. There are two pathways leading to indirect

emissions. The first pathway results from volatilization of N as  $\text{NO}_x$  and  $\text{NH}_3$  following application of synthetic fertilizer, organic amendments (e.g., manure, biosolids), and deposition of PRP manure. Nitrogen made available from mineralization of soil organic matter and residue, including N incorporated into crops and forage from symbiotic N fixation, and input of N from asymbiotic fixation also contributes to volatilized N emissions. Volatilized N can be returned to soils through atmospheric deposition, and a portion of the deposited N is emitted to the atmosphere as  $\text{N}_2\text{O}$ . The second pathway occurs via leaching and runoff of soil N (primarily in the form of  $\text{NO}_3^-$ ) that is made available through anthropogenic activity on managed lands, mineralization of soil organic matter and residue, including N incorporated into crops and forage from symbiotic N fixation, and inputs of N into the soil from asymbiotic fixation. The  $\text{NO}_3^-$  is subject to denitrification in water bodies, which leads to  $\text{N}_2\text{O}$  emissions. Regardless of the eventual location of the indirect  $\text{N}_2\text{O}$  emissions, the emissions are assigned to the original source of the N for reporting purposes, which here includes croplands and grasslands.

### *Tier 1 and 3 Approaches for Indirect $\text{N}_2\text{O}$ Emissions from Atmospheric Deposition of Volatilized N*

The Tier 3 DAYCENT model and IPCC (2006) Tier 1 methods are combined to estimate the amount of N that is volatilized and eventually emitted as  $\text{N}_2\text{O}$ . DAYCENT is used to estimate N volatilization for land areas whose direct emissions are simulated with DAYCENT (i.e., most commodity and some specialty crops and most grasslands). The N inputs included are the same as described for direct  $\text{N}_2\text{O}$  emissions in the Tier 3 Approach for Mineral Cropland Soils and Direct  $\text{N}_2\text{O}$  Emissions from Grassland Soils sections. Nitrogen volatilization from all other areas is estimated using the Tier 1 method and default IPCC fractions for N subject to volatilization (i.e., N inputs on croplands not simulated by DAYCENT, PRP manure N excreted on federal grasslands, biosolids [i.e., sewage sludge] application on grasslands). For the volatilization data generated from both the DAYCENT and Tier 1 approaches, the IPCC (2006) default emission factor is used to estimate indirect  $\text{N}_2\text{O}$  emissions occurring due to re-deposition of the volatilized N (see Table 5-18).

### *Tier 1 and 3 Approaches for Indirect $\text{N}_2\text{O}$ Emissions from Leaching/Runoff*

As with the calculations of indirect emissions from volatilized N, the Tier 3 DAYCENT model and IPCC (2006) Tier 1 method are combined to estimate the amount of N that is subject to leaching and surface runoff into water bodies, and eventually emitted as  $\text{N}_2\text{O}$ . DAYCENT is used to simulate the amount of N transported from lands in the Tier 3 Approach. Nitrogen transport from all other areas is estimated using the Tier 1 method and the IPCC (2006) default factor for the proportion of N subject to leaching and runoff. This N transport estimate includes N applications on croplands that are not simulated by DAYCENT, biosolids amendments on grasslands, and PRP manure N excreted on federal grasslands. For both the DAYCENT Tier 3 and IPCC (2006) Tier 1 methods, nitrate leaching is assumed to be an insignificant source of indirect  $\text{N}_2\text{O}$  in cropland and grassland systems in arid regions, as discussed in IPCC (2006). In the United States, the threshold for significant nitrate leaching is based on the potential evapotranspiration (PET) and rainfall amount, similar to IPCC (2006), and is assumed to be negligible in regions where the amount of precipitation plus irrigation does not exceed 80 percent of PET. For leaching and runoff data estimated by the Tier 3 and Tier 1 approaches, the IPCC (2006) default emission factor is used to estimate indirect  $\text{N}_2\text{O}$  emissions that occur in groundwater and waterways (see Table 5-18).

Indirect soil  $\text{N}_2\text{O}$  emissions from 2013 to 2017 are estimated using the splicing method that is described in Box 5-4. As with the direct  $\text{N}_2\text{O}$  emissions, the time series will be recalculated in a future Inventory report when new activity data are compiled (see Planned Improvements section).

## **Uncertainty and Time-Series Consistency**

Uncertainty is estimated for each of the following five components of  $\text{N}_2\text{O}$  emissions from agricultural soil management: (1) direct emissions simulated by DAYCENT; (2) the components of indirect emissions (N volatilized and leached or runoff) simulated by DAYCENT; (3) direct emissions calculated with the IPCC (2006) Tier 1 method; (4) the components of indirect emissions (N volatilized and leached or runoff) calculated with the IPCC (2006) Tier 1 method; and (5) indirect emissions estimated with the IPCC (2006) Tier 1 method. Uncertainty in direct emissions, which account for the majority of  $\text{N}_2\text{O}$  emissions from agricultural management, as well as the components of indirect emissions calculated by DAYCENT are estimated with a Monte Carlo Analysis, addressing uncertainties in model inputs and structure (i.e., algorithms and parameterization) (Del Grosso et al. 2010). For 2013

to 2017, there is additional uncertainty propagated through the Monte Carlo Analysis associated with the splicing method (See Box 5-4).

Simple error propagation methods (IPCC 2006) are used to estimate confidence intervals for direct emissions calculated with the IPCC (2006) Tier 1 method, the proportion of volatilization and leaching or runoff estimated with the IPCC (2006) Tier 1 method, and indirect N<sub>2</sub>O emissions. Uncertainty in the splicing method is also included in the error propagation for 2013 to 2017 (see Box 5-4). Additional details on the uncertainty methods are provided in Annex 3.12. Table 5-19 shows the combined uncertainty for direct soil N<sub>2</sub>O emissions ranged from 17 percent below to 18 percent above the 2017 emission estimate of 227.7 MMT CO<sub>2</sub> Eq., and the combined uncertainty for indirect soil N<sub>2</sub>O emissions range from 58 percent below to 143 percent above the 2017 estimate of 38.8 MMT CO<sub>2</sub> Eq.

**Table 5-19: Quantitative Uncertainty Estimates of N<sub>2</sub>O Emissions from Agricultural Soil Management in 2017 (MMT CO<sub>2</sub> Eq. and Percent)**

Source	Gas	2017 Emission Estimate (MMT CO <sub>2</sub> Eq.)	Uncertainty Range Relative to Emission Estimate (%)			
			Lower Bound	Upper Bound	Lower Bound	Upper Bound
Direct Soil N <sub>2</sub> O Emissions	N <sub>2</sub> O	227.7	185.7	265.0	-17%	18%
Indirect Soil N <sub>2</sub> O Emissions	N <sub>2</sub> O	38.8	16.3	94.9	-58%	143%

Notes: Due to lack of data, uncertainties in managed manure N production, PRP manure N production, other organic fertilizer amendments, and biosolids (i.e., sewage sludge) amendments to soils are currently treated as certain; these sources of uncertainty will be included in future Inventory reports.


Additional uncertainty is associated with an incomplete estimation of N<sub>2</sub>O emissions from managed croplands and grasslands in Hawaii and Alaska. The Inventory currently includes the N<sub>2</sub>O emissions from mineral fertilizer and PRP N additions in Alaska and Hawaii, and drained organic soils in Hawaii. Land areas used for agriculture in Alaska and Hawaii are small relative to major commodity cropping states in the conterminous United States, so the emissions are likely to be small for the other sources of N (e.g., crop residue inputs), which are not currently included in the Inventory.

Methodological recalculations were applied to the entire time series to ensure time-series consistency from 1990 through 2017. Details on the emission trends through time are described in more detail in the Methodology section.

## QA/QC and Verification

DAYCENT results for N<sub>2</sub>O emissions and NO<sub>3</sub><sup>-</sup> leaching are compared with field data representing various cropland and grassland systems, soil types, and climate patterns (Del Grosso et al. 2005; Del Grosso et al. 2008), and further evaluated by comparing the model results to emission estimates produced using the IPCC (2006) Tier 1 method for the same sites. Nitrous oxide measurement data are available for 41 sites, which mostly occur in the United States, with five in Europe and three in Australia, representing over 200 different combinations of fertilizer treatments and cultivation practices. Nitrate leaching data are available for four sites in the United States, representing 10 different combinations of fertilizer amendments/tillage practices. DAYCENT estimates of N<sub>2</sub>O emissions are closer to measured values at most sites compared to the IPCC Tier 1 estimate (see Figure 5-10). In general, the IPCC Tier 1 methodology tends to over-estimate emissions when observed values are low and under-estimate emissions when observed values are high, while DAYCENT estimates have less bias. DAYCENT accounts for key site-level factors (i.e., weather, soil characteristics, and management) that are not addressed in the IPCC Tier 1 method, and thus the model is better able to represent the variability in N<sub>2</sub>O emissions. DAYCENT does have a tendency to under-estimate very high N<sub>2</sub>O emission rates; and estimates are adjusted using the statistical model derived from the comparison of model estimates to measurements (see Annex 3.12 for more information). Regardless, the comparison demonstrates that DAYCENT provides relatively high predictive capability for N<sub>2</sub>O emissions, and is an improvement over the IPCC Tier 1 method.

**Figure 5-10: Comparison of Measured Emissions at Field Sites and Modeled Emissions Using the DAYCENT Simulation Model and IPCC Tier 1 Approach (kg N<sub>2</sub>O per ha per year)**



Spreadsheets containing input data and probability distribution functions required for DAYCENT simulations of croplands and grasslands and unit conversion factors have been checked, in addition to the program scripts that are used to run the Monte Carlo uncertainty analysis. Links between spreadsheets have also been checked, updated, and corrected when necessary. Spreadsheets containing input data, emission factors, and calculations required for the Tier 1 method have been checked and updated as needed.

## Recalculations Discussion

Methodological recalculations in the current Inventory are associated with the following updates to the time series: (1) manure N deposition from grazing animals in PRP systems; (2) managed livestock manure N application data; and (3) biosolid soil amendment data. These changes resulted in an average decrease in emissions of 0.4 percent from 1990 to 2016 relative to the previous Inventory.

## Planned Improvements

New land representation data have not been compiled for this Inventory, and a splicing method has been applied to estimate emissions in the latter part of the time series, which introduces additional uncertainty in the emissions data. Therefore, a key improvement for a future Inventory will be to recalculate the time series from 2013 to 2017 with the latest land use data from the National Resources Inventory and related management statistics, particularly data compiled through the USDA-NRCS Conservation Effects Assessment Program (CEAP). CEAP data will be used to update the times series and fill several gaps in the management data, including more specific data on fertilizer rates, updated tillage practices, and more information on planting and harvesting dates for crops.

Several planned improvements are underway associated with improving the DAYCENT biogeochemical model. These improvements include a better representation of plant phenology, particularly senescence events following grain filling in crops. In addition, crop parameters associated with temperature and water stress effects on plant production will be further improved in DAYCENT with additional model calibration. Model development is underway to represent the influence of nitrification inhibitors and slow-release fertilizers (e.g., polymer-coated fertilizers) on N<sub>2</sub>O emissions. An improved representation of drainage as well as freeze-thaw cycles are also under

development. Experimental study sites will continue to be added for quantifying model structural uncertainty. Studies that have continuous (daily) measurements of N<sub>2</sub>O (e.g., Scheer et al. 2013) will be given priority.

Improvements are underway to simulate crop residue burning in the DAYCENT model based on the amount of crop residues burned according to the data that is used in the Field Burning of Agricultural Residues source category (see Section 5.7). Alaska and Hawaii are not included for all sources in the current Inventory for agricultural soil management, with the exception of N<sub>2</sub>O emissions from drained organic soils in croplands and grasslands for Hawaii, synthetic fertilizer and PRP N amendments for grasslands in Alaska and Hawaii. A planned improvement to add the remaining sources for these states into the Inventory analysis. There is also an improvement based on updating the Tier 1 emission factor for N<sub>2</sub>O emissions from drained organic soils by using the revised factor in the 2013 Supplement to the *2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands* (IPCC 2013).

These improvements are expected to be completed for the next Inventory (i.e., 2020 submission to the UNFCCC, 1990 through 2018 Inventory). However, the time line may be extended if there are insufficient resources to fund all or part of these planned improvements.

## 5.5 Liming (CRF Source Category 3G)

Crushed limestone (CaCO<sub>3</sub>) and dolomite (CaMg(CO<sub>3</sub>)<sub>2</sub>) are added to soils by land managers to increase soil pH (i.e., to reduce acidification). Carbon dioxide emissions occur as these compounds react with hydrogen ions in soils. The rate of degradation of applied limestone and dolomite depends on the soil conditions, soil type, climate regime, and whether limestone or dolomite is applied. Emissions from liming of soils have fluctuated over the past 25 years in the United States, ranging from 3.2 MMT CO<sub>2</sub> Eq. to 6.0 MMT CO<sub>2</sub> Eq. In 2017, liming of soils in the United States resulted in emissions of 3.2 MMT CO<sub>2</sub> Eq. (0.9 MMT C), representing a 32 percent decrease in emissions since 1990 (see Table 5-20 and Table 5-21). The trend is driven by variation in the amount of limestone and dolomite applied to soils over the time period.

**Table 5-20: Emissions from Liming (MMT CO<sub>2</sub> Eq.)**

Source	1990	2005	2013	2014	2015	2016	2017
Limestone	4.1	3.9	3.6	3.3	3.5	2.9	2.9
Dolomite	0.6	0.4	0.3	0.3	0.3	0.3	0.3
<b>Total</b>	<b>4.7</b>	<b>4.3</b>	<b>3.9</b>	<b>3.6</b>	<b>3.7</b>	<b>3.2</b>	<b>3.2</b>

Note: Totals may not sum due to independent rounding.

**Table 5-21: Emissions from Liming (MMT C)**

Source	1990	2005	2013	2014	2015	2016	2017
Limestone	1.1	1.1	1.0	0.9	0.9	0.8	0.8
Dolomite	0.2	0.1	0.1	0.1	0.1	0.1	0.1
<b>Total</b>	<b>1.3</b>	<b>1.2</b>	<b>1.1</b>	<b>1.0</b>	<b>1.0</b>	<b>0.9</b>	<b>0.9</b>

Note: Totals may not sum due to independent rounding.

## Methodology

Carbon dioxide emissions from application of limestone and dolomite to soils were estimated using a Tier 2 methodology consistent with IPCC (2006). The annual amounts of limestone and dolomite applied (see Table 5-22) were multiplied by CO<sub>2</sub> emission factors from West and McBride (2005). These emission factors (0.059 metric ton C/metric ton limestone, 0.064 metric ton C/metric ton dolomite) are lower than the IPCC default emission factors because they account for the portion of carbonates that are transported from soils through hydrological processes and eventually deposited in ocean basins (West and McBride 2005). This analysis of lime dissolution is based on studies in the Mississippi River basin, where the vast majority of lime application occurs in the United States (West 2008). Moreover, much of the remaining lime application is occurring under similar precipitation regimes, and so



the emission factors are considered a reasonable approximation for all lime application in the United States (West 2008).

The annual application rates of limestone and dolomite were derived from estimates and industry statistics provided in the *Minerals Yearbook* and *Mineral Industry Surveys* (Tepordei 1993 through 2006; Willett 2007a, 2007b, 2009, 2010, 2011a, 2011b, 2013a, 2014, 2015, 2016, 2017, 2018; USGS 2008 through 2018). The U.S. Geological Survey (USGS; U.S. Bureau of Mines prior to 1997) compiled production and use information through surveys of crushed stone manufacturers. However, manufacturers provided different levels of detail in survey responses so the estimates of total crushed limestone and dolomite production and use were divided into three components: (1) production by end-use, as reported by manufacturers (i.e., “specified” production); (2) production reported by manufacturers without end-uses specified (i.e., “unspecified” production); and (3) estimated additional production by manufacturers who did not respond to the survey (i.e., “estimated” production).

#### Box 5-5: Comparison of the Tier 2 U.S. Inventory Approach and IPCC (2006) Default Approach

Emissions from liming of soils were estimated using a Tier 2 methodology based on emission factors specific to the United States that are lower than the IPCC (2006) emission default factors. Most lime application in the United States occurs in the Mississippi River basin, or in areas that have similar soil and rainfall regimes as the Mississippi River basin. Under these conditions, a significant portion of dissolved agricultural lime leaches through the soil into groundwater. Groundwater moves into channels and is transported to larger rivers and eventually the ocean where  $\text{CaCO}_3$  precipitates to the ocean floor (West and McBride 2005). The U.S.-specific emission factors (0.059 metric ton C/metric ton limestone and 0.064 metric ton C/metric ton dolomite) are about half of the IPCC (2006) emission factors (0.12 metric ton C/metric ton limestone and 0.13 metric ton C/metric ton dolomite). For comparison, the 2017 U.S. emission estimate from liming of soils is 3.2 MMT  $\text{CO}_2$  Eq. using the U.S.-specific factors. In contrast, emissions would be estimated at 6.5 MMT  $\text{CO}_2$  Eq. using the IPCC (2006) default emission factors.

Data on “specified” limestone and dolomite amounts were used directly in the emission calculation because the end use is provided by the manufacturers and can be used to directly determine the amount applied to soils. However, it is not possible to determine directly how much of the limestone and dolomite is applied to soils for manufacturer surveys in the “unspecified” and “estimated” categories. For these categories, the amounts of crushed limestone and dolomite applied to soils were determined by multiplying the percentage of total “specified” limestone and dolomite production that is applied to soils, by the total amounts of “unspecified” and “estimated” limestone and dolomite production. In other words, the proportion of total “unspecified” and “estimated” crushed limestone and dolomite that was applied to soils is proportional to the amount of total “specified” crushed limestone and dolomite that was applied to soils.

In addition, data were not available for 1990, 1992 and 2017 on the fractions of total crushed stone production that were limestone and dolomite, and on the fractions of limestone and dolomite production that were applied to soils. To estimate the 1990 and 1992 data, a set of average fractions were calculated using the 1991 and 1993 data. These average fractions were applied to the quantity of “total crushed stone produced or used” reported for 1990 and 1992 in the 1994 *Minerals Yearbook* (Tepordei 1996). To estimate 2017 data, 2016 fractions were applied to a 2017 estimate of total crushed stone presented in the USGS *Mineral Industry Surveys: Crushed Stone and Sand and Gravel in the First Quarter of 2018* (USGS 2018).

The primary source for limestone and dolomite activity data is the *Minerals Yearbook*, published by the Bureau of Mines through 1994 and by the USGS from 1995 to the present. In 1994, the “Crushed Stone” chapter in the *Minerals Yearbook* began rounding (to the nearest thousand metric tons) quantities for total crushed stone produced or used. It then reported revised (rounded) quantities for each of the years from 1990 to 1993. In order to minimize the inconsistencies in the activity data, these revised production numbers have been used in all of the subsequent calculations.

**Table 5-22: Applied Minerals (MMT)**

Mineral	1990	2005	2013	2014	2015	2016	2017
Limestone	19.0	18.1	16.4	15.3	16.0	13.5	13.4
Dolomite	2.4	1.9	1.5	1.3	1.2	1.2	1.2

## Uncertainty and Time-Series Consistency

Uncertainty regarding the amount of limestone and dolomite applied to soils was estimated at  $\pm 15$  percent with normal densities (Tepordei 2003; Willett 2013b). Analysis of the uncertainty associated with the emission factors included the fraction of lime dissolved by nitric acid versus the fraction that reacts with carbonic acid, and the portion of bicarbonate that leaches through the soil and is transported to the ocean. Uncertainty regarding the time associated with leaching and transport was not addressed in this analysis, but is assumed to be a relatively small contributor to the overall uncertainty (West 2005). The probability distribution functions for the fraction of lime dissolved by nitric acid and the portion of bicarbonate that leaches through the soil were represented as triangular distributions between ranges of zero and 100 percent of the estimates. The uncertainty surrounding these two components largely drives the overall uncertainty.

A Monte Carlo (Approach 2) uncertainty analysis was applied to estimate the uncertainty in CO<sub>2</sub> emissions from liming. The results of the Approach 2 quantitative uncertainty analysis are summarized in Table 5-23. Carbon dioxide emissions from carbonate lime application to soils in 2017 were estimated to be between -0.35 and 6.01 MMT CO<sub>2</sub> Eq. at the 95 percent confidence level. This confidence interval represents a range of 111 percent below to 89 percent above the 2017 emission estimate of 3.2 MMT CO<sub>2</sub> Eq. Note that there is a small probability of a negative emissions value leading to a net uptake of CO<sub>2</sub> from the atmosphere. Net uptake occurs due to the dominance of the carbonate lime dissolving in carbonic acid rather than nitric acid (West and McBride 2005).

**Table 5-23: Approach 2 Quantitative Uncertainty Estimates for CO<sub>2</sub> Emissions from Liming (MMT CO<sub>2</sub> Eq. and Percent)**

Source	Gas	2017 Emission Estimate (MMT CO <sub>2</sub> Eq.)	Uncertainty Range Relative to Emission Estimate <sup>a</sup>			
			(MMT CO <sub>2</sub> Eq.)		(%)	
			Lower Bound	Upper Bound	Lower Bound	Upper Bound
Liming	CO <sub>2</sub>	3.2	(0.35)	6.01	-111%	+89%

<sup>a</sup>Range of emission estimates predicted by Monte Carlo Stochastic Simulation for a 95 percent confidence interval.

Methodological recalculations were applied to the entire time series to ensure time-series consistency from 1990 through 2017.

## QA/QC and Verification

A source-specific QA/QC plan for liming has been developed and implemented, and the quality control effort focused on the Tier 1 procedures for this Inventory. No errors were found.

## Recalculations Discussion

Adjustments were made in the current Inventory to improve the results. First, limestone and dolomite application data for 2015 and 2016 were updated with the recently published data from USGS (2018), rather than being approximated by a ratio method for 2016. With this revision in the activity data, the emissions decreased by 1.1 and 17 percent for 2015 and 2016, respectively, relative to the previous Inventory estimates.

## 5.6 Urea Fertilization (CRF Source Category 3H)

The use of urea (CO(NH<sub>2</sub>)<sub>2</sub>) as a fertilizer leads to greenhouse gas emissions through the release of CO<sub>2</sub> that was fixed during the industrial production process. In the presence of water and urease enzymes, urea is converted into ammonium (NH<sub>4</sub><sup>+</sup>), hydroxyl ion (OH<sup>-</sup>), and bicarbonate (HCO<sub>3</sub><sup>-</sup>). The bicarbonate then evolves into CO<sub>2</sub> and water. Emissions from urea fertilization in the United States totaled 5.1 MMT CO<sub>2</sub> Eq. (1.4 MMT C) in 2017 (Table 5-24 and Table 5-25). Carbon dioxide emissions have increased by 109 percent between 1990 and 2017 due to an increasing amount of urea that is applied to soils.

**Table 5-24: CO<sub>2</sub> Emissions from Urea Fertilization (MMT CO<sub>2</sub> Eq.)**

Source	1990	2005	2013	2014	2015	2016	2017
Urea Fertilization	2.4	3.5	4.4	4.5	4.7	4.9	5.1

**Table 5-25: CO<sub>2</sub> Emissions from Urea Fertilization (MMT C)**

Source	1990	2005	2013	2014	2015	2016	2017
Urea Fertilization	0.7	1.0	1.2	1.2	1.3	1.3	1.4

## Methodology

Carbon dioxide emissions from the application of urea to agricultural soils were estimated using the IPCC (2006) Tier 1 methodology. The method assumes that all CO<sub>2</sub> fixed during the industrial production process of urea are released after application. The annual amounts of urea applied to croplands (see Table 5-26) were derived from the state-level fertilizer sales data provided in *Commercial Fertilizer* reports (TVA 1991, 1992, 1993, 1994; AAPFCO 1995 through 2018).<sup>20</sup> These amounts were multiplied by the default IPCC (2006) emission factor (0.20 metric tons of C per metric ton of urea), which is equal to the C content of urea on an atomic weight basis. Because fertilizer sales data are reported in fertilizer years (July previous year through June current year), a calculation was performed to convert the data to calendar years (January through December). According to monthly fertilizer use data (TVA 1992b), 35 percent of total fertilizer used in any fertilizer year is applied between July and December of the previous calendar year, and 65 percent is applied between January and June of the current calendar year.

Fertilizer sales data for the 2016 and 2017 fertilizer years (i.e., July 2015 through June 2016 and July 2016 through June 2017) were not available for this Inventory. Therefore, urea application in the 2016 and 2017 fertilizer years were estimated using a linear, least squares trend of consumption over the data from the previous five years (2011 through 2015) at the state scale. A trend of five years was chosen as opposed to a longer trend as it best captures the current inter-state and inter-annual variability in consumption. State-level estimates of CO<sub>2</sub> emissions from the application of urea to agricultural soils were summed to estimate total emissions for the entire United States. The fertilizer year data is then converted into calendar year data using the method described above.

**Table 5-26: Applied Urea (MMT)**

	1990	2005	2013	2014	2015	2016	2017
Urea Fertilizer <sup>a</sup>	3.3	4.8	6.1	6.2	6.5	6.7	6.9

<sup>a</sup>These numbers represent amounts applied to all agricultural land, including *Cropland Remaining Cropland*, *Land Converted to Cropland*, *Grassland Remaining Grassland*, *Land Converted to Grassland*, *Settlements Remaining Settlements*, *Land Converted to Settlements*, *Forest Land Remaining Forest Land* and *Land Converted to Forest Land*, as it is not currently possible to apportion the data by land-use category.

## Uncertainty and Time-Series Consistency

Uncertainty estimates are presented in Table 5-27 for urea fertilization. An Approach 2 Monte Carlo analysis was completed. The largest source of uncertainty was the default emission factor, which assumes that 100 percent of the C in CO(NH<sub>2</sub>)<sub>2</sub> applied to soils is ultimately emitted into the environment as CO<sub>2</sub>. This factor does not incorporate the possibility that some of the C may be retained in the soil, and therefore the uncertainty range was set from 0 percent emissions to the maximum emission value of 100 percent using a triangular distribution. In addition, urea consumption data also have uncertainty that is propagated through the emission calculation using a Monte Carlo simulation approach as described by the IPCC (2006). Carbon dioxide emissions from urea fertilization of agricultural soils in 2017 were estimated to be between 2.89 and 5.21 MMT CO<sub>2</sub> Eq. at the 95 percent confidence

<sup>20</sup> The amount of urea consumed for non-agricultural purposes in the United States is reported in the Industrial Processes and Product Use chapter, Section 4.6 Urea Consumption for Non-Agricultural Purposes.

level. This indicates a range of 43 percent below to 3 percent above the 2017 emission estimate of 5.1 MMT CO<sub>2</sub> Eq.

**Table 5-27: Quantitative Uncertainty Estimates for CO<sub>2</sub> Emissions from Urea Fertilization (MMT CO<sub>2</sub> Eq. and Percent)**

Source	Gas	2017 Emission Estimate (MMT CO <sub>2</sub> Eq.)	Uncertainty Range Relative to Emission Estimate <sup>a</sup>			
			(MMT CO <sub>2</sub> Eq.)		(%)	
			Lower Bound	Upper Bound	Lower Bound	Upper Bound
Urea Fertilization	CO <sub>2</sub>	5.1	2.89	5.21	-43%	3%

<sup>a</sup>Range of emission estimates predicted by Monte Carlo Stochastic Simulation for a 95 percent confidence interval.

There are additional uncertainties that are not quantified in this analysis. Urea for non-fertilizer use, such as aircraft deicing, may be included in consumption totals, but the amount is likely very small. For example, research on aircraft deicing practices based on a 1992 survey found a known annual usage of approximately 2,000 tons of urea for deicing; this would constitute 0.06 percent of the 1992 consumption of urea (EPA 2000). Similarly, surveys conducted from 2002 to 2005 indicate that total urea use for deicing at U.S. airports is estimated to be 3,740 metric tons per year, or less than 0.07 percent of the fertilizer total for 2007 (Itle 2009). In addition, there is uncertainty surrounding the underlying assumptions behind the calculation that converts fertilizer years to calendar years. These uncertainties are negligible over multiple years because an over- or under-estimated value in one calendar year is addressed with corresponding increase or decrease in the value for the subsequent year.

Methodological recalculations were applied to the entire time series to ensure time-series consistency from 1990 through 2017. Details on the emission trends through time are described in more detail in the Introduction, above.

## QA/QC and Verification

A source-specific QA/QC plan for Urea Fertilization has been developed and implemented, and no errors were found in the calculation. Based on the quality control review, it was not clear if Urea Ammonium Nitrate (UAN) should also be included as a source of CO<sub>2</sub> emissions. This will be further investigated in a future Inventory.

## Recalculations Discussion

Recalculations resulted from updated urea application estimates in a new AAPFCO report (2018). New activity data for 2015 were applied to all states; 2016 and 2017 estimates were derived using the new data for 2011 and 2015. This resulted in an emissions decrease for the United States of 0.5 percent in 2014, 3.3 percent in 2015, and 4.3 percent in 2016.

## Planned Improvements

A key planned improvement is to investigate of the composition of Urea Ammonium Nitrate (UAN), and determine if UAN should be included in the estimation of Urea CO<sub>2</sub> emissions. In addition, the analytical solutions to the IPCC Tier 1 method have been used as the estimates in this report. It would be more robust to use the uncertainty analysis as the basis for the estimates, rather than a separate analytical calculation. These improvements will be implemented in a future Inventory.

## 5.7 Field Burning of Agricultural Residues (CRF Source Category 3F)

Crop production creates large quantities of agricultural crop residues, which farmers manage in a variety of ways. For example, crop residues can be left in the field and possibly incorporated into the soil with tillage; collected and used as fuel, animal bedding material, supplemental animal feed, or construction material; composted and applied to soils; transported to landfills; or burned in the field. Field burning of crop residues is not considered a net source of CO<sub>2</sub> emissions because the C released to the atmosphere as CO<sub>2</sub> during burning is reabsorbed during the next growing season by the crop. However, crop residue burning is a net source of CH<sub>4</sub>, N<sub>2</sub>O, CO, and NO<sub>x</sub>, which are released during combustion.

In the United States, field burning of agricultural residues commonly occurs in southeastern states, the Great Plains, and the Pacific Northwest (McCarty 2011). The primary crops that are managed with residue burning include corn, cotton, lentils, rice, soybeans, and wheat (McCarty 2009). In 2017, CH<sub>4</sub> and N<sub>2</sub>O emissions from field burning of agricultural residues were 0.2 MMT CO<sub>2</sub> Eq. (8 kt) and 0.1 MMT CO<sub>2</sub> Eq. (0.3 kt), respectively (Table 5-28 and Table 5-29). Annual emissions of CH<sub>4</sub> and N<sub>2</sub>O have increased from 1990 to 2017 by 82 percent and 72 percent, respectively. The increase in emissions over time is partly due to larger amounts of residue production with higher yielding crop varieties and fuel loads, but also linked with an increase in area burned for some of the crops.

**Table 5-28: CH<sub>4</sub> and N<sub>2</sub>O Emissions from Field Burning of Agricultural Residues (MMT CO<sub>2</sub> Eq.)**

Gas/Crop Type	1990	2005	2013	2014	2015	2016	2017
<b>CH<sub>4</sub></b>	<b>0.1</b>	<b>0.2</b>	<b>0.2</b>	<b>0.2</b>	<b>0.2</b>	<b>0.2</b>	<b>0.2</b>
Maize	+	+	0.1	0.1	0.1	0.1	0.1
Rice	+	0.1	+	+	+	+	+
Wheat	+	+	+	+	+	+	+
Barley	+	+	+	+	+	+	+
Oats	+	+	+	+	+	+	+
Other Small Grains	+	+	+	+	+	+	+
Sorghum	+	+	+	+	+	+	+
Cotton	+	+	+	+	+	+	+
Grass Hay	+	+	+	+	+	+	+
Legume Hay	+	+	+	+	+	+	+
Peas	+	+	+	+	+	+	+
Sunflower	+	+	+	+	+	+	+
Tobacco	+	+	+	+	+	+	+
Vegetables	+	+	+	+	+	+	+
Chickpeas	+	+	+	+	+	+	+
Dry Beans	+	+	+	+	+	+	+
Lentils	+	+	+	+	+	+	+
Peanuts	+	+	+	+	+	+	+
Soybeans	+	+	+	+	+	+	+
Potatoes	+	+	+	+	+	+	+
Sugarbeets	+	+	+	+	+	+	+
<b>N<sub>2</sub>O</b>	<b>+</b>	<b>0.1</b>	<b>0.1</b>	<b>0.1</b>	<b>0.1</b>	<b>0.1</b>	<b>0.1</b>
Maize	+	+	+	+	+	+	+
Rice	+	+	+	+	+	+	+
Wheat	+	+	+	+	+	+	+

Barley	+		+		+	+	+	+
Oats	+		+		+	+	+	+
Other Small Grains	+		+		+	+	+	+
Sorghum	+		+		+	+	+	+
Cotton	+		+		+	+	+	+
Grass Hay	+		+		+	+	+	+
Legume Hay	+		+		+	+	+	+
Peas	+		+		+	+	+	+
Sunflower	+		+		+	+	+	+
Tobacco	+		+		+	+	+	+
Vegetables	+		+		+	+	+	+
Chickpeas	+		+		+	+	+	+
Dry Beans	+		+		+	+	+	+
Lentils	+		+		+	+	+	+
Peanuts	+		+		+	+	+	+
Soybeans	+		+		+	+	+	+
Potatoes	+		+		+	+	+	+
Sugarbeets	+		+		+	+	+	+
<b>Total</b>	<b>0.2</b>		<b>0.3</b>		<b>0.3</b>	<b>0.3</b>	<b>0.3</b>	<b>0.3</b>

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

Note: Totals may not sum due to independent rounding.

**Table 5-29: CH<sub>4</sub>, N<sub>2</sub>O, CO, and NO<sub>x</sub> Emissions from Field Burning of Agricultural Residues (kt)**

Gas/Crop Type	1990	2005	2013	2014	2015	2016	2017
<b>CH<sub>4</sub></b>	<b>4</b>	<b>7</b>	<b>8</b>	<b>8</b>	<b>8</b>	<b>8</b>	<b>8</b>
Maize	+	1	3	3	3	3	3
Rice	2	2	2	1	2	1	2
Wheat	1	2	1	1	1	1	1
Barley	+	+	+	+	+	+	+
Oats	+	+	+	+	+	+	+
Other Small Grains	+	+	+	+	+	+	+
Sorghum	+	+	+	+	+	+	+
Cotton	+	1	1	1	1	1	1
Grass Hay	+	+	+	+	+	+	+
Legume Hay	+	+	+	+	+	+	+
Peas	+	+	+	+	+	+	+
Sunflower	+	+	+	+	+	+	+
Tobacco	+	+	+	+	+	+	+
Vegetables	+	+	+	+	+	+	+
Chickpeas	+	+	+	+	+	+	+
Dry Beans	+	+	+	+	+	+	+
Lentils	+	+	+	+	+	+	+
Peanuts	+	+	+	+	+	+	+
Soybeans	+	1	1	1	1	1	1
Potatoes	+	+	+	+	+	+	+
Sugarbeets	+	+	+	+	+	+	+



<b>N<sub>2</sub>O</b>	+		+		+	+	+	+	+
Maize	+		+		+	+	+	+	+
Rice	+		+		+	+	+	+	+
Wheat	+		+		+	+	+	+	+
Barley	+		+		+	+	+	+	+
Oats	+		+		+	+	+	+	+
Other Small Grains	+		+		+	+	+	+	+
Sorghum	+		+		+	+	+	+	+
Cotton	+		+		+	+	+	+	+
Grass Hay	+		+		+	+	+	+	+
Legume Hay	+		+		+	+	+	+	+
Peas	+		+		+	+	+	+	+
Sunflower	+		+		+	+	+	+	+
Tobacco	+		+		+	+	+	+	+
Vegetables	+		+		+	+	+	+	+
Chickpeas	+		+		+	+	+	+	+
Dry Beans	+		+		+	+	+	+	+
Lentils	+		+		+	+	+	+	+
Peanuts	+		+		+	+	+	+	+
Soybeans	+		+		+	+	+	+	+
Potatoes	+		+		+	+	+	+	+
Sugarbeets	+		+		+	+	+	+	+
<b>CO</b>	<b>89</b>		<b>154</b>		<b>157</b>	<b>152</b>	<b>148</b>	<b>144</b>	<b>141</b>
<b>NO<sub>x</sub></b>	<b>4</b>		<b>6</b>		<b>6</b>	<b>6</b>	<b>6</b>	<b>6</b>	<b>6</b>

+ Does not exceed 0.5 kt

Note: Totals may not sum due to independent rounding.

## Methodology

A U.S.-specific Tier 2 method is used to estimate greenhouse gas emissions from field burning of agricultural residues from 1990 to 2012 (for more details comparing the U.S.-specific approach to the IPCC (2006) default approach, see Box 5-6). In order to estimate the amounts of C and N released during burning, the following equation is used:

$$\text{C or N released} = \sum \text{for all crop types and states} \left[ \frac{\text{AB}}{\text{CAH} \times \text{CP} \times \text{RCR} \times \text{DMF} \times \text{BE} \times \text{CE} \times (\text{FC or FN})} \right]$$

where,

- Area Burned (AB) = Total area of crop burned, by state
- Crop Area Harvested (CAH) = Total area of crop harvested, by state
- Crop Production (CP) = Annual production of crop in kt, by state
- Residue: Crop Ratio (RCR) = Amount of residue produced per unit of crop production
- Dry Matter Fraction (DMF) = Amount of dry matter per unit of biomass for a crop
- Fraction of C or N (FC or FN) = Amount of C or N per unit of dry matter for a crop
- Burning Efficiency (BE) = The proportion of prefire fuel biomass consumed<sup>21</sup>
- Combustion Efficiency (CE) = The proportion of C or N released with respect to the total amount of C or N available in the burned material, respectively

<sup>21</sup> In IPCC/UNEP/OECD/IEA (1997), the equation for C or N released contains the variable ‘fraction oxidized in burning’. This variable is equivalent to (burning efficiency × combustion efficiency).

Crop production data are available by state and year from USDA (2017) for twenty-one crops that are burned in the conterminous United States, including maize, rice, wheat, barley, oats, other small grains, sorghum, cotton, grass hay, legume hay, peas, sunflower, tobacco, vegetables, chickpeas, dry beans, lentils, peanuts, soybeans, potatoes, and sugarbeets. Crop area data are based on the 2012 National Resources Inventory (NRI) (USDA-NRCS 2015). In order to estimate total crop production, the crop yield data from USDA Quick Stats crop yields is multiplied by the NRI crop areas. The production data for the crop types are presented in Table 5-30. Alaska and Hawaii are not included in the current analysis, but there is a planned improvement to estimate residue burning emissions for these two states in a future Inventory.

The amount of elemental C or N released through oxidation of the crop residues is used in the following equation to estimate CH<sub>4</sub>, CO, N<sub>2</sub>O, and NO<sub>x</sub> emissions from the Field Burning of Agricultural Residues:

$$\text{CH}_4 \text{ and CO, or N}_2\text{O and NO}_x = \text{C or N Released} \times \text{ER} \times \text{CF}$$

where,

$$\begin{aligned} \text{Emissions Ratio (ER)} &= \text{g CH}_4\text{-C or CO-C/g C released, or g N}_2\text{O-N or NO}_x\text{-N/g N released} \\ \text{Conversion Factor (CF)} &= \text{conversion, by molecular weight ratio, of CH}_4\text{-C to C (16/12), or CO-C to C (28/12), or N}_2\text{O-N to N (44/28), or NO}_x\text{-N to N (30/14)} \end{aligned}$$

Emissions from Field Burning of Agricultural Residues are calculated using a Tier 2 methodology that is based on the method developed by the IPCC/UNEP/OECD/IEA (1997). The rationale for using the IPCC/UNEP/OECD/IEA (1997) approach rather than the method provided in the 2006 IPCC Guidelines is as follows: (1) the equations from both guidelines rely on the same underlying variables (though the formats differ); (2) the IPCC (2006) equation was developed to be broadly applicable to all types of biomass burning, and, thus, is not specific to agricultural residues; (3) the IPCC (2006) method provides emission factors based on the dry matter content rather emission rates related to the amount of C and N in the residues; and (4) the IPCC (2006) default factors are provided only for four crops (corn, rice, sugarcane, and wheat) while this Inventory includes emissions from twenty-one crops.

A comparison of the methods and factors used in: (1) the current Inventory and (2) the default IPCC (2006) approach was undertaken for the time series from 1990 through 2012 to determine the difference in overall estimates between the two approaches. To estimate greenhouse gas emissions from field burning of agricultural residues using the IPCC (2006) methodology, the following equation—cf. IPCC (2006) Equation 2.27—was used:

$$\text{Emissions (kt)} = \text{AB} \times (\text{M}_B \times \text{C}_f) \times \text{G}_{\text{ef}} \times 10^{-6}$$

where,

$$\begin{aligned} \text{Area Burned (AB)} &= \text{Total area of crop burned (ha)} \\ \text{Mass Burned (M}_B \times \text{C}_f) &= \text{IPCC (2006) default carbon fractions with fuel biomass consumption US-Specific Values using NASS Statistics}^{22} \text{ (metric tons dry matter burnt ha}^{-1}\text{)} \\ \text{Emission Factor (G}_{\text{ef}}) &= \text{IPCC (2006) emission factor (g kg}^{-1}\text{ dry matter burnt)} \end{aligned}$$

The IPCC (2006) Tier 1 method approach that utilizes default combustion factors and emission factors with mass of fuel values derived from national datasets resulted in 22 percent lower emissions of CH<sub>4</sub> and 44 percent lower emissions of N<sub>2</sub>O compared to this Inventory. In summary, the IPCC/UNEP/OECD/IEA (1997) method is considered more appropriate for U.S. conditions because it is more flexible for incorporating country-specific data and emissions are estimated based on specific C and N content of the fuel, which is converted into CH<sub>4</sub>, CO, N<sub>2</sub>O and NO<sub>x</sub>, compared to IPCC (2006) approach that is based on dry matter rather than elemental composition.

<sup>22</sup> NASS yields are used to derive mass of fuel values because IPCC (2006) only provides default values for 4 of the 21 crops included in the Inventory.

**Table 5-30: Agricultural Crop Production (kt of Product)**

Crop	1990	2005	2011	2012
Maize	294,558	370,338	389,657	341,374
Rice	9,476	11,914	10,172	9,829
Wheat	79,920	68,919	61,609	70,867
Barley	9,087	5,042	3,778	5,293
Oats	5,972	2,632	1,625	1,631
Other Small Grains	2,639	2,007	1,237	1,681
Sorghum	23,688	13,080	9,344	10,422
Cotton	4,609	6,227	5,343	5,666
Grass Hay	+	+	+	464,050
Legume Hay	64	706	302	534
Peas	992	1,397	792	1,204
Sunflower	1,151	347	264	521
Tobacco	+	903	1,189	2,027
Vegetables	+	7	+	+
Chickpeas	638	1,084	1,079	1,159
Dry Beans	+	119	59	121
Lentils	1,822	2,242	1,906	2,649
Peanuts	56,613	87,164	86,839	84,805
Soybeans	18,960	19,471	20,296	20,517
Potatoes	25,017	26,604	28,922	28,488
Sugarbeets	+	+	+	+

The area burned is determined based on an analysis of remote sensing products (McCarty et al. 2009, 2010, 2011). The presence of fires have been analyzed at 3600 survey locations in the NRI from 1990 to 2002 with LANDFIRE data products developed from 30m Landsat imagery (LANDFIRE 2014), and from 2003 through 2012 using 1km Moderate Resolution Imaging Spectroradiometer imagery (MODIS) Global Fire Location Product (MCD14ML) using combined observations from Terra and Aqua satellites (Giglio et al. 2006). A sample of states are included in the analysis with high, medium and low burning rates for agricultural residues, including Arkansas, California, Florida, Indiana, Iowa and Washington. The area burned is determined directly from the analysis for these states.

For other states within the conterminous United States, the area burned is estimated from a logistical regression model that has been developed from the data collected from the remote sensing products for the six states. The logistical regression model is used to predict occurrence of fire events. Several variables are tested in the logistical regression including a) the historical level of burning in each state (high, medium or low levels of burning) based on an analysis by McCarty et al. (2011), b) existence of state laws limiting burning of fields, in addition to c) mean annual precipitation and mean annual temperature from a 4 kilometer gridded product developed by the PRISM Climate Group (2015). A K-fold model fitting procedure is used due to low frequency of burning and likelihood that outliers could influence the model fit. Specifically the model is trained with a random selection of sample locations and evaluated with the remaining sample. This process is repeated ten times to select a model that is most common among the set of ten, and avoid models that appear to be influenced by outliers due to the random draw of survey locations for training the model. In order to address uncertainty, a Monte Carlo analysis is used to sample the parameter estimates for the logistical regression model and produce one thousand estimates of burning for each crop in the remaining forty-two states included in this Inventory. State-level area burned data are divided by state-level crop area data to estimate the percent of crop area burned by crop type for each state. Table 5-31 shows the resulting percentage of crop residue burned at the national scale by crop type. State-level estimates are also available upon request.

**Table 5-31: U.S. Average Percent Crop Area Burned by Crop (Percent)**

Crop	1990	2005	2011	2012
Maize	+	+	+	+

Rice	5%		6%		2%	3%
Wheat	+		+		1%	1%
Barley	1%		+		+	+
Oats	+		+		+	1%
Other Small Grains	+		+		+	+
Sorghum	+		+		1%	1%
Cotton	+		1%		1%	1%
Grass Hay	+		+		+	+
Legume Hay	+		+		+	+
Peas	+		+		1%	+
Sunflower	+		+		+	+
Tobacco	+		1%		1%	1%
Vegetables	+		+		+	+
Chickpeas	+		+		+	+
Dry Beans	+		+		+	+
Lentils	+		+		+	+
Peanuts	1%		1%		2%	2%
Soybeans	+		+		+	+
Potatoes	+		+		+	+
Sugarbeets	+		+		+	+

+ Does not exceed 0.5%

Additional parameters are needed to estimate the amount of burning, including residue:crop ratios, dry matter fractions, carbon fractions, nitrogen fractions, burning efficiency and combustion efficiency. Residue:crop product mass ratios, residue dry matter fractions and the residue N contents are obtained from several sources (IPCC 2006 and sources at bottom of Table 5-32). The residue C contents for all crops are based on IPCC (2006) default value for herbaceous biomass. The burning efficiency is assumed to be 93 percent, and the combustion efficiency is assumed to be 88 percent, for all crop types (EPA 1994). See Table 5-32 for a summary of the crop-specific conversion factors. Emission ratios and mole ratio conversion factors for all gases are based on the *Revised 1996 IPCC Guidelines* (IPCC/UNEP/OECD/IEA 1997) (see Table 5-33).

**Table 5-32: Parameters for Estimating Emissions from Field Burning of Agricultural Residues**

Crop	Residue/Crop Ratio	Dry Matter Fraction	Carbon Fraction	Nitrogen Fraction	Burning Efficiency (Fraction)	Combustion Efficiency (Fraction)
Maize	0.707	0.56	0.47	0.01	0.93	0.88
Rice	1.340	0.89	0.47	0.01	0.93	0.88
Wheat	1.725	0.89	0.47	0.01	0.93	0.88
Barley	1.181	0.89	0.47	0.01	0.93	0.88
Oats	1.374	0.89	0.47	0.01	0.93	0.88
Other Small Grains	1.777	0.88	0.47	0.01	0.93	0.88
Sorghum	0.780	0.60	0.47	0.01	0.93	0.88
Cotton	7.443	0.93	0.47	0.01	0.93	0.88
Grass Hay	0.208	0.90	0.47	0.02	0.93	0.88
Legume Hay	0.290	0.67	0.47	0.01	0.93	0.88
Peas	1.677	0.91	0.47	0.01	0.93	0.88
Sunflower	1.765	0.88	0.47	0.01	0.93	0.88
Tobacco	0.300	0.87	0.47	0.01	0.93	0.88
Vegetables	0.708	0.08	0.47	0.01	0.93	0.88
Chickpeas	1.588	0.91	0.47	0.01	0.93	0.88

Dry Beans	0.771	0.90	0.47	0.01	0.93	0.88
Lentils	1.837	0.91	0.47	0.02	0.93	0.88
Peanuts	1.600	0.94	0.47	0.02	0.93	0.88
Soybeans	1.500	0.91	0.47	0.01	0.93	0.88
Potatoes	0.379	0.25	0.47	0.02	0.93	0.88
Sugarbeets	0.196	0.22	0.47	0.02	0.93	0.88

Notes:

Chickpeas: IPCC 2006, Table 11.2; Beans & pulses

Cotton: Combined sources (Heitholt et al. 1992, Halevy 1976, Wells and Meredith 1984, Sadras and Wilson 1997, Pettigrew and Meredith 1997, Torbert and Reeves 1994, Gerik et al. 1996, Brouder and Cassmen 1990, Fritschi et al. 2003, Pettigrew et al. 2005, Bouquet and Breitenbeck 2000, Mahroni and Aharonov 1964, Bange and Milroy 2004, Hollifield et al. 2000, Mondino et al. 2004, Wallach et al. 1978) Lentils: IPCC 2006, Table 11.2; Beans & pulses

Peas: IPCC 2006, Table 11.2; Beans & pulses

Peanuts: IPCC 2006; Table 11.2; Root ratio and belowground N content are from Root crops, other

Sugarbeets: IPCC 2006; Table 11.2; values are for Tubers

Sunflower: IPCC 2006, Table 11.2; values are from Grains

Sugarcane: combined sources (Wiedenfels 2000, Dua and Sharma 1976, Singels & Bezuidenhout 2002, Stirling et al. 1999, Sitompul et al. 2000)

Tobacco: combined sources (Beyaert 1996, Moustakas and Ntzanis 2005, Crafts-Brandner et al. 1994, Hopkinson 1967, Crafts-Brandner et al. 1987)

Vegetables:

Carrots: McPharlin et al. 1992; Gibberd et al. 2003; Reid and English 2000; Peach et al. 2000; see IPCC Tubers for R:S and N frac

Lettuce, cabbage: combines sources (Huett and Dettman 1991; De Pinheiro Henriques & Marcelis 2000; Huett and Dettman 1989; Peach et al. 2000; Kage et al. 2003; Tan et al. 1999; Kumar et al. 1994; MacLeod et al. 1971; Jacobs et al. 2004; Jacobs et al. 2001; Jacobs et al. 2002); IPCC Grains for N frac

Melons: Valantin et al. 1999; squash for R:S; IPCC Grains for N frac

Onion: Peach et al. 2000, Halvorson et al. 2002; IPCC 2006 Tubers for N fractions

Peppers: combined sources (Costa and Gianquinto 2002; Marcussi et al. 2004; Tadesse et al. 1999; Diaz-Perez et al. 2008); IPCC Grains for N frac

Tomatoes: Scholberg et al. 2000a,b; Akintoye et al. 2005; AGR-N and BGR-N are from Grains

**Table 5-33: Greenhouse Gas Emission Ratios and Conversion Factors**

Gas	Emission Ratio	Conversion Factor
CH <sub>4</sub> :C	0.005 <sup>a</sup>	16/12
CO:C	0.060 <sup>a</sup>	28/12
N <sub>2</sub> O:N	0.007 <sup>b</sup>	44/28
NO <sub>x</sub> :N	0.121 <sup>b</sup>	30/14

<sup>a</sup> Mass of C compound released (units of C) relative to mass of total C released from burning (units of C).

<sup>b</sup> Mass of N compound released (units of N) relative to mass of total N released from burning (units of N).

For this Inventory, new activity data on crop areas are not available for 2013 to 2017 from the USDA National Resources Inventory (USDA-NRCS 2015). To complete the emissions time series, a linear extrapolation of the trend is applied to estimate the emissions in the last five years of the inventory. Specifically, a linear regression model with autoregressive moving-average (ARMA) errors is used to estimate the trend in emissions over time from 1990 through 2012, and the trend is used to approximate the CH<sub>4</sub>, N<sub>2</sub>O, CO and NO<sub>x</sub> for the last 5 years in the time series from 2013 to 2017 (Brockwell and Davis 2016). The Tier 2 method described previously will be applied to recalculate the emissions in a future Inventory.

## Uncertainty and Time-Series Consistency

Emissions are estimated using a linear regression model with autoregressive moving-average (ARMA) errors for 2017. The linear regression ARMA model produced estimates of the upper and lower bounds to quantify uncertainty

(Table 5-34), and the results are summarized in Table 5-34. Methane emissions from field burning of agricultural residues in 2017 are between 0.10 and 0.29 MMT CO<sub>2</sub> Eq. at a 95 percent confidence level. This indicates a range of 51 percent below and 49 percent above the 2017 emission estimate of 0.2 MMT CO<sub>2</sub> Eq. Nitrous oxide emissions are between 0.04 and 0.11 MMT CO<sub>2</sub> Eq., or approximately 47 percent below and 46 percent above the 2017 emission estimate of 0.1 MMT CO<sub>2</sub> Eq.

**Table 5-34: Approach 2 Quantitative Uncertainty Estimates for CH<sub>4</sub> and N<sub>2</sub>O Emissions from Field Burning of Agricultural Residues (MMT CO<sub>2</sub> Eq. and Percent)**

Source	Gas	2017 Emission Estimate (MMT CO <sub>2</sub> Eq.)	Uncertainty Range Relative to Emission Estimate <sup>a</sup>			
			(MMT CO <sub>2</sub> Eq.)		(%)	
			Lower Bound	Upper Bound	Lower Bound	Upper Bound
Field Burning of Agricultural Residues	CH <sub>4</sub>	0.2	0.10	0.29	-51%	49%
Field Burning of Agricultural Residues	N <sub>2</sub> O	0.1	0.04	0.11	-47%	46%

Due to data limitations, there are additional uncertainties in agricultural residue burning, particularly the omission of burning associated with Kentucky bluegrass and “other crop” residues.

## QA/QC and Verification

A source-specific QA/QC plan for field burning of agricultural residues was implemented with Tier 1 analyses. Errors were identified and corrected in the analysis of the remote sensing product related to scaling of the data for the entire state for Iowa and Indiana, and also the application of the logical regression model.

## Recalculations Discussion

Methodological recalculations are associated with the new analysis to estimate the area burned based on the LANDFIRE data products (LANDFIRE 2014) and MODIS Global Fire Location Product (Giglio et al. 2006). The emissions decreased on average across the times by 43 percent and 28 percent for CH<sub>4</sub> and N<sub>2</sub>O, respectively. The new analysis is considered more robust with an evaluation of burned area across the entire time series using the remote sensing products, rather than only subset of years, which was used in the previous Inventory.

## Planned Improvements

The key planned improvement to estimate the emissions associated with field burning of agricultural residues in the states of Alaska and Hawaii. In addition a new method is in development that will directly link agricultural residue burning with the Tier 3 methods that are used in several other source categories, including Agricultural Soil Management, *Cropland Remaining Cropland*, and *Land Converted to Cropland* chapters of the Inventory. The method is based on the DAYCENT model, and burning events will be simulated directly within the process-based model framework using information derived from remote sensing fire products as described in the Methodology section. This improvement will lead to greater consistency in the methods for these sources, and better ensure mass balance of C and N in the Inventory analysis.