


Systematic Optimization of High Activity Neutralization Materials for Bulk Chemical Agent Detoxification



PRESENTED BY

Patrick D. Burton¹; Mark K. Kinnan¹; Calen J. Leverant¹
David J. McGarvey²; William R. Creasy³

1 Sandia National Laboratories

2 CCDC CBC

3 Leidos, supporting CCDC CBC



Sandia National Laboratories is a multimission laboratory managed and operated by National Technology & Engineering Solutions of Sandia, LLC, a wholly owned subsidiary of Honeywell International Inc., for the U.S. Department of Energy's National Nuclear Security Administration under contract DE-NA0003525.

SAND2019-13538 C

Detoxification is a fundamentally different process from decontamination



Several methods and chemistries have been developed to remediate chemical warfare agents or toxic industrial chemicals on a variety of surfaces.

All of them share a common set of assumptions and practices:

1. The agent is a minority contaminant in a comparatively much larger system.
 - Eliminate toxicity posed by the agent while leaving the system (relatively) intact.
2. The precise quantity and location of agent is unknown.
 - Excess decontamination reagent is needed to ensure effective decontamination.
3. Many CWAs are poorly soluble in aqueous solution
 - Surfactants or substantial volumetric excess increase contact of the decontaminant with hydrophobic agents.




These practices are not practical for a bulk detoxification scenario

Detoxification of bulk chemicals requires specific considerations



Chemical agent
comprises the
majority or entirety
of the system




A smaller volume of more
highly reactive reagents is
needed for practical
detoxification.


Since great excesses are
not used, physical mixing
and solubility of the
reagent becomes a more
significant issue.

Treatment with a substantial excess of
mild reagents is not feasible.

Under bulk conditions, the agent itself is the reaction solvent



As the reaction progresses, products accumulate and change the distribution of agent and reactants.



Products may passivate the reactant, limiting overall reaction

Achieving effective detoxification requires a highly efficient approach


The 12 Principles of Green Chemistry are guidelines to achieve the greatest product output while minimizing detrimental environmental impact.





Lump of mustard (HD) recovered by fishing in the Baltic Sea

Missiaen *et al.*, *Science of The Total Environment* 2010, 408 (17), 3536-3553.

- Prevention
- Atom Economy
- Less Hazardous Chemical Syntheses
- Designing Safer Chemicals*
- Safer Solvents and Auxiliaries
- Design for Energy Efficiency
- Use of Renewable Feedstocks
- Reduce Derivatives*
- Catalysis
- Design for Degradation
- Real-time analysis for Pollution Prevention
- Inherently Safer Chemistry for Accident Prevention












*This effort is to *decompose* existing materials into a less toxic form. The concepts are still valid, but are approached differently.

Synergistic benefits of Li_3N compared to LiOH and NH_3 







Agent	RXN Vol (mL)	Additive Quantity (% vol)	Agent Remaining with Time
GB	10	13	9% (1 day)
GD	10	13	<0.4% (5 days)
VX	10	13	14% (13 days)
VX	100	18	17% (5 days)
QL	10	18	ND (7 day)
DF	10	19	16% (7 days)










The logistics of transporting reagents to detoxify CWAs can be challenging. Making efficient use out of every atom becomes essential to minimize transportation burden.



9 Challenges specific to HD





HD hydrolyzes easily under optimal conditions, but is very insoluble in aqueous solution.



Tailoring the solvent used to initiate Li_3N reaction provides control over the reactants produced.

Larger aliphatic alcohols reduce the reaction rate.

Cooperative effect of high pH and ammonia production



Substoichiometric Li_3N is still reactive, but not as fast. Since there is not enough NH_3 to fully react, (slower) hydrolysis is likely taking place.

Requires a surfactant to disperse surrogate



- Detoxification of bulk scale CWAs ($\text{Volume}_{\text{CWA}} > \text{Volume}_{\text{neutralization reagent}}$) is challenging.
- The agent, neutralization reagent, and breakdown products must be collectively optimized.
- Applying green chemistry practices provides the needed efficiency perspective.
- Li_3N , a pyrophoric material, provides multifunctional detoxification efficacy.

Questions?

This research was funded and supported by the Defense Threat Reduction Agency (DTRA)/Joint Science and Technology Office (JSTO) under project CB10412. We thank Dr. Glenn Lawson of DTRA/HSTO for his support of this project and Dr. Kevin Morrissey of the Decontamination Sciences Branch for contract support. The views expressed in this presentation are those of the authors and do not necessarily reflect the official policy or position of the Department of Defense or the U.S. Government.