


Guidance on the Preparation of Clean Air Act Section 179B Demonstrations for Nonattainment Areas Affected by International Transport of Emissions



Guidance on the Preparation of Clean Air Act Section 179B Demonstrations for Nonattainment Areas Affected by International Transport of Emissions

> U.S. Environmental Protection Agency Office of Air Quality Planning and Standards Air Quality Policy Division Research Triangle Park, NC

TABLE OF CONTENTS

Acronyms		ii
1. INTRO	DUCTION	1
	pose	
	tutory and Regulatory Framework	
	pe and Definition of International Emissions	
	alytical Considerations	
	NATIONAL TRANSPORT OF POLLUTION	
	ar-border Transport	
	ng-range Transport	8
	JATING THE SECTION 179B RELATIONSHIP TO OTHER RELATED	10
	VSension of Attainment Date	
	ension of Attainment Dateeptional Events	
	tions 110(a)(2)(D) and 126 – Interstate Transport	
	EMONSTRATION DEVELOPMENT PROCESS UNDER SECTION 179B	
	ly Engagement with EPA Regional Offices	
	spective Demonstrations under Section 179B(a)	
4.2.1.	Section 179B(a) Process and Key Questions	
4.2.2.	Section 179B(a) Demonstration Submission Schedule	15
4.2.3.	Flow Chart for Section 179B(a) Prospective Demonstrations (SIP approval)	15
4.3. Ret	rospective Demonstrations under Sections 179B(b)–(d)	15
4.3.1.	Marginal Nonattainment Areas for the O ₃ NAAQS	16
4.3.2.	Schedule for Retrospective Demonstrations for O ₃ Nonattainment Areas	17
4.3.3.	Schedule for Retrospective Demonstrations for PM ₁₀ , PM _{2.5} , and CO	
Nonatta	inment Areas	17
4.3.4.	Flow Chart for Section 179B(b)–(d) Retrospective Demonstrations (potential	
	ication)	
	·	
	EPTUAL MODEL OF INTERNATIONAL INFLUENCE	
6. TECHN 23	ICAL ANALYSES IN SUPPORT OF THE SECTION 179B DEMONSTRAT	ION
_	comming the Ammoniste Analytic Ammond for the Section 170D Demonstra	tion 22
	termining the Appropriate Analytic Approach for the Section 179B Demonstrated Institute of the Section 179B Demonstrations	
	alyses to Demonstrate International Contribution	
6.3.1.	Measured Air Quality Data Analysis	
6.3.2.	Comprehensive Emission Analysis	
6.3.3.	Modeling to Quantify International Contribution	40
6.3.4.	Receptor Modeling Analysis	44
	ample Conclusion Statement in the Demonstration	
	Comment Process	
	ces	
Appendix A.	Statutory Text	52

Acronyms

AQS Air Quality System CAA Clean Air Act

CFR Code of Federal Regulations

CO Carbon monoxide

CTM Chemical Transport Model
ENSO El Nino / Southern Oscillation
EPA Environmental Protection Agency

FR Federal Register FT Free troposphere

HTAP Hemispheric Transport of Air Pollutants

HYSPLIT HYbrid Single-Particle Lagrangian Integrated Trajectory

K Kelvin km Kilometer

LRTAP The Convention on Long-range Transboundary Air Pollution

mb Millibar

NAAQS National ambient air quality standard or standards NASA National Aeronautics and Space Administration

NNSR Nonattainment New Source Review

NOAA National Oceanic and Atmospheric Administration

NO Nitric oxide NOx Nitrogen oxides

NPES Normalized Potential Emissions Sensitivity
NPSC Normalized Potential Source Contribution

O₃ Ozone

PA Policy Assessment

PBL Planetary boundary layer PES Potential Emissions Sensitivity

PM Particulate matter ppb Parts per billion

PSC Potential Source Contribution

PT Potential temperature

RACM Reasonably Available Control Measures
RACT Reasonably Available Control Technology

RFP Reasonable Further Progress SIP State Implementation Plan WAQS Western Air Quality Study

UN United Nations

1. <u>INTRODUCTION</u>

1.1. Purpose

The purpose of this guidance document¹ is to assist air agencies² that are considering the development of a demonstration, under section 179B of the Clean Air Act (CAA), that a nonattainment area would be able to attain and maintain, or would have attained, the relevant National Ambient Air Quality Standard (NAAQS) but for emissions emanating from outside the United States (U.S.). To help air agencies better understand how to satisfy the requirements of section 179B, the guidance describes and provides examples of the kinds of information and analyses that the U.S. Environmental Protection Agency (EPA) recommends air agencies consider including in a section 179B demonstration. The guidance also describes a weight-of-evidence approach that EPA intends to use when evaluating section 179B demonstrations. This non-binding guidance is intended to assist air agencies in the preparation of demonstrations but does not limit the types of information and analysis that could be used to develop such demonstrations under the CAA.

An air agency has the authority under section 179B to develop and submit to EPA a demonstration that its state implementation plan (SIP) would be adequate to attain and maintain the NAAQS in a designated nonattainment area, or that the nonattainment area would have attained the NAAQS, but for emissions emanating from outside the U.S. EPA has the authority under section 179B to assess such an international transport demonstration when evaluating a SIP revision submitted in response to a nonattainment designation or reclassification of an area, or when EPA determines whether a nonattainment area has failed to attain the standard by the attainment date and thus becomes subject to additional CAA requirements. If upon such assessment, the demonstration is to the Administrator's satisfaction, EPA will provide specified regulatory relief as indicated in section 179B.

In addition to describing the kinds of information and analyses that may be helpful to include in a section 179B demonstration, this guidance provides:

• A review of the existing regulatory framework for considering section 179B demonstrations;

¹ This document is intended only to provide clarity to the public regarding existing requirements under the law or agency policies. This document is not a rule or regulation, and the guidance it contains may not apply to a particular situation based upon the individual facts and circumstances. This guidance does not change or substitute for any law, regulation, or other legally binding requirement and is not legally enforceable. The use of non-mandatory language such as "guidance," "recommend," "may," "should," and "can" is intended to describe EPA's policies and recommendations. The use of mandatory terminology such as "must" and "required" is intended to describe controlling legal requirements under the terms of the CAA and of EPA regulations. Neither such language nor anything else in this document is intended to or does establish legally binding requirements in and of itself. None of the recommendations in this guidance are binding or enforceable against any person, and neither any part of the guidance nor the guidance as a whole constitutes final agency action that affects the rights and obligations of any person or represents the consummation of agency decision making. Only final actions taken to approve or disapprove state implementation plan (SIP) submissions or final findings by the Administrator under section 179B(b)–(d) that implement any of the recommendations in this guidance would be final actions for purposes of CAA section 307(b).

² References to "air agencies" include state, local, and tribal air agencies.

- A review of other existing regulatory mechanisms that may be more appropriate alternatives to section 179B in certain situations;
- Recommended timeframes for the section 179B demonstration development process; and
- Background on the nature of intracontinental and intercontinental transport of air pollution.

1.2. Statutory and Regulatory Framework

After EPA promulgates a new or revised NAAQS, the CAA requires EPA to designate all areas of the country as either Nonattainment, Attainment, or Unclassifiable, with respect to that NAAQS. The process for these initial area designations is outlined in CAA section 107(d)(1).

Under the CAA, air agencies are required to develop and submit SIPs to EPA that provide for the implementation, attainment, maintenance, and enforcement of the NAAOS through control programs directed at various sources of emissions. When designated as nonattainment, areas for the ozone (O_3) , particulate matter equal to or less than 2.5 microns in diameter $(PM_{2.5})$, particulate matter equal to or less than 10 microns in diameter (PM₁₀), and carbon monoxide (CO) NAAQS are each assigned a classification that identifies the latest allowable attainment date and associated requirements to be addressed in the SIP. The core statutory requirements³ to be addressed in the SIP may include the following: an accurate inventory of current emissions for all sources within the nonattainment area; a Nonattainment New Source Review (NNSR) permit program; regulations providing for implementation of Reasonably Available Control Measures (RACM), including Reasonably Available Control Technology (RACT); a demonstration that the plan provides for Reasonable Further Progress (RFP) toward attainment; a demonstration of attainment by the attainment date, commonly in the form of air quality modeling analyses; and contingency measures to be implemented in certain circumstances should the area fail to attain by the attainment date.⁴ These obligations are detailed in various CAA sections, including sections 110 and 171 through 193.

After the attainment date has passed, the air agency and EPA evaluate ambient air quality data for the nonattainment area to determine whether each area has attained by the attainment date. Once EPA has determined that an O₃, PM_{2.5}, PM₁₀, or CO nonattainment area has failed to attain the NAAQS by the attainment date, that area by operation of law must be reclassified to a higher classification, which will trigger additional planning obligations for the area, potentially resulting in further emissions control requirements.⁵

_

³ Specific statutory SIP due dates and requirements for SIPs vary depending on the specific NAAQS and an area's classification. Related implementation requirements and deadlines associated with these statutory requirements may be established by EPA regulations. The requirements listed here only refer to the core statutory requirements generally applicable to most nonattainment areas.

⁴ Note that O₃ nonattainment areas classified as Marginal are not subject to most of the requirements that apply to higher classifications. Compare CAA section 182(a) with sections 182(b)-(e).

⁵ Reclassification to a higher classification for failure to attain the NAAQS by the attainment date do not apply to areas that are already classified at the highest classification and, in the case of ozone, a classification of Severe (which is the second-highest level of classification for ozone). Although failure to attain the SO₂, NO₂, or Pb NAAQS does not result in reclassification, it does also result in additional planning and attainment demonstration requirements.

Section 179B⁶ provides EPA with authority to consider whether state demonstrations of the impacts from international emissions are satisfactory in two contexts: (1) at the implementation plan review stage, when EPA determines prospectively whether a SIP adequately demonstrates that a nonattainment area will attain the NAAQS by its future attainment date (in section 179B(a)); and (2) after each applicable attainment date, when EPA determines retrospectively whether a nonattainment area has attained the NAAQS by that attainment date (in section 179B(b)–(d)).

In the first context, section 179B(a) provides that, "[N]otwithstanding any other provision of law, an implementation plan or plan revision required under this chapter shall be approved by the Administrator if (1) such plan or revision meets all the requirements applicable to it...other than a requirement that such plan or revision demonstrate attainment and maintenance of the relevant national ambient air quality standards by the attainment date specified under the applicable provision of this chapter, or in a regulation promulgated under such provision, and (2) the submitting state establishes to the satisfaction of the Administrator that the implementation plan of such state would be adequate to attain and maintain the relevant national ambient air quality standards by the attainment date ... but for emissions emanating from outside of the United States." (Emphasis added). For the purpose of this guidance, we refer to such section 179B demonstrations as section 179B(a) or "prospective" demonstrations because they typically involve as assessment of future air quality. Thus, an EPA-approved prospective demonstration provides relief from demonstrating future attainment.

In the second context, sections 179B(b), (c), and (d) provide that, for O₃, CO, and PM, respectively, "[n]otwithstanding any other provision of law, any State that establishes to the satisfaction of the Administrator that ... such State would have attained the national ambient air quality standard ... by the applicable attainment date but for emissions emanating from outside of the United States" shall not be subject to reclassification to a higher classification category by operation of law, as otherwise required in CAA sections 181(b)(2), 186(b)(2), and 188(b)(2), respectively. (Emphasis added.) For the purpose of this guidance, we refer to such section 179B demonstrations as section 179B(b)—(d) or "retrospective" demonstrations because they involve analysis of past air quality. Thus, an EPA-approved retrospective demonstration provides relief from reclassification that would have resulted from EPA determining that the area failed to attain the NAAQS by the relevant attainment date.

While section 179B can be an important tool for providing specified regulatory relief⁸ for air agencies. EPA's approval of a section 179B demonstration does not relieve air agencies with


⁶ All references to CAA section 179B are to 42 U.S.C. § 7509a. International border areas, as added Pub. L. No. 101-549, title VIII, § 818, 104 Stat. 2697 (Nov. 15, 1990). See Appendix A for full statutory text.

⁷ EPA's longstanding view is that CAA section 179B(b) contains an erroneous reference to section 181(a)(2), and that Congress actually intended to refer here to section 181(b)(2). *See* "State Implementation Plans; General Preamble for the Implementation of Title I of the Clean Air Act Amendments of 1990," 57 Fed. Reg. 13,498, 13,569 n.41 (Apr. 16, 1992).

 $^{^8}$ The regulatory relief gained if the state's section 179B(a) prospective demonstration is to the satisfaction of the Administrator and approved by EPA is the ability for a state to submit an approvable attainment plan that does not demonstrate attainment and maintenance of the relevant NAAQS. If the state's section 179B(b) – (d) retrospective demonstration for an O₃, CO, PM_{2.5}, or PM₁₀ area that does not attain the standard by the attainment date is to the

nonattainment areas of having to meet the remaining applicable planning or emission reduction requirements in the CAA. It also does not provide a basis for either excluding air monitoring data influenced by international transport from regulatory determinations related to attainment and nonattainment, or redesignating an area to attainment.

If an air agency is contemplating a section 179B demonstration in either the section 179B(a) "prospective" context or the section 179B(b)–(d) "retrospective" context, EPA encourages early consultation and communication throughout the demonstration development and submission process, along the lines of these basic steps:



More detailed, context-specific recommended process diagrams can be found in Sections 4.2.3 and 4.3.4 of this guidance.

1.3. Scope and Definition of International Emissions

This guidance provides examples of recommended technical analyses that an air agency can consider demonstrating either that:

- 1) its SIP would be adequate to show attainment and maintenance of the NAAQS by the attainment date but for international emissions; or
- 2) the area would have attained the NAAQS by the area's attainment date but for international emissions.

EPA expects section 179B demonstrations to be developed in a manner consistent with the CAA principles and practices used in attainment plans. The overall plan requirements for nonattainment areas, as identified in sections 171-193 of Part D of the Act⁹, call for plans that: (i) include a comprehensive inventory of actual emissions from all sources; (ii) identify and quantify the allowable emissions from major new or modified stationary sources, and require permits for major stationary sources; (iii) provide for implementation of reasonably available control measures (including reasonably available control technology); (iv) provide for reasonable further progress toward attainment; (v) include emission limitations and such other control measures as may be necessary or appropriate to provide for attainment; (vi) a demonstration of

_

satisfaction of the Administrator and is approved by EPA, EPA will not reclassify the area to the next higher classification. A state with an approved section 179B(b) retrospective demonstration for a Severe or Extreme ozone nonattainment area that fails to attain the standard by the attainment date also will not be subject to CAA section 185 fees.

⁹ CAA sections 171-193, 42 U.S.C. §§ 7501-7515.

attainment by the attainment date, commonly in the form of air quality modeling analyses; and (vii) contingency measures to be implemented in certain circumstances should the area fail to attain by the attainment date. Each one of these mandatory planning requirements is linked to anthropogenic emissions. To promote consistency between the nonattainment planning requirements and the corresponding section 179B provisions for relief from certain elements of those requirements, EPA recommends that section 179B demonstrations focus on the contribution to ambient concentrations attributable to international *anthropogenic* emissions. Since states are required to evaluate and adopt controls on domestic anthropogenic sources as necessary to fulfill their nonattainment planning requirements. consistent treatment in section 179B demonstrations would focus on contributions from non-U.S. anthropogenic sources as opposed to nonanthropogenic sources.¹⁰

For purposes of this guidance, the terms "international sources" and "international emissions" therefore mean, respectively, anthropogenic sources located outside of the U.S, and anthropogenic emissions emanating from sources located outside of the U.S. Emissions from offshore areas where U.S. federal laws govern emission sources are not considered to be international emissions for purposes of this guidance.¹¹

1.4. Analytical Considerations

Despite the title of section 179B ("International Border Areas"), EPA has twice affirmed in recent years its interpretation that this provision is not restricted to areas adjoining international borders. ¹² As explained in these instances, and as further detailed later in this document, domestic O₃ air quality can be affected by sources of emissions located across U.S. borders in Canada and Mexico, and under certain circumstances, from sources in other continents. Additionally, in his April 12, 2018, Memorandum for the Administrator of the Environmental Protection Agency ("Promoting Domestic Manufacturing and Job Creation – Policies and Procedures Relating to Implementation of Air Quality Standards"), the President directed EPA to "not limit its consideration of demonstrations or petitions to those submitted by States located on the borders of the United States with Mexico or Canada, but rather consider[] section 179B demonstrations or petitions submitted by any State, including but not limited to those located in

¹⁰ EPA believes anthropogenic emissions should be the focus of section 179B demonstrations for the reasons stated here. A state's weight-of-evidence presentation in a section 179B demonstration may refer to nonanthropogenic emissions. For example, certain analyses described herein, e.g., back trajectories, may not be able to distinguish the source of emissions. Therefore, it may be useful to identify days with contributions primarily from natural sources to contrast with days where ambient concentrations may be enhanced by contributions from anthropogenic sources.

¹¹ For example, emissions within United States portion of the North American Emission Control Area, as is defined in Annex VI of MARPOL, are regulated under the Act to Prevent Pollution from Ships. 33 U.S.C. §§ 1905-1915; see 40 C.F.R. Part 1043. Annex VI of MARPOL is available at http://www.imo.org/en/OurWork/Environment/PollutionPrevention/AirPollution/Pages/Air-Pollution.aspx. Air emissions from sources located on the Outer Continental Shelf are also subject to federal regulation. 42 U.S.C. §

emissions from sources located on the Outer Continental Shelf are also subject to federal regulation. 42 U.S.C. § 328, 43 U.S.C. § 1334(a)(8). States considering offshore sources as part of their section 179B demonstration should consult with their EPA Regional office.

¹² Although section 179B is titled "International Border Areas," EPA indicated in the preambles to the implementation rules for both the 2015 and 2008 O₃ NAAQS that the statutory language of section 179B does not prohibit air agencies from submitting such demonstrations for nonattainment areas not located on the border with another country. 83 FR 63010 (December 6, 2018); see also 80 FR 12294 (March 6, 2015). To date, EPA has not acted on a section 179B demonstration for any nonattainment area not located on the Mexican border.

the Western United States." As explained in this document, technical demonstrations for non-border areas may involve additional technical rigor and resources compared to demonstrations for border areas.

This guidance provides examples of information and analyses that are recommended for inclusion in section 179B demonstrations. Air agencies may also use other well-documented, appropriately applied, and technically sound information and analyses not identified in this guidance. EPA recommends that the air agency consult with its EPA Regional office as early as possible to reach a common understanding of the types of information and analyses that would be most appropriate for a section 179B demonstration for a particular area. EPA is hopeful that early consultation will help air agencies develop high-quality technical analyses and enable EPA to conduct expeditious reviews of section 179B demonstrations.

EPA recommends that section 179B demonstrations include a conceptual model that describes the conditions causing the exceedance(s)¹³ at the ambient monitor(s), discusses how emissions from international anthropogenic sources led to the exceedances at the affected monitor(s), and identifies the specific EPA regulatory decision (SIP approval or determination of attainment) that is intended to be addressed by the demonstration. The conceptual model would generally appear at or near the beginning of a demonstration to help readers and the reviewing EPA staff understand the role of international transport before more detailed technical evidence is presented. It would include much of the information that the air agency provided or discussed with EPA during initial consultations regarding a possible demonstration. Section 4 of this guidance describes the recommended scope of a conceptual model in more detail.

As described in Section 1.2 of this guidance, CAA section 179B contemplates two different types of demonstrations that could be developed by an air agency at different points in the air quality management process. If the air agency intends to submit a section 179B(a) prospective demonstration (i.e., one intended to support approval of a SIP submission by showing that the plan would be adequate to *attain and maintain the standard by the attainment date* but for international emissions), it should submit the demonstration prior to or as part of the overall nonattainment area SIP submission. A retrospective demonstration pursuant to sections 179B(b)–(d) (i.e., one intended to avoid a reclassification by showing that an area *would have attained the standard* but for international emissions) should illustrate that air quality was influenced by international emissions on specific days during the years that contribute to the design value calculation for the area. This retrospective demonstration would be submitted after air quality data collected pursuant to federal reference or equivalent monitoring methods and indicating that the area failed to attain by the attainment date are certified. As noted in Section 4.3 of this guidance, the submittal of a retrospective demonstration in some cases could occur before the attainment date. ¹⁴

_

¹³ For purposes of this guidance, unless otherwise specified, the term "exceedance" also refers to non-exceedance-level concentrations that contribute to a design value that violates certain NAAQS (e.g., one year's 4th highest daily maximum 8-hour average value could be below the level of the O₃ NAAQS, but the 3-year design value is above the NAAQS).

¹⁴ Note that an area that has submitted a retrospective demonstration but that ultimately attains the NAAQS by the relevant attainment date would not be subject to reclassification and, therefore, relief offered under section 179B(b) – (d) would no longer be applicable.

Given the extensive number of technical factors and meteorological conditions that can affect international transport of air pollution, EPA believes that section 179B demonstrations should be evaluated based on the weight of evidence ¹⁵ of all information and analyses provided by the air agency. The appropriate level of supporting documentation will vary on a case-by-case basis depending on the nature and severity of international influence. EPA will consider and qualitatively weigh all evidence based on its relevance to section 179B and the nature of international contributions as described in the demonstration's conceptual model. Every demonstration should include fact-specific analyses tailored to the nonattainment area in question. When a section 179B demonstration shows that international contributions are larger than domestic contributions, the weight of evidence will be more compelling than if the demonstration shows domestic contributions exceeding international contributions. Section 6 includes additional discussion of weight of evidence considerations.

The demonstration should also consider and examine any contradictory evidence that may indicate that influences other than international emissions caused or contributed to the subject NAAQS exceedances or violations. Such evidence, for example, may include data indicating that NAAQS exceedances or violations could be predominantly attributed to local, intrastate, or interstate U.S. sources which may otherwise be addressed by other CAA authorities. EPA will weigh the body of available evidence to determine whether it collectively indicates that the SIP would be adequate to attain and maintain the NAAQS, or the area would have attained the NAAQS, but for emissions emanating from outside of the U.S.

2. INTERNATIONAL TRANSPORT OF POLLUTION

The inclusion of section 179B in the 1990 CAA Amendments was an acknowledgement of the long-standing recognition of international transport of pollution to the U.S. In the legislative history to the 1990 Amendments (US Senate, 1993), the Senate committee on Environment and Public Works recognized that EPA and air agencies would need to develop technical analyses that attempted to quantify the impact of international transport of pollution (US Senate, 1993, p. 5742 and 10110). EPA has actively engaged in various efforts since that time to quantify and understand the impacts to the U.S. of both nearby and more distant international emissions from around the world. This section provides an overview of the type of transport that occurs at intracontinental and intercontinental scales.

2.1. Near-border Transport

¹⁵ Throughout this guidance document, the term "weight of evidence" is used to describe the collective analysis by which we evaluate CAA section 179B demonstrations. The guidance describes certain recommended analyses and other supplemental analyses that can be provided as part of a section 179B demonstration. Because each nonattainment area is unique, area-specific factors may affect the types of analyses that would be appropriate for any particular area. If the state provides multiple analyses as part of a section 179B demonstration, EPA recommends the state describe the analyses performed, databases used, key assumptions and outcomes of each analysis, and why an air agency believes that the evidence, viewed as a whole, supports a conclusion that the area would not attain, or would have attained, but for international emissions. The EPA will consider the weight of this evidence in considering whether to approve any section 179B demonstration.

Near-border transport to the U.S. from Canada and Mexico is more easily observed and documented than intercontinental transport. Pollutants from near-border international emissions sources, such as industrial facilities and motor vehicles, are transported on a scale comparable to the distance extending across large metropolitan areas. For example, international near-border emissions from Juarez, Mexico, have been demonstrated to have an impact on El Paso, Texas, located directly across the border (TNRCC, 1994; EPA, 2009). Conceptually, analyses for near-border areas can be similar to analyses performed to assess interstate transport under separate CAA authorities and requirements. ¹⁶ In addition, North American intracontinental transport can also occur over greater distances, and it can potentially affect interior locations in the U.S. to a lesser degree than border areas.

EPA has previously estimated the level of O₃ and haze pollution contributed to U.S. air quality monitors from near-border Canadian sources, Mexican sources, and international shipping emissions from outside offshore areas governed by federal law. For example, air quality modeling for the 2015 O₃ NAAQS Policy Assessment (PA) estimated Canadian and Mexican contributions to U.S. pollution for the calendar year 2007 (EPA, 2014). In addition, modeling to support the Cross-State Air Pollution Rule and the Regional Haze Rule quantified projected 2017 O₃ contributions and 2028 particulate matter contributions from an approximately 300,000 square mile region of northern Mexico and an approximately 600,000 square mile region of southern Canada, respectively (EPA, 2016a; EPA, 2018b). These two studies use two techniques (i.e.: (1) "zero-out"; and (2) source apportionment, both discussed in more detail in Section 6.3.3.1 of this guidance) that provide quantitative estimates of international transport.

2.2. Long-range Transport

Long-range international transport of air pollution has been recognized for decades, as illustrated by Byrne (2015). Particularly with O₃, modeling analyses have typically included boundary conditions that implicitly included "background" O₃. "Background" is a generic term that has been used to refer to O₃ formed from any non-local or regional source (EPA, 2015), but recent publications have refined the term to focus on uncontrollable sources (Jaffe et al., 2018). "Background" or "Uncontrollable sources," as used in discussions of domestic air pollution, include natural ¹⁷ sources globally (i.e., domestic and international), plus international anthropogenic sources. EPA has been aware of global natural and international anthropogenic contributions while setting prior O₃ NAAQS, as illustrated by the use of boundary conditions in modeling. ¹⁸

The characterization of international transport of pollution has significantly advanced in the last decade with the development of improved international emissions inventories and global-scale chemical transport models (CTMs). While global scale CTM simulations alone have been used to estimate international transport to the U.S., global-scale models typically have coarse spatial resolution and perform poorly for estimating concentrations in U.S. urban areas (Jaffe, et al.,

¹⁷ Natural sources are inclusive of all non-anthropogenic sources, which include biogenic, geogenic, oceanic etc. Some natural sources have complex relationships with human activity, such that some anthropogenic enhancement of natural sources will be considered anthropogenic.

¹⁶ For example, CAA sections 110(a)(2)(D) and 126.

¹⁸ Boundary conditions accounted for all sources that were not explicitly accounted for within the modeling domain.

2018). To better estimate impacts of international transport of emissions to U.S. urban areas, recent modeling studies have used higher spatial resolution models of the North American "region" with boundary conditions derived from global-scale CTMs.

EPA has also endeavored to characterize and quantify contributions from more distant international sources. EPA has been actively engaged in the task force on Hemispheric Transport of Air Pollutants (HTAP) as part of the Convention on Long-range Transboundary Air Pollution (LRTAP). The HTAP effort has included two phases that focused on characterizing transported air pollution for the years 2000 (Dentener, Keating, & Akimoto, 2010) and 2010. These studies quantify the sensitivity of air pollution levels in the U.S. to anthropogenic emissions from other regions, with an emphasis on long-range transport from Europe, Russia, and Asia. EPA also supported the National Research Council's 2010 report "Global Sources of Local Pollution: An Assessment of Long-Range Transport of Key Air Pollutants to and from the United States." (NRC, 2010)

Synthesis of the literature, including citations above and other studies, show that contributions from international emissions to U.S. O₃ concentrations come from a combination of diffuse background and identifiable plumes from intercontinental transport. Intercontinental transport plumes occur more efficiently between certain locations on the globe than others (Dentener, Keating, & Akimoto, 2010; NRC, 2010). For example, semi-permanent pressure systems can set up an atmospheric "conveyor belt" between the Asian east coast and the U.S. west coast. Off the coast of China, a semi-permanent low-pressure system lofts air and associated pollutants to the middle and upper free troposphere. In this region of the atmosphere, fast winds can move pollution eastward toward the U.S. Pacific coast over the course of days to weeks. During transport in the upper troposphere, O₃ has a long chemical lifetime because the low temperatures in that part of the atmosphere are not conducive to O₃ destruction.

The U.S. Pacific coast has a semi-permanent high-pressure system that draws down air from the upper troposphere. During vertical mixing of air from the upper troposphere with air closer to the surface, O₃ transported in the upper troposphere is diluted and chemically destroyed (NRC, 2010). The amount of international O₃ that is mixed down from the upper troposphere with air near the surface depends on location and local meteorology. Generally, the further a location is from an international source the less O₃ will be available to mix down due to dispersion and chemical lifetime. Local meteorology, however, creates exceptions to this rule due to the dynamics of the planetary boundary layer (PBL), which is the lowest layer of the troposphere, and topography. Areas with exceptionally deep PBLs can more rapidly transport free tropospheric air to the surface. Complex topography can include mountain peaks that are routinely exposed to free tropospheric air. The peaks can also create winds that enhance O₃ mixing down mountain slopes (Zaveri, Saylor, Peters, McNider, & Song, 1995). As a result, high-altitude locations with complex topography may experience greater impacts from intercontinental transport of O₃ as compared to locations at lower elevations.

There is a large body of research exploring the role of international transport and its evolution. For example, it is well known that Asian NOx emissions increased quickly in the 1990s and 2000s (van der A, et al., 2017) and that transport of O₃ also increased (Lin, Horowitz, Payton, Fiore, & Tonnesen, 2017; Verstraeten, et al., 2015). Several recent studies (Huang, et al., 2017;

Nopmongcol, Liu, Stoeckenius, & Yarwood, 2017) have quantified interhemispheric transport in 2008 or 2010 near the peak contribution (circa 2011). There is broad consistency in the literature that transpacific transport increased and has since either flattened or decreased (Jaffe, et al., 2018).

3. EVALUATING THE SECTION 179B RELATIONSHIP TO OTHER RELATED PROVISIONS

As discussed in this document, section 179B addresses sources of emissions originating outside of the U.S. and provides qualifying nonattainment areas with specified regulatory relief from certain otherwise applicable planning and emissions control requirements. This section discusses limitations to section 179B applicability and identifies related regulatory mechanisms that air agencies may also find useful for addressing their obligations.

Section 179B applies only to nonattainment areas, and only provides relief from (1) the attainment demonstration requirement and (2) the requirement that EPA determine whether an area failed to attain by the attainment date (and associated reclassification as appropriate). It does not provide EPA with the authority to do any of the following:

- Exclude monitoring data influenced by international transport from regulatory determinations related to an area's designation as attainment or nonattainment (however, if an exceedance or violation is event-related, it may be able to qualify as an exceptional event, as described in Section 3.2 of this guidance);
- Classify an area with a lower classification than indicated by actual air quality;
- Relax any mandatory control measures associated with the area's classification;
- Redesignate a nonattainment area to attainment without meeting the other attainment plan requirements of CAA section 107(d)(3); or
- Determine that a state satisfied its interstate transport SIP obligations under CAA section 110(a)(2)(D)(i), the "good neighbor" provision.

Air agencies should also consider whether the regulatory mechanisms discussed below would be a more appropriate mechanism before initiating a section 179B demonstration. Air agencies with questions regarding these separate mechanisms are encouraged to consult their EPA Regional office.

3.1. Extension of Attainment Date

A nonattainment area that fails to attain the O₃, CO, PM_{2.5}, or PM₁₀ NAAQS by its attainment date but has achieved a threshold level of air quality and has met all requirements and commitments pertaining to the area in the applicable implementation plan may be eligible for a 1-year extension of the attainment date. For example, an area that fails to attain the 2015 O₃ NAAQS by its attainment date may be eligible to request a 1-year extension if it has complied

with all requirements in its SIP and if, for the attainment year, the area's fourth highest daily maximum 8-hour average is at or below the level of the standard. ¹⁹ A 1-year extension is also possible for nonattainment areas for CO, $PM_{2.5}$, and PM_{10} . ²⁰ In addition, up to 5-year extensions are also possible under certain conditions for $PM_{2.5}$ and PM_{10} areas classified as Serious. ²¹

As an attainment date approaches, an air agency may consider whether to develop a retrospective section 179B demonstration while waiting to see whether certified air quality data will ultimately qualify the area for an attainment date extension. For an area with a high likelihood of attaining a NAAQS with the additional time provided by an extension, the relief provided by section 179B may be unnecessary. However, due to the lead time to prepare a retrospective section 179B demonstration, the air agency should consider beginning development of a retrospective demonstration well before the attainment year air quality data are certified, in the event that the area does not ultimately qualify for the extension.

3.2. Exceptional Events

Section 319(b) of the CAA recognizes that, when making certain NAAQS-related regulatory determinations, including determinations of attainment by the attainment date, it may be appropriate to exclude ambient monitoring data that are influenced by exceptional events. The 2016 Exceptional Events Rule provides the regulatory mechanism for this purpose.²²

When exceptional events influence monitoring data and cause exceedances or violations of the NAAQS, air agencies can develop and submit a technical demonstration to request the exclusion of certain event-influenced data. If the demonstration satisfies the Exceptional Events Rule criteria, EPA can exclude these data from the data set used for certain regulatory decisions. Specifically, transported pollution, which may include pollution from outside the U.S., must be event-related and be either natural or caused by a human activity unlikely to recur at a particular location (see 40 CFR 50.14(c)(3)(iv)(E)).

Exceptional events from natural causes may include wildfires, stratospheric O₃ intrusions, high wind dust events, and volcanic activity. Exceptional events that are human activities unlikely to recur at a particular location may include prescribed fires on wildland, chemical spills, industrial accidents, or terrorist activities. In accordance with the CAA, routine emissions generated by and transported from anthropogenic sources, foreign or domestic, are not exceptional events.²³

3.3. Sections 110(a)(2)(D) and 126 – Interstate Transport

²² 81 FR 68216 (October 3, 2016).

¹⁹ See CAA section 181(a)(5). More information regarding extending attainment dates for the O₃ NAAQS is available in Implementation of the 2015 National Ambient Air Quality Standards for Ozone: Nonattainment Area State Implementation Plan Requirements, 83 FR 62998 (December 6, 2018).

²⁰ See CAA section 186(a)(4) for CO, and CAA section 188(d) for Moderate PM_{2.5} and PM₁₀ areas.

²¹ See CAA section 188(e).

²³ An example of routine emissions generated by and transported from anthropogenic sources might include emissions of O₃ precursors or directly emitted particulate matter (or PM precursors) from one state or foreign country's power plants transported into another state or the U.S.

For some nonattainment areas, the contribution to poor air quality from interstate pollution transport may be more substantial than contributions from international sources. The "good neighbor" provisions in CAA sections 110(a)(2)(D) and 126 may apply in addressing out of state pollution sources that may be interfering with attainment and maintenance of a NAAQS. CAA section 110(a)(2)(D) requires all states, to have SIPs containing adequate provisions prohibiting any source or other type of emissions activity within the state from emitting any air pollutant in amounts that will contribute significantly to nonattainment in, or interfere with maintenance by, any other state with respect to any NAAQS. CAA section 126 provides a mechanism for states to petition EPA for a finding that any out-of-state major source or group of stationary sources emits or would emit any air pollutant in violation of the prohibition of section 110(a)(2)(D).

EPA notes that an upwind state that is considering or has an approved section 179B demonstration must still meet its obligations for contributions to downwind states under CAA section 110(a)(2)(D). Likewise, an upwind state that contributes to a downwind state that is considering or has an approved section 179B demonstration must still meet its obligations for contributions to that downwind state, per CAA section 110(a)(2)(D). Policy governing whether an air agency can take into account emissions emanating from outside the U.S. in developing a plan to meet its interstate transport obligation²⁴ for any NAAQS is outside the scope of this guidance.

4. <u>THE DEMONSTRATION DEVELOPMENT PROCESS UNDER SECTION 179B</u>

This section describes the recommended section 179B demonstration development and submission process, including additional details about prospective and retrospective demonstrations. The Clean Air Act timelines for consequential SIP obligations and regulatory determinations associated with nonattainment areas give rise to different submission schedules for each type of demonstration.

4.1. Early Engagement with EPA Regional Offices

EPA recommends that an air agency engage with its EPA Regional office as early as possible when considering development of a demonstration under section 179B. The air agency and EPA should discuss the conceptual model for characterizing the international impacts, identify the types of analyses that would be most appropriate for the demonstration, and establish expectations for timing and other considerations. An air agency also may choose to share an early engagement draft demonstration with its EPA Regional office for review and feedback prior to submitting a final demonstration to EPA. This early engagement draft may include preliminary air quality data but a final retrospective demonstration submitted for EPA action should rely on final certified air quality data including the attainment year as further explained in section 4.3 of this guidance.

4.2. Prospective Demonstrations under Section 179B(a)

_

²⁴ See CAA section 110(a)(2)(D)(i) (aka the "good neighbor" provision) for SIP requirements addressing interstate transport.

As discussed in Section 1.2 of this document, section 179B(a) states that a *required* implementation plan or plan revision shall be approved by the Administrator if such plan or revision meets all the requirements applicable to it, other than a requirement that such plan or revision demonstrate attainment and maintenance of the relevant NAAQS by the attainment date, if the air agency establishes to the satisfaction of the Administrator that the SIP *would be adequate to attain and maintain* the relevant NAAQS by the applicable attainment date, but for emissions emanating from outside of the U.S. The wording in paragraph (a) of section 179B (the air agency must show that the SIP "would be" adequate to attain and maintain the NAAQS but for international emissions) indicates that this demonstration is to be forward-looking or prospective in nature. For this reason, we refer to such demonstrations as "prospective demonstrations" under section 179B(a).

Section 179B(a) does not include language specific to only certain NAAQS, and thus an air agency theoretically could develop such a demonstration for any NAAQS. If an air agency is required to submit a SIP demonstrating attainment of the NAAQS by the relevant attainment date and is seeking to submit a SIP that shows that the area would attain and maintain the NAAQS by the attainment date but for international emissions, then the air agency should submit a demonstration under the process and schedule described in this section.

With regard to O₃ nonattainment areas classified Marginal, ²⁵ a prospective attainment demonstration is not a *required* SIP obligation for a Marginal area, and therefore there is no prospective relief to be granted by EPA under section 179B(a). Accordingly, section 179B(a) prospective demonstrations are not relevant for Marginal O₃ areas, and EPA does not intend to evaluate requests for prospective relief in these instances.

4.2.1. Section 179B(a) Process and Key Questions

A SIP that includes a prospective demonstration under section 179B(a) must meet all requirements of the CAA Chapter applicable to it other than a demonstration that the area will attain the NAAQS by the attainment date. ²⁶ In general, the applicable requirements for any nonattainment area SIP include an accurate inventory of current emissions for all sources within the nonattainment area; a Nonattainment New Source Review (NNSR) permit program; regulations providing for implementation of Reasonably Available Control Measures (RACM), including Reasonably Available Control Technology (RACT); a demonstration that the plan provides for Reasonable Further Progress (RFP) toward attainment; a demonstration of attainment by the attainment date, commonly in the form of air quality modeling analyses; and contingency measures to be implemented in certain circumstances should the area fail to attain by the attainment date. Marginal O₃ nonattainment areas are the exception: applicable requirements are limited to an emissions inventory, source emission statements, transportation and general conformity programs, and a nonattainment new source review program.

-

²⁶ See CAA section 179B(a).

Requirements stated elsewhere in the CAA chapter 85 (Air Pollution Prevention and Control) but that are not applicable to specific nonattainment area SIP submissions include infrastructure SIP requirements such as the interstate transport program under section 110(a)(2)(D), which as indicated above are referred to as "good neighbor" SIPs. These "good neighbor" SIP obligations apply to *all* states and are due 3 years after a NAAQS is newly established or revised.

Section 179B(a)(2) defines the demonstration in terms of attainment "but for" international emissions. If the plan, in meeting all the otherwise applicable requirements of the CAA, results in attainment, it would be attaining "despite" international emissions rather than "but for" international emissions. Thus, to support the 179B(a) demonstration, an air agency should show that even after fulfilling other relevant obligations, such as imposition of required controls, the area is still projected not to attain the NAAQS. The air agency should therefore first evaluate whether the area can attain the standard by the attainment date based on required domestic emission reductions only.

An air agency should follow the steps below when considering and developing a demonstration under CAA section 179B(a). Step 1 is a required exercise as part of the attainment SIP development process. Step 2 is conditional upon the outcome of the analyses for Step 1.

Step 1. SIP Attainment Modeling for Potential Domestic Reductions: For the subject nonattainment area in the relevant SIP, the air agency should first model projected air quality concentrations for the attainment year based on on-the-books domestic net emission reductions and growth by the outermost attainment date (e.g., including mobile source standards, interstate transport programs, rules already adopted by the state for other nonattainment plans, etc.), and potential reductions associated with required controls for that particular NAAQS and classification that can be implemented by the attainment date.^{27, 28}

Go to Step 2 *only if* the Step 1 modeling shows the area could not attain with on-the-books measures and potential reductions associated with required controls for that particular NAAQS and classification (e.g., RACM/RACT) that can be implemented by the attainment date.

Step 2. Section 179B(a) Analyses for International Anthropogenic Emissions: The air agency may consider developing a demonstration of the impact of international emissions using the analyses identified in Section 6 of this document.

An air agency should include documentation of completing the analyses for Step 1 as part of its section 179B(a) demonstration.


²⁷ For further information on how to conduct these analyses, see Modeling Guidance for Demonstrating Air Quality Goals for Ozone, PM_{2.5}, and Regional Haze, November 29, 2018. Available at: https://www3.epa.gov/ttn/scram/guidance/guide/O3-PM-RH-Modeling Guidance-2018.pdf.

²⁸ Note that details on the RACM/RACT analytical process for a particular NAAQS can be found in other implementation rules and guidance. See, e.g., PM_{2.5} NAAQS implementation rule at 81 FR 58129-58132; O₃ NAAQS implementation rules at 70 FR 71659, 80 FR 12264, and 83 FR 62998; General Preamble for Implementation of Title I of the Clean Air Act Amendments of 1990 at 57 FR 13498.

4.2.2. Section 179B(a) Demonstration Submission Schedule

EPA considers a section 179B(a) demonstration as part of an attainment SIP. Therefore, the demonstration should be submitted to EPA no later than the date when all other SIP elements are submitted. An air agency may also elect to submit a section 179B(a) demonstration before the bulk of the attainment SIP is completed. EPA encourages air agencies to consult with their EPA Regional office to discuss timing associated with development and submission of a section 179B(a) demonstration to avoid potentially missing CAA deadlines.

4.2.3. Flow Chart for Section 179B(a) Prospective Demonstrations (SIP approval)



4.3. Retrospective Demonstrations under Sections 179B(b)–(d)

As discussed in Section 1 of this guidance, sections 179B(b)–(d) (relating to nonattainment areas for O₃, CO, and PM, respectively) provide that "any State that establishes to the satisfaction of the Administrator that such State would have attained the national ambient air quality standard by the applicable attainment date, but for emissions emanating from outside of the United States, shall not be subject to a mandatory reclassification to the next highest

classification by operation of law."²⁹ Because the wording in each of these section 179B paragraphs is presented in the past tense, EPA believes that such demonstrations should be retrospective in nature. In other words, the demonstration should include analyses showing that the air quality data on specific days in the past were affected by international emissions to an extent that prevented the area from attaining the standard by the attainment date.

A nonattainment area that may be eligible to submit a section 179B(b)–(d) retrospective demonstration would have previously been required under the CAA to submit an attainment SIP. The CAA requires an air agency to meet all SIP requirements applicable to the area during the SIP development, submission, and implementation process. Section 179B does not relieve an air agency of its planning or control obligations.

EPA acknowledges that for some NAAQS the timeline is challenging for air agencies to submit retrospective demonstrations in advance of the CAA deadline for EPA to determine whether an area attained by the attainment date. This is one reason why EPA recommends that the air agency communicate with the EPA Regional office as soon as possible after determining that it would like to explore the possibility of submitting a section 179B(b)—(d) retrospective demonstration. Much of the relevant air quality data and other information for a retrospective demonstration may be available to the air agency in advance of the attainment date. However, to ensure the integrity of air quality data used in demonstrations, EPA recommends that the air agency submit a final retrospective demonstration only after all air quality data used to calculate the attainment year design value are certified, but before the date by which EPA is required to make determinations of whether areas attained by the attainment date (i.e., 6 months after the attainment date). As further described in Section 4.3.2 of this guidance, an air agency may optionally pursue early certification of ambient data to help ensure timely submittal of a final retrospective demonstration prior to the date by which EPA must make its determinations of attainment. Provided that the attainment year air quality data are certified, in some cases air agencies may be able to submit a retrospective demonstration prior to the relevant attainment date.

As noted above, if EPA approves a section 179B(b)–(d) retrospective demonstration, EPA will not be required to issue a determination regarding whether the area attained by the attainment date pursuant to sections 181(b)(2), 186(b)(2), or 188(b)(2) of the CAA. Additionally, EPA does not intend to take action on any retrospective section 179B demonstrations until after all air quality data used to calculate the attainment year design value as of the attainment date are certified.

4.3.1. Marginal Nonattainment Areas for the O₃ NAAQS

As discussed in Section 4.2 of this guidance, the list of applicable requirements for O₃ nonattainment areas classified as Marginal is limited to: emissions inventory, source emission statements, transportation and general conformity programs, and a nonattainment new source review program. As described in the 2015 O₃ NAAQS Implementation Rule, section 182(a) of the CAA does not require states to implement RACM/RACT in Marginal O₃ nonattainment

16

²⁹ See CAA sections 179B(b), (c), and (d). Any such state (or area) would also not be subject to a determination of failure to attain by the attainment date.

areas, and nothing in section 179B alters the statutory requirements with respect to RACM/RACT obligations in subpart 2.³⁰

However, an air agency with a Marginal O₃ nonattainment area that is affected by international emissions may wish to evaluate whether implementing emission reduction measures on domestic sources in the nonattainment area can bring the area into attainment prior to the attainment date. If an air agency submits a section 179B(b) demonstration that is approved, an area with air quality violating a NAAQS will remain designated nonattainment and retain its current classification, until such time as the area attains the standard and the air agency submits, and EPA approves, a request for redesignation. The area will continue to be subject to NNSR and the other requirements noted above until the area meets and EPA approves an air agency submission addressing the redesignation criteria of CAA section 107(d)(3)(E).

4.3.2. Schedule for Retrospective Demonstrations for O₃ Nonattainment Areas

Although EPA encourages an air agency to consult with its EPA Regional office regarding a possible section 179B retrospective demonstration as soon as practicable, a complete section 179B retrospective demonstration should rely on certified air quality data for the entire period evaluated. EPA recommends that the air agency submit such a demonstration for O₃ as soon as possible after the data from the attainment year are certified.

Air agencies are required to certify O₃ air quality data for a given calendar year by May 1 of the following year. An air agency may also elect to pursue early certification of the preceding year's air quality data, which could improve the possibility of the air agency submitting a complete a section 179B retrospective demonstration in advance of the area's attainment date. Attainment dates for Marginal through Extreme O₃ NAAQS nonattainment areas extend from 3 to 20 years from the effective date of area designations by EPA. For the nonattainment areas for the 1997, 2008, and 2015 O₃ NAAQS, attainment dates fall on June 15, July 20, and August 3 (respectively) of various years, depending on an area's classification. EPA is obligated to make determinations of attainment within 6 months of each attainment date based on the area's design value (CAA section 181(b)(2)(A)). An area's design value is based on air quality data for the 3 calendar years preceding the attainment date. Therefore, EPA recommends that an air agency submit its section 179B retrospective demonstration as soon as practicable, and well ahead of EPA's deadline to make a determination of attainment, because EPA approval of such a demonstration would relieve the area from being subject to EPA's determination of whether the area attained by its attainment date.

4.3.3. Schedule for Retrospective Demonstrations for PM₁₀, PM_{2.5}, and CO Nonattainment Areas

Although EPA encourages the air agency to consult with its EPA Regional office regarding a possible section 179B retrospective demonstration as soon as practicable, a complete section 179B retrospective demonstration should rely on certified air quality data for the entire period evaluated. Thus, like the recommendation for O₃, EPA recommends that the state submit such a

_


³⁰ 83 FR 63010 (December 6, 2018).

³¹ 40 CFR § 58.15 - Annual air monitoring data certification.

demonstration for PM_{10} , $PM_{2.5}$, and CO as soon as possible after the data from the most recent year are certified.

Air agencies are required to certify air quality data for a given calendar year by May 1 of the following year. An air agency may also elect to pursue early certification of the preceding year's air quality data, which could improve the possibility of the air agency submitting a complete a section 179B retrospective demonstration in advance of the area's attainment date. The attainment date for PM₁₀, PM_{2.5}, and CO NAAQS nonattainment areas typically falls on December 31 of a given year (although there are some exceptions). EPA is required to make determinations of attainment within 6 months of the attainment date (i.e., by June 30th of the year following the December 31 attainment date). See CAA section 186(b)(2)(A) for carbon monoxide; CAA section 188(b)(2) for PM_{2.5} and PM₁₀. Therefore, EPA recommends that the air agency submit its section 179B retrospective demonstration as soon as practicable, and well ahead of EPA's deadline to make a determination of attainment, because EPA approval of such a demonstration would relieve the area from being subject to EPA's determination of whether the area attained by its attainment date.

4.3.4. Flow Chart for Section 179B(b)–(d) Retrospective Demonstrations (potential reclassification)



5. CONCEPTUAL MODEL OF INTERNATIONAL INFLUENCE

Section 179B demonstrations should include a conceptual model that is intended to frame the "state of the knowledge" for air quality in the nonattainment area. The conceptual model includes information regarding the influence of emissions, meteorology, transport, and/or other relevant atmospheric processes on air quality in given nonattainment area (EPA, 2018c). A well-constructed conceptual model of pollutant formation and transport for the area can assist in the

determination of international transport impacts by highlighting the contrast between locally formed pollutant days and the internationally influenced days in question. The conceptual model should provide the context for reviewing the more detailed analyses provided by the state, described in Section 3 of this guidance.

The conceptual model should clearly identify which type of transport and international sources/source regions directly affect the nonattainment area and whether that transport type is distinct from locally influenced high pollution conditions. Clearly identifying the conceptual distinction between local and international contributions forms the basis for which analyses should be included in the demonstration. The conceptual model should also identify which regions and sources meaningfully contribute to the international portion of emissions that influence ambient concentrations in the area of interest.

The information included in a conceptual model should help establish the context for the overall analysis and should be consistent with more detailed evidence and analyses provided in the overall demonstration³². To promote a shared understanding and interpretation of this information, the list below provides examples of the kinds of information that would typically be useful to include in a conceptual model.

- A summary of the affected area's NAAQS attainment and classification information.
- A description of the regulatory determination the state believes is influenced by the international emissions.
- A map of the existing ambient monitors in the area, and a description of the sites (e.g., site ID, current measured design value, elevation, recent pollutant trends), and any other relevant information.
- A list of the monitor(s) and days that the air agency has identified as influenced by international anthropogenic emissions.
- A description of the key differences between the measured exceedances influenced by international emissions concentrations and typical exceedances influenced by local, non-international emissions. It would be helpful to include a table of the relevant monitor data (e.g., date, hours, monitor values, and design value calculations with and without the international emissions).
- A summary of the meteorological and atmospheric conditions that lead to high concentrations at the monitor on days <u>influenced by international anthropogenic emissions</u> and days <u>not influenced by international anthropogenic emissions</u>. The contents of this summary will vary by area, but could include:
 - o the months in which high concentration days usually occur
 - o the diurnal evolution of a typical exceedance in the area
 - o the source sector(s) that contribute to the local and regional contribution
 - o the source sector(s) that contribute to the international contribution (if known)
 - o for PM, the pollutant species that contribute to typical exceedances in the area
 - o typical spatial patterns of exceedance days

_

³² All comments received on this guidance can be found at: https://www.regulations.gov/docket?D=EPA-HQ-OAR-2019-0668.

- o the meteorological conditions associated with high concentration days influenced by international emissions, including a description of the route traveled by transported pollution, such as distance and altitude
- o the meteorological conditions associated with high concentration days not influenced by international emissions.
- Identification of specific international anthropogenic emissions sources (e.g., an international emitting facility) or source regions (e.g., an international metropolitan area) that predominantly impact the monitor location on internationally influenced days
- Where available, a description of how controls on the upwind international anthropogenic sources differ from those required within the U.S. and how that difference could have affected the regulatory determination.
- A synthesis of literature characterizing international transport contributions to the geographic region of interest.

The California Air Resources Board's 2013 SIP for Imperial County for the 2006 24-Hour PM_{2.5} standard (CARB, 2014)³³ contains useful examples for several components of a conceptual model for a section 179B demonstration.

Figure 1 below, taken from the referenced Imperial County SIP, shows one example of an overview map where the border with Mexico is clearly visible, as are major topographical features, major roads, and cities. In addition, the lower left-hand inset presents the broader context of the nonattainment area location within California. This type of graphic should also show where area monitors are located relative to key international anthropogenic sources and provide context for other geographical considerations.

-

³³ The citation of specific sections of a plan or demonstration as useful examples is not intended as an endorsement of the entire document.




Figure 1. Example overview map, Imperial County $PM_{2.5}$ nonattainment area overview map $(CARB, 2014)^{34}$

That same SIP also included a graphic that connected time and space trends for PM_{2.5} concentrations. Figure 2 below, also taken from the referenced Imperial County SIP, indicates both that substantial air quality improvements have been made in the region and that the southern-most monitor (Calexico) has the highest design values with the smallest reduction between 2001 and 2012. This result would need to be corroborated with discussion of emissions in a later section to establish that domestic emissions affecting Calexico have been reduced commensurate with domestic emissions affecting El Centro and Brawley.

Other components of the conceptual model will be case-specific. In addition to the Imperial County demonstration for the 2006 PM_{2.5} NAAQS referenced in Figures 1 and 2, components of other section 179B demonstrations referenced in this document may provide helpful examples for how elements can be addressed.³⁵ Each situation regarding international contributions is unique and every section 179B demonstration will necessitate information and analyses to be tailored accordingly.

³⁵ This guidance includes these illustrative examples for the benefit of air agencies developing section 179B demonstrations. Most of these examples were taken from demonstrations that have been submitted to EPA. As previously mentioned, EPA has not yet acted on a demonstration for an area not adjoining an international border. As additional demonstrations are submitted to EPA, more illustrative examples may become available.

³⁴ The citation of specific sections of a plan or demonstration as useful examples is not intended as an endorsement of the entire document.




Figure 2. 2001 and 2012 24-hour Design Values for Brawley, El Centro and Calexico

Figure 2. 2001 and 2012 24-hour $PM_{2.5}$ Design Values for three Monitoring Sites in Imperial County, California (CARB, 2014, pp. 15, Figure 2.5)³⁶

Identification of sources and how controls could affect attainment provides context to the reviewer and informs potential international engagement. When international sources are uncontrolled or under-controlled, this helps a demonstration reviewer understand why the source is expected to have a large contribution. If potential controls of international sources could lead to attainment, this information may highlight an area for EPA international engagement and/or negotiation.

Finally, the conceptual model should be put in the context of historical and contemporary literature. There have been multiple projects and analyses completed that focus on U.S. background O₃ or on international transport that may be relevant. The results from these projects are often provided as annual averages or in metrics that are not specific to the NAAQS form, but they still provide important insights and establish expectations. For example, a literature synthesis should summarize international transport magnitudes, variability, and seasonality. The synthesis should highlight where modeling results are consistent and where there is divergence among models. Specific focus should be given to comparing and contrasting the literature with the conceptual model.

-

³⁶ The citation of specific sections of a plan or demonstration as useful examples is not intended as an endorsement of the entire document.

6. <u>TECHNICAL ANALYSES IN SUPPORT OF THE SECTION 179B</u> DEMONSTRATION

6.1. Determining the Appropriate Analytic Approach for the Section 179B Demonstration

As noted in Section 1.3 of this guidance, EPA recognizes that the relationship between certain NAAQS exceedances and associated international transport is clearer in some cases than in others. Table 1 details analyses recommended for all demonstrations, and additional analyses for demonstrations needing additional lines of evidence.

While each section 179B demonstration will involve a unique set of conditions, EPA believes that the general characteristics listed below would suggest the need for a demonstration with fewer lines of evidence to be appropriate:

- Affected monitors located near an international border.
- Large international emission sources located across the border near the affected monitors.
- Meteorology and international transport patterns connect emissions from identified international sources to monitors on days with monitored exceedances.
- Exceedances do not occur on days with similar conditions when transport to monitors is domestic in origin.

Conversely, the following characteristics would suggest the need for a more detailed demonstration with additional lines of evidence:

- Affected monitors not located near an international border.
- Specific international sources and/or their contributing emissions are not identified or are difficult to identify.
- Exceedances on internationally influenced days are in the range of typical exceedances attributable to local sources.
- Exceedances occurred in association with other processes and sources of pollutants, or on days where meteorological conditions were conducive to local pollutant formation (e.g., for O₃, clear skies and elevated temperatures).

Air quality in areas with proximity to borders and coasts often has clear and strong relationships with nearby international anthropogenic sources. Figure 3 shows the combined O₃ contributions (the fourth highest 8-hour daily maximum value within the U.S.) from anthropogenic emissions in Canada and Mexico based on source apportionment modeling results for 2023 (EPA, 2018b). This modeling estimates future aggregate international contributions of emissions from Canada and Mexico and is not necessarily representative of the international contributions that occur on specific days (e.g., specific exceedances that may influence a SIP determination and be critical for section 179B purposes) in a given area. Therefore, this type of analysis, by itself, would not be sufficient to demonstrate international contributions for the purposes of section 179B, but it may be helpful as part of a broader set of analyses used to support the weight of evidence in a section 179B demonstration.

In the 2023 modeling results discussed above, 24 of the 26 monitors with contributions greater than 5 ppb were located within 30 miles of the Canada or Mexico international border, and all 26 were within 40 miles of the border. Additionally, the influence of international sources on near-

border U.S. monitors appears to be substantially larger in locations with a large industrial source or urban center directly on the international side of the border.




Figure 3. O_3 (fourth highest maximum daily 8-hour average) contributions from in-domain Mexican and Canadian anthropogenic emissions at AQS monitors in 2023 projections. Monitors that are projected as nonattainment in 2023 are outlined squares and all other monitors are smaller circles.

Consistent with the general pattern illustrated in Figure 3, Figure 4 shows a strong exponential reduction of Mexico/Canada contribution as distance from the border increases. This modeling is only able to quantify emissions from within the contiguous U.S. modeling domain (i.e., "indomain"), which extends from 23 to 52 degrees north. Outside the domain, international shipping and transpacific international contributions could not be quantified in this modeling.




Figure 4. Mexico/Canada source contribution (S) in 2023 at monitors by distance from border (dx).

Impacts from other international sources (not from Canada, Mexico, or shipping) are related to longer-range transport efficiency, which is related to an area's elevation. Figure 5 shows the influence of O₃ and precursors from outside the modeling domain as represented by simulation of initial and lateral boundary conditions as a function of elevation at monitors more than 100 miles from the boundary. The boundary conditions do not distinguish between international anthropogenic and natural sources, but here serve as a proxy for efficiency of transport from the boundaries to the monitor. Although the relationship is not trivial (r²=0.55), it is driven by a cluster of low-elevation monitors and a scattering of high-elevation monitors. If the low-elevation monitors (<500 m) are excluded, the relationship significantly weakens (r²=0.08). To contrast, removing the monitors further from the border strengthens the relationship with distance from border. Proximity to the boundary alone is not sufficient to explain the variability in international contribution between sites. Instead, there is a complex relationship between location, local topography, elevation, and season.




Figure 5. Initial and Boundary Conditions contribution (ICBC) to projected 2023 design values as a function of elevation at AQS monitors that are farther than 100 miles from the border.

6.2. Analyses to Consider in Weight-of-Evidence Demonstrations

As with all intended section 179B demonstrations, the submitting air agency and its EPA Regional office should discuss the appropriate level of evidence before developing a demonstration. As a result of this discussion, EPA and the air agency can jointly identify the appropriate analytical approach for the section 179B demonstration.

There is no rigid set of rules regarding which specific analytical elements will demonstrate (or refute) the influence of international anthropogenic emissions. Each case is unique. Table 1 provides a checklist of recommended analyses that could support the demonstration of international anthropogenic influence. Each analysis is described in more detail in Section 6.3 of this guidance. In addition to these recommended analyses, other assessments not specifically mentioned in this guidance may also be valuable for certain areas. An approvable demonstration will generally contain a consistent analytical narrative that shows international anthropogenic emissions meaningfully contributed to an exceedance at the monitor.

Table 1. Recommended analyses to demonstrate international contribution for all demonstrations, with specific analyses listed for demonstrations needing additional lines of evidence. All analyses should be selected in consultation with the EPA Regional office.

Type of Analysis	Description of Analyses
Conceptual Model (Section 5)	Description and basic supporting information regarding the conditions that lead to high pollutant concentrations in the area and what conditions are conducive to international influence.
Ambient Observational Analysis (Section 6.3.1)	 5 years (or more) of peak concentration data with internationally influenced days flagged. Table with percentile ranks of days for international influence. Relationship between high levels of pollution (i.e., criteria or precursor pollutants) and meteorological/air quality conditions characteristic of international transport. Back trajectories and/or backward dispersion analyses (or where trajectory/dispersion analyses are not available, in-situ wind roses) showing internationally influenced source-receptor relationships on exceedance days Summary narrative including a comparison of locally and internationally influenced days. Demonstrations needing additional lines of evidence: Historical diurnal profile comparison on local and non-local contribution days.
Comprehensive Emissions Analysis (Section 6.3.2)	 Develop a domestic emission inventory. Obtain or develop an international emission inventory for proximate sources. Projections of both inventories to the relevant year (in some cases it may be helpful to project prior-year inventories even for historical analyses). Demonstrations needing additional lines of evidence: Obtain or develop a broader international emission inventory.

Type of Analysis	Description of Analyses
	Reference existing modeling that has been evaluated, typically for a related analysis.
Modeling to Quantify International Contribution (Section 6.3.3)	 Demonstrations needing additional lines of evidence (choose one in consultation with the Region): Dispersion modeling quantifying international contribution and culpability (for inert pollutants such as direct PM_{2.5}) Photochemical modeling quantifying contribution via sensitivity modeling or source apportionment. (for chemically reactive pollutants such as O₃ or secondary PM_{2.5})
Chemical Fingerprint Analysis (aka "receptor modeling" or "filter analysis") (Section 6.3.4)	Where available and applicable (see section 6.3.4 of this guidance).

6.3. Analyses to Demonstrate International Contribution

After determining the appropriate analytical approach for a section 179B demonstration, the next step in establishing a relationship between international emissions and a monitored exceedance is to develop the analyses that describe how international emissions were transported to the monitor in sufficient quantities to cause the exceedance. The analyses necessary to establish this relationship should be tailored to the area and pollutant of interest, and should generally include pollutant measurements, meteorological observations, emissions quantification, and air quality modeling. The 1994 Addendum to the General Preamble of the Implementation of Title I of the Clean Air Act Amendments of 1990 (59 FR 42001) suggested a set of five analyses that may help support section 179B demonstrations (a, d) for PM₁₀:

- 1. Place several ambient PM-10 monitors and a meteorological station; measuring wind speed and direction, in the U.S. nonattainment area near the international border.³⁷ Evaluate and quantify any changes in monitored PM-10 concentrations with a change in the predominant wind direction.
- 2. Comprehensively inventory PM-10 emissions within the U.S. in the vicinity of the nonattainment area and demonstrate that the impact of those sources, after application of reasonably available controls, does not cause the NAAQS to be exceeded. This analysis must include an influx of background PM-10 in the area. Background PM-10 levels could be based on concentrations measured in a similar area not influenced by emissions from outside the U.S.

_

³⁷ See 40 CFR part 58 for guidance on locating PM-10 monitors and "On-site Meteorological Program Guidance for Regulatory Modeling Applications" (EPA, 1987) for guidance on locating meteorological stations.

- 3. Analyze ambient sample filters for specific types of particles emanating from across the border. (Although not required, characteristics of emissions from sources may be helpful).
- 4. Inventory the sources on both sides of the border and compare the magnitude of PM-10 emissions originating within the U.S. to those emanating from outside the U.S.
- 5. Perform air dispersion and/or receptor modeling to quantify the relative impacts on the nonattainment area of sources located within the U.S., and of foreign sources of PM_{10} emissions (this approach combines information collected from the international emission inventory and meteorological stations, ambient monitoring network, and analysis of filters).

With the exception of item 3 above, these methods, with minor modifications, are also generally applicable to other criteria pollutants, including carbon monoxide and O₃ which are explicitly addressed in section 179B(b)–(d). For O₃ and potentially PM_{2.5}, the type of modeling performed will be chemical transport and not dispersion. Again, air agencies can describe the mechanics of the international transport in a variety of ways. Although the specific analyses appropriate for a section 179B demonstration will vary on a case-by-case basis, these methods are generally aligned with the five factors considered in the present-day designations process: air quality data, emissions and emissions-related data, meteorological data, geography/topography, and jurisdictional boundaries.³⁸

This guidance on analyses is structured into four types of supporting analyses. Section 6.3.1 of this guidance discusses measured air quality data analyses. Section 6.3.2 of this guidance describes comprehensive emissions analyses. Section 6.3.3 characterizes source-receptor modeling analyses. Section 6.3.4 of this guidance highlights receptor modeling analyses. The examples provided in each section focus on PM_{2.5} and O₃, which EPA anticipates will be of wider interest than CO and PM₁₀. However, the examples may be applicable to other criteria pollutants.

6.3.1. Measured Air Quality Data Analysis

The first component of establishing a relationship between international anthropogenic emissions and the monitored exceedance is to prepare an analysis showing how the measured concentration compares to the distribution or time series of historical concentrations measured at the same monitor and/or at other monitors in the area. Air agencies can show the relationship between the days with internationally influenced concentration(s) and historical concentrations across all days in a variety of ways. Table 2 describes example analyses that could be completed to show

29

³⁸ Memorandum to Regional Administrators entitled "Area Designations for the 2015 Ozone National Ambient Air Quality Standards" February 2016 at: https://www.epa.gov/sites/production/files/2016-02/documents/ozone-designations-guidance-2015.pdf.

whether the international-influenced exceedances were outside the bounds of generally expected pollutant levels.

Table 2. Suggested Analyses for Comparing Historical and Internationally Influenced Concentrations

Historical Concentration Evidence	Types of Analyses/Supporting Information
1. NAAQS concentration data	Plot the NAAQS concentration matrix at the affected monitor(s) for the most recent 5-year period ³⁹ that includes the international transport influence(s) of regulatory significance. This can also be supplemented with a table that briefly describes percentile ranks of internationally influenced days with a comparison against historical means and maxima.
2. Identify transport influences	Distinguish any high concentrations associated with previously approved international transport demonstrations, suspected international transport, and other unusual occurrences (e.g., exceptional events) from other high pollution days that are primarily due to normal domestic emissions (provide evidence to support the identification when possible).

6.3.1.1. Demonstration Days and Historical Context

A brief overview of the measured concentration data and the transport patterns that governed the suspected international influence should be provided near the beginning of a section 179B demonstration as part of the conceptual model. For illustrative purposes, Figure 6 shows an example time-series plot that highlights (in red) the days that are the focus of a hypothetical demonstration and puts them in the context of other O₃ days. In this case, 2015 and 2016, which are the focus of the evaluation of international sources, do not fully cover the 5-year data period. Therefore, there could be other historical days (i.e., a subset of the days in blue) that also have international influence. Note that Figure 6 shows that days being evaluated for international influence are not unlike other exceedance days in the historical record. An example of tabular day identification is shown in Table 3 that could complement a figure like Figure 6.

30

³⁹ Section 8.4.2.e of appendix W (82 FR 5182, January 17, 2017) recommends using 5 years of adequately representative meteorology data from the National Weather Service (NWS) to ensure that worst-case meteorological conditions are represented. Similarly, for international influence purposes, EPA believes that 5 years of ambient air data better represent the range of "normal" air quality than do shorter periods.




Figure 6. Measured O_3 as a function of day-of-year with example days highlighted as influenced by international sources marked in red and other days marked in green.

Table 3. Example tabular summary of exceedance days for internationally influenced O₃ data.

Transport Day	Concentration
	(ppm)
5/30/2015	0.071
6/8/2015	0.076
3/1/2016	0.074
4/19/2016	0.072
5/13/2016	0.072
6/28/2016	0.072
7/5/2016	0.071
7/12/2016	0.075

6.3.1.2. Establishing an International Source-Receptor Relationship

Measured exceedances should be connected to international source emissions by meteorological analysis. Fleming, Monks, and Manning (2012) published a particularly relevant review paper highlighting in-situ wind direction, trajectory models, particle dispersion models, and chemical transport models. The two most widely used analyses in section 179B demonstrations are in-situ wind analyses and trajectory analyses. Fleming, Monks, and Manning (2012) highlight that more sophisticated methods have largely replaced local wind direction as a method of identifying air mass history. Further, they note that dispersion models provide a more useful field for composition

analysis. Trajectory analyses rely on the use of meteorological models and their interpretation of observed conditions (aka reanalysis), which provides meteorological information over a broad area, but introduces biases inherent in the meteorological model. For example, previous section 179B demonstration submissions have highlighted that reanalysis may be too coarse to properly represent wind flow patterns in locations with complex topography. HYSPLIT now includes several 12 km resolved products and a 3 km resolved product from the NOAA's High-Resolution Rapid Refresh system (HRRR).⁴⁰ The HRRR product and others like it will help future demonstrations that require fine resolution for complex topography. Despite their respective limitations, winds and trajectory analyses can provide useful information as part of the weight-of-evidence in a section 179B demonstration.

<u>In-situ Winds Analyses</u>: In-situ winds analyses infer sources of pollution based on the speed and direction from which the wind blows. The strength of the method is in its simplicity to perform, but the simplicity is achieved by assuming that locally measured winds are representative of wind directions between the source and monitor. One weakness of this approach is the known existence of meandering or circular wind patterns. Wind analyses should only be used when more sophisticated methods are precluded. If included in a demonstration, in-situ wind analyses will hold less weight in the evidence assessment.

In-situ wind analysis was used in the section 179B demonstration for the Nogales, Arizona, PM₁₀ nonattainment area (ADEQ, 2012). That demonstration suggested that local topography limited the application of trajectory analyses. Each section 179B demonstration should review the increasingly accurate and fine-scale meteorology data available at the time. When opting out of trajectory or dispersion modeling, the air agency should still compare trajectory or dispersion modeling to its proposed technique. In the case of Nogales, the demonstration instead included analysis of measured local winds. Table 4 and Figure 7 highlight that wind direction on low concentration days was disproportionately from the north and west, and that high concentration days were disproportionately from the southerly direction with lower wind speeds. By itself, this wind vector analysis is merely suggestive. It is possible that meteorology conducive for transport from international sources is associated with southerly winds, or southerly winds may indicate a larger-scale flow pattern that connects this monitor to other domestic sources. Therefore, additional analyses are needed to help confirm conclusions from the wind vector analysis.

_

⁴⁰ Although the web interface only has July 2019 forward, the University of Utah HRRR archive (https://rapidrefresh.noaa.gov/hrrr/) has been rebuilt going back as far as July 2016. HYSPLIT provides tools for creating HYSPLIT compatible meteorological inputs from the available grib2 file (https://www.ready.noaa.gov/documents/Tutorial/html/meteo cnvrt.html).

*Table 4. Nogales, Arizona Hourly ambient PM10 concentrations sorted by concentration and wind direction, 2007–2009 exceedance days. (ADEQ, 2012, p. Appendix A Table 11)*⁴¹

	Range of Ambient Concentration Values (microgram/m³)							
Wind Direction Quadrant	< 150	150 - 250	250 - 350	350 - 450	450 - 550	>= 550	Share of All Wind Direction Observations	
Northerly NW to NNE	27%	6%	3%	3%	3%	0%	17%	
Easterly NE to ESE	15%	16%	16%	11%	3%	8%	14%	
Southerly SE to WSW	41%	71%	72%	84%	92%	92%	57%	
Westerly SW to WNW	18%	6%	8%	3%	3%	0%	12%	
Total	100%	100%	100%	100%	100%	100%	100%	




Figure 7. Nogales, Arizona PM10 nonattainment area: wind roses for non-exceedance and exceedance days (ADEQ, 2012, pp. Appendix A, Figure 9)

<u>Trajectory Analyses</u>: Backward trajectory analyses promote a fuller understanding of transport between sources and measured concentrations. A trajectory analysis follows an air mass represented by a single point as it moves through space due to atmospheric forces, and a backward trajectory retraces the path a particle would have taken to arrive at a point. The location of trajectories can be paired with additional meteorological data, geopolitical boundaries, and emission inventories to help distinguish between the influence of domestic and international emissions.


33

⁴¹ The citation of specific sections of a plan or demonstration as useful examples is not intended as an endorsement of the entire document.

The examples shown in Figure 8 and Figure 9 present results of HYSPLIT trajectories to assess the potential international and domestic impacts in the Imperial County, California, area. 42 Trajectories were initialized every 2 hours through each 8-hour maximum O₃ daily value (from start, dt=0,2,4,6,8 hours) from 2015-01-01 to 2016-08-24 using North American Mesoscale Forecast System archived on the Sigma coordinate (NAMS) data downloaded from the HYSPLIT archive. Each initialization uses three altitudes: 100, 500, and 1000 meters. The three altitudes help to characterize air throughout the PBL. Ideally, each altitude release would be paired with hourly trajectories (8 trajectories/day for O₃, 24 trajectories/day for PM_{2.5}) instead of the five used here. The trajectories used in these examples were also configured to include PBL height as an output along with coordinates. The altitude of the trajectory can be compared to the PBL to identify when a trajectory is likely to interact with emissions. If the PBL is not known, using a low assumed PBL may be appropriate. The residence time of trajectories within the PBL or below a fixed level can qualitatively be considered a potential emissions sensitivity (PES, f_x). The sensitivity can then be combined with emission inventories to develop an expected potential source contribution (PSC, f_xE_x). Because PES and PSC do not account for pollutant production and loss processes, the results should not be interpreted as absolute contributions and may be most useful in section 179B demonstrations when normalized. (For example, $NPES_x = PES_x$ / ΣPES_x ; and $NPSC_x = PSC_x / \Sigma PSC_x$). The normalized values show the expected relative contributions.

Figure 8 simply compares trajectories from (a) all exceedance day trajectories; (b) days that are primarily U.S. influenced; and (c) days that are primarily Mexican influenced. Trajectories are rarely conclusive on their own because they are individually subject to model configuration artifacts and represent probable trajectories. When a large majority (e.g., 75 percent) of trajectories pass over an international source region of interest with known emissions (here Mexicali), this is part of a weight of evidence that an international contribution exists on that day. When the international contribution is more prevalent on exceedance days than other days, that reinforces the weight of evidence.

⁴² Trajectories were developed to support this guidance document and have not been used in a submission.



(a) All Exceedance Days




Figure 8. El Centro 72-hour back trajectories for (a) all exceedance days, (b) an example primarily domestically influenced trajectory, and (c) an example more internationally influenced exceedance day.

Figure 9 shows back trajectory-based NPES (a,b) and NPSC (c,d). The percent attributed to various regions (NPES: U.S., Mexico, or Ocean; NPSC: U.S., Mexico, or Natural) is shown in the figure panel titles. Comparing NPES panels (a) and (b) reveals that exceedance day trajectories spent relatively more time in the Mexican PBL than non-exceedance days (24)

percent vs. 17 percent), but also show a density of PBL time along the U.S. Pacific coast. This identifies both international and domestic sources on exceedance days. Comparing NPSC panels (c,d) to NPES (a,b) shows that emissions across the border play a bigger role than suggested by just looking at trajectories. Like NPES, the NPSC attributed to Mexico is greater on exceedance days (52 percent) than all days (39 percent).

One challenge associated with using backward trajectories is that a large number of trajectories is necessary to create a PES or PSC. The NPES and NPSC in Figure 9 are relatively sparse and currently include 23 days with three altitudes and five releases (345 trajectories). Further, the trajectory could pass near a source and the narrow nature of the trajectory would not include the source. For this reason, trajectory-based NPES and NPSC can only provide supporting evidence. When exceedance days show larger fractions of NPSC from international anthropogenic sources, this adds to the weight of evidence that international anthropogenic sources contribute to exceedances.




Figure 9. Imperial County, California O₃ nonattainment area: El Centro 72-hour back trajectory Normalized Potential Emission Sensitivity maps (a,b) and Normalized Potential Source

Contribution maps (c,d) for all days (a,c) and exceedance days (b,d). The frequency is shown by the color and identifies the fraction of hourly locations from all trajectories that are in each cell. The USA, MEX, and OCN (ocean) labels identify the percentage of NPES within each region. The USA, MEX and NAT labels identify the percentage of NPSC from regional sources.

For areas farther from international sources, the HYSPLIT analysis will provide less direct evidence and can be used only to assess general transport patterns. Back trajectory analysis can provide information about the transport patterns, including the time spent in the local boundary layer, the extent of vertical transport, and the transport time to upwind areas. The time spent in the boundary layer will often be inversely related to international impact, while the vertical transport connects the parcel to regions of the troposphere with efficient transport. This type of information can be provided to support a section 179B demonstration. Fleming, Monks, and Manning (2012) discusses several applications to long-range transport that include cluster analysis. Cluster analysis may be useful for demonstrating similarity between types of days and, therefore, to distinguish between days of varying international influence.

<u>Backward Dispersion</u>: Backward dispersion analyses follow plumes rather than single-point air masses in backward trajectories. Backward dispersion results can improve upon trajectory analyses in several ways. First, backward dispersion includes the effects of turbulent motion. Each trajectory is no longer defined by just its average pathway, but instead is described by a probability distribution of particles. Second, the backward dispersion model uses a continuous release so that the backward trajectory includes many different initial conditions. The backward dispersion creates a "retroplume" that is proportional to residence time (Seibert and Frank 2004; 10.5194/acp-4-51-2004). The retroplume does not account for all processes and is subject to certain assumptions (Lin et al. 2003; 10.1175/BAMS-D-14-00110.1). When assumptions are satisfied, the retroplume can be converted to a NPES and NPSC (similar residence time from back trajectories).


The example shown in Figure 10 is based on HYSPLIT backward dispersion simulations. Similar to the back-trajectory section (above), simulations were configured based on maximum daily 8-hour O₃ measurements on 587 days at the El Centro monitor. For backward dispersion, source locations were configured to release a vertical line-source of a generic gas tracer between 100 and 1000 m over the monitor. The release lasted for the full 8-hour period. Compared to a series of backward trajectories, the 8 hours of emissions provide a more continuous view of possible sources, the dispersion model accounts for turbulent motion, and explicitly weights closer sources more heavily. In the case of secondarily-formed O₃ and PM_{2.5}, the weighting toward closer sources may overestimate their influence. In an ideal case, these receptor-oriented backward dispersion results could be combined with source-oriented forward dispersion results to produce a more complete source-receptor relationship.⁴³


Figure 10 shows the backward dispersion NPES and NPSC. Compared to backward trajectories (see Figure 9), the backward dispersion NPES and NPSC have a smoother distribution that results from turbulent dispersion. The percent attributed to various regions (NPES: U.S., Mexico, or Ocean; NPSC: U.S., Mexico, or Natural) is also similar to backward trajectories. A key difference is that the dispersion-based continuous surface lends itself to creating day-specific

_

⁴³ https://ready.arl.noaa.gov/documents/Tutorial/html/src recp.html.

NPES and NPSC in a way that trajectories do not. Figure 10 also shows two days that illustrate primarily U.S.-influenced (Figure 10e) and primarily Mexico-influenced (Figure 10f) exceedances. Despite the quantitative results, there are many uncertainties and the results depend upon the configuration and choices made in the processing. When a large majority of NPSC are from a source region (here Mexico), these results could be part of a weight of evidence that these days have international influence. When the fraction of NPSC is substantially larger on exceedance days than typical days, this strengthens the weight of evidence.






(a) All days NPES


Points below 1000m - NAT: 7%; US: 51%; MEX: 42%; 10⁻² 10⁻³ 10⁻⁴ (⁵⁵₂₀ ⁴₂₀) ⁵⁶₂₀ (⁵⁵₂₀ ⁵⁶₂₀) ⁵⁶₂₀ (⁵⁵₂₀ (⁵⁵₂₀) ⁵⁶₂₀ (⁵⁵₂₀) ⁵⁶₂₀ (⁵⁵₂₀) ⁵⁶


(b) Exceedance days NPES



(c) All days NPSC

(d) Exceedance days NPSC





(e) Primarily U.S. Influenced Exceedance

(f) Primarily Mexico Influenced Exceedance

Figure 10. El Centro 72-hour back dispersion-based NPES (a,b) and NPSC (c-f) for all days (a,c), exceedance days (b,d), a primarily U.S. influenced exceedance (e) and a primarily Mexico influenced exceedance (f).

6.3.2. Comprehensive Emission Analysis

A comprehensive emissions analysis is an important component of a section 179B demonstration. The emissions analysis provides most compelling direct evidence when the international emissions are immediately abutting the nonattainment area. In this case, the emissions analysis, including domestic and international emissions, will cover a geographic region of similar size and proximity to a metropolitan area. Emissions within the region would be separated into international and domestic components. A comprehensive emissions analysis should consider:

- What emission sources currently exist, and what is the magnitude of domestic versus international emissions?
- What controls are in place currently for the international sources (where available)?
- What change in emissions is expected in the foreseeable future (where available)?
- Are there international agreements that are already addressing these emissions?

Domestic and international emissions inventories should be developed in a manner consistent with EPA's emission inventory guidance (EPA, 2017). Domestic emissions should have been developed by the state during the SIP development process and should conform to that guidance. Not all emission guidance that is relevant to the domestic emissions is practical for international sources. The international emissions database may either be developed as a part of the section 179B demonstration or leveraged from a pre-existing public database. In either case, the process for developing the database should be based on the same principles of emission inventory development as are included in the guidance for domestic emissions. When using a pre-existing database, the provenance and methodology used to build the international emissions database should be well documented. Ideally, the database should also be compared to well-established databases (e.g., EDGAR, EDGAR-HTAP, CEDS).

A prospective attainment demonstration is based on future emissions estimates for international and domestic sources. Thus, on-the-books emissions controls should be considered when projecting future air quality. In some cases, it may be difficult to forecast international emission changes. The section 179B demonstration should document a good faith effort to account for expected international emissions growth and emission reduction measures. If there are significant uncertainties in international emissions estimates, it may be useful to provide a range of estimates.

A Nogales, Arizona, PM₁₀ section 179B demonstration (ADEQ, 2012)⁴⁴ illustrates the value of comparing domestic and international emissions for metropolitan areas. Nogales is the name of two cities (one in Arizona and one in Mexico) that share a border. The Nogales nonattainment area included only the U.S. side. The majority of the population and PM₁₀ emissions, however, occur on the Mexican side of the border. Table 5 shows an example PM₁₀ emissions from the Nogales nonattainment area, a range of emissions estimates from the cross-border municipality, and a range of percent contribution to the combined airshed emissions. ADEQ quantified emissions bounding the years (2008-2011) of exceedances being evaluated. The results in Table 5 are based on aggregated estimates for several broad source categories (not shown), including point, agricultural, residential wood combustion, waste burning, construction, onroad mobile, and nonroad sources.

Table 5. Annual Emission Inventories for Nogales, Arizona, and Nogales, Sonora, Mexico. Adapted from ADEQ (2012) Appendix A Clean Air Act Section 179B Attainment Determination

for the Nogales, Arizona, PM₁₀ Nonattainment Area; Tables 6-9.

	200) 8	2011		
	PM ₁₀ (tons)	Percent (%)	PM ₁₀ (tons)	Percent (%)	
Nogales NAA, Arizona	1,531	18-36.1	1,528	17.1-35.7	
Nogales Municipality, Mexico	[2,713-6,987]	[63.9-82]	[2,757-7,420]	[64.3-82.9]	
Total	4,244-8,518	100	4,285-8,948	100	

When the emissions inventory shows large emissions in a nearby international metropolitan area (large both in total and relative to local emissions), this supports a weight of evidence that international emissions are contributing to exceedances.

6.3.3. Modeling to Quantify International Contribution

Using air pollution modeling techniques—such as chemical transport models or dispersion models—is the most complete way to estimate the contribution of international emissions to monitors exceeding the NAAQS. In some cases, sufficient modeling may be readily available from the relevant SIP or past EPA analyses. The key factors for determining which modeling technique to use include: proximity of the nonattainment area to the emissions source or source region, and whether the NAAQS exceedances at the monitor are the result of emissions that react in the atmosphere (such as O₃ or secondarily formed PM_{2.5}) or primary emissions (such as direct

40

⁴⁴ The citation of specific sections of a plan or demonstration as useful examples is not intended as an endorsement of the entire document.

emissions of PM_{2.5}). Air agencies should consult with their EPA Regional office to determine the need for and applicability of modeling techniques.

6.3.3.1. Chemical Transport Modeling

Chemical Transport Modeling (CTM) is the preferred approach for quantifying international contribution for pollutants with a secondary component (such as O₃ and PM_{2.5}, which are formed, at least in part, as a result of photochemical reactions of precursor gases in the atmosphere). CTMs are necessary for quantifying long-range and secondary pollutant international contributions, and Baker & Kelly (2014) showed that CTMs can also reliably estimate single-source impacts relatively near the source. Thus, CTM application is not limited to regional-scale or national-scale international source contribution estimates. For more information on modeling single sources of secondarily formed pollutants see EPA (2016b) section 2.2. EPA has released guidance on performing CTM simulations (EPA, 2018c), which will be referred to hereafter as the modeling guidance. The applications described in the modeling guidance focus on SIP demonstration modeling, which is most directly applicable to the section 179B(a) demonstration. For a section 179B(a) demonstration, the observations and modeling should be consistent with the recommendations for SIP modeling. For a section 179B(b)–(d) demonstration, the observations and modeling would be from the attainment period rather than the designation period used in SIP modeling. The attainment period is the design value period used to determine whether the area attained by the applicable attainment date, while the designation period is the design value period used to designate the area as nonattainment. For either type of demonstration, the technical approach is similar.

Simulations to support section 179B demonstrations benefit from day-specific and aggregate analyses. Model performance evaluation often focuses on aggregate statistics, but may also include day-specific evaluation and analysis. This is particularly useful for section 179B demonstrations where the high international days may not align seasonally with other high concentration days. Ideally, the modeling episode would cover all the relevant observations (designation or attainment periods), but can use a representative modeling year to evaluate sensitivity to sources (domestic and international). When using a surrogate year, a demonstration should include an analysis to examine the impact of year-specific meteorology and transport patterns. The analysis to establish year-specific similarity will depend on the specific situation, but may include modeling, back trajectory cluster analyses, or other non-modeling analyses. The choice of modeling episodes to perform will depend upon the conceptual model and should be determined in coordination with the EPA Regional office.

Contributions from international sources can be estimated using various techniques described in the modeling guidance. The modeling guidance describes "brute-force" sensitivity modeling (i.e., rerunning the model simulation with emissions that have either been reduced or zeroed out from sources of interest) and source apportionment modeling that can be used to identify contributions from specific types of sources or source sectors, including international emissions. It will be useful to provide day-specific results as both absolute model contributions and relative contributions (Relative Contribution = Observation * Absolute Contribution / Total Prediction). To summarize the results, the contribution estimates may be applied to the design value from the designation or attainment period in a relative manner that is consistent with the Relative

Reduction Factor approach (RRF). For an example, see the Imperial County section 179B(b) demonstration for the 2008 O₃ standard (CARB, 2018).⁴⁵ This combination of contribution estimates will provide context for variability and for the magnitude of corrections applied via relative attribution.

Any modeling analysis for evaluating long-range impacts of international emissions will typically include a global simulation and a regional simulation. The global component is often used to supply boundary conditions to the regional simulation. To the extent practical, the two scales should use consistent meteorology, vertical resolution, emissions and representation of chemical species. The need for consistency is particularly true for the international emissions. It is also recommended to use consistent gas-phase chemical mechanisms and aerosol physics/chemistry components when conducting analyses with both global and regional simulations. Maintaining this consistency is critical when conducting analyses that zero-out anthropogenic emissions or rely on source apportionment modeling that targets sources present in both modeling scales.

Before estimating source contributions, the base case simulation 46 should be able to reasonably reproduce historical exceedances and typical differences between internationally influenced days and other days. For example, if internationally influenced days (identified by meteorology or trajectory analysis) are observed to have a higher concentration than on other days, then the base case should similarly predict higher concentrations on those days. This relationship is important because international contribution is often considered in a fractional sense. As a fraction, the denominator is total O₃. If the total O₃ on internationally influenced days is biased low, that could be due to international or local sources. Predicting the right pattern related to the international contribution does not guarantee accurate fractional prediction, but accuracy improves confidence in model results. Thus, the air agency should conduct both a model performance evaluation and diagnostic evaluation as described in the modeling guidance. These evaluations will provide confidence that variability in international contributions is reasonably well represented.

After the base case has been evaluated and shown capable of representing observations on internationally influenced days, then quantification of the international sources can commence. The choice of base case should match the application. For prospective demonstrations, the modeling should use the same base case and future year, consistent with the SIP modeling. For retrospective demonstrations, the modeling should focus on years used in the attainment evaluation. Quantifying the impact from international sources, as previously stated, may be done using a combination of sensitivity and/or source apportionment model runs. Depending on the scale of the analysis, this work may require coordinated efforts between a global and regional simulation. When this is the case, it is especially important to understand the consistency of inventories developed at the different scales. Regardless of the technique used for quantification of source contributions (e.g., sensitivity or source apportionment modeling), the most important aspect is appropriate implementation, discussed next.

⁴⁶ Base case is used to denote a simulation of a historical year that represents a baseline period (e.g., design value period) before emission projections and hypothetical controls are applied to project a future year air quality state.

⁴⁵ The citation of specific sections of a plan or demonstration as useful examples is not intended as an endorsement of the entire document.

The appropriate implementation of both sensitivity and source apportionment simulations begins with identifying the sources to be quantified. The sources to be quantified should be consistent with those described in the conceptual model. If the conceptual model links proximate international sources (e.g., for near-border receptors), then sensitivity runs would include perturbations to just these emissions and source apportionment should isolate the influence of these emissions as a separate tag. When the emissions cross scales (global to regional), the emissions should be consistent between the two scales and the sensitivity or source apportionment modeling must also cross scales. For sensitivity analyses, the boundary conditions for the regional perturbation simulation would be provided by a consistent perturbation in the global model similar to what has been done for U.S. Background (Emery, et al., 2012; Dolwick, et al., 2015; Zhang, et al., 2011). For international anthropogenic assessments that cross scales, examples are available in HTAP/AQMEII (Im, et al., 2018), the updated modeling for regional haze (EPA, 2019a), and EPA's most recent Ozone Policy Assessment (EPA, 2020). Most of these examples use zero-out or perturbation at multiple scales, but the regional haze application fused zero-out at hemispheric scale with source apportionment at regional scale. For either method, the appropriate emission perturbations are a critical step. Documentation on the hemispheric-scale perturbations for the two EPA applications is described in EPA (2019b).

In addition, the sensitivity or source apportionment modeling results should include an estimate of contribution from the U.S. for comparison. It may also be useful to separate the U.S. contribution into the nonattainment area's own state contribution and contributions from all other U.S. states. The collection of modeled results can be used to estimate the contribution of domestic and international anthropogenic sources.

Model contributions will be imperfect, and an estimate of a range should be considered and discussed in the context of the demonstration. Particularly for sensitivity modeling, the order of emission perturbations influences the result (zeroing the international source or the local source give different answers). Thus, at least two estimates of international source contribution should be developed and used to help characterize a range. Similarly, model attribution is often done using a relative approach (i.e., the Relative Reduction Factor or RRF). Photochemical modeling used in section 179B demonstrations use the RRF approach. Dolwick et al. (2015) showed that a relative approach for U.S. Background O₃ led to convergence in model results that had raw biases of differing sign. Similar convergence is expected for international contributions. Relative model application can mask problems when the model performance is poor due disproportionately to specific sources (e.g., missing stratosphere or oil & gas contributions). As a result, it is often useful to compare the raw results to the relative results because relative results drawn from poor performing days may lead to contribution estimates that are unreasonable. The range of results should demonstrate that international anthropogenic sources were large contributors relative to U.S. contributions on exceedance days.

Simulation results may be available from analyses conducted for a related regulatory program in lieu of developing modeling specific for the application. For example, EPA often performs source apportionment analyses with its modeling platform. EPA's 2011 modeling platform and the Western Air Quality Study ("WAQS") 2011 platform have both made source apportionment

modeling publicly available (EPA, 2016a). ⁴⁷ These model results include an estimate of future-year contribution from "in-domain" Mexico and Canada. If the conceptual model identifies near-source Mexican or Canadian sources as key contributors to nonattainment area exceedances, results from EPA modeling analyses may provide useful information in support of a prospective demonstration. For example, EPA's source apportionment results can be used as an upper bound estimate used to constrain adjustments proposed by trajectory analysis. EPA source apportionment results could also supplement state developed estimates to help characterize the credible range. In addition, the inputs for EPA's analysis provide a foundation for additional analyses that could be developed for a demonstration.

When results show that international contributions are larger on exceedance days and meaningfully larger than domestic contributions, the weight of evidence will be more compelling. The appropriateness of using pre-existing modeling for a section 179B demonstration should be discussed with the appropriate EPA Regional office.

6.3.3.2. Dispersion Modeling

Dispersion modeling is the preferred approach for quantifying contributions of near-monitor international emission sources of primary pollutants. For situations where the international contribution is from a single or group of industrial sources in close proximity (less than 50 km) to the impacted monitor, EPA has established several preferred dispersion models that can be used (The *Guideline on Air Quality Models*, i.e., Appendix W⁴⁸), depending on the application. In most applications, EPA's preferred near-field model, AERMOD will be used. When applied in a section 179B demonstration, a dispersion model should be applied with actual emissions from the international source(s) and modeled for the time period of the monitor design value calculation. For these applications, EPA's Appendix W offers guidance on many modeling inputs and procedures. Additionally, the SO₂ NAAQS Designations Modeling Technical Assistance Document (EPA, 2016c), provides recommendations on modeling domain, receptor placement, emissions inputs, meteorological data, and other inputs (EPA, 2018d). While the TAD is for SO₂, many of the recommendations would apply to other pollutants in section 179B demonstrations.

6.3.4. Receptor Modeling Analysis

This section describes receptor modeling and chemical finger-print approaches that are only applicable to identifying contributions from sources of particulate matter (or sources of pollutants that correlate well with particulate matter). Many tools are available for receptor modeling and discussed on the Support Center for Regulatory Atmospheric Modeling website (https://www.epa.gov/scram/air-pollutant-receptor-modeling). In an ideal case, there may be a unique tracer emission from a specific source across the border that can be used to identify international contribution. In more complex conditions, a Chemical Mass Balance or Positive

_

⁴⁷ Newer data can be made available upon request from the Ozone Policy Assessment. EPA: Policy Assessment for the Review of the Ozone National Ambient Air Quality Standards (EPA-452/R-20-001). U.S. Environmental Protection Agency, Research Triangle Park, NC, 2020b.

⁴⁸ Appendix W to 40 CFR Part 51, Guideline on Air Quality Models

Matrix Factorization may be used to identify international influence. With either approach, the connection of the receptor modeling to specific international sources will be critical.

6.3.4.1. Unique Tracer Analysis

A unique tracer analysis will need to include evidence that the tracer was emitted from the identified international source and that there are no sources of the tracer on the U.S. side of the border. Further, the tracer will need to be measurable above the detection limit in measurements made at or near the exceeding monitor.

The tracer choice will be extremely specific to the specific source and geographic area, but it should have certain properties. It should have a proportional relationship to total emissions of the pollutant and precursor potential from the source of interest. It should also be either chemically inert or have a lifetime much longer than the time spent in transport from the source to the monitor. If both of these properties are met, then the measured tracer concentration on exceeding days should be proportional to the international source contribution on exceeding days.

An analysis of tracer and international source contribution should be done on both internationally influenced exceedance days and days believed to be influenced by local sources (see Section 3.4.4 of this guidance). The contribution should be clearly larger on international transport days to demonstrate that there are not local sources of the tracer. Then, the contribution on exceeding, internationally influenced days would be subtracted from measured total to isolate the non-international contributions. If the non-international contribution is below the level of the NAAQS and the international (by difference) is large, then this analysis would support a weight of evidence that international anthropogenic emissions caused the exceedance.

6.3.4.2. Chemical Mass Balance or Positive Matrix Factorization


Chemical mass balance (CMB) (Schauer & Cass, 2000; Schauer, et al., 1996; Watson, Chow, & Mathai, 1989; Watson, et al., 2015) and positive matrix factorization (PMF) (Aiken, et al., 2009; Lanz, et al., 2007; Larsen & Baker, 2003; Ulbrich, Canagaratna, Zhang, & Worsnop, 2009) are two examples of receptor modeling methods which can be used to estimate source contributions to particulate matter. Both techniques combine ambient observations of speciated particulate matter with a set of minimization equations to produce an estimate of the source or factor (which can be associated with a type of source) contribution to PM_{10} or $PM_{2.5}$.

Using these techniques in a section 179B demonstration necessitates a careful selection of source profiles (for CMB) or interpretation of factors (for PMF) to properly attribute which sources are international. For CMB, source profiles specific to or dominated by emission sources located outside of the U.S. need to be used along with domestic source profiles for proper attribution of the international contribution. For PMF, additional measurements such as wind direction may be used as input to strengthen the confidence that one of the factors is international in origin.

Like the unique tracer analysis, CMB or PMF should be performed on both exceedance days and days believed to be influenced primarily by local emissions. If a source or factor is identified that is solely or mostly international in origin, the contribution should be clearly larger on the days

identified to be dominated by international transport. The contribution on exceeding, internationally influenced days would then be subtracted from measured total to isolate the non-international contributions. If the non-international contribution is below the level of the NAAQS and the international (by difference) is large, then this analysis would support a weight of evidence that international anthropogenic emissions caused the exceedance.

PMF was applied for the Calexico-Ethel monitor in a 2014 section 179B demonstration. In their analysis, they identified industrial sources, secondary nitrate/sulfate, motor vehicle sources, airborne soil, and refuse burning. Refuse burning, which was attributed exclusively to Mexico, was identified by a high organic carbon, elemental carbon and chlorine signature. Figure 11 shows the results of that analysis pooled for 2010 to 2012 and for just the days during that period with high PM_{2.5} concentrations. The demonstration also showed that the individual days identified as international transport-influenced had high refuse concentrations.






Figure 11. Average source contributions between 2010 and 2012 (CARB, 2014, p. Figure B2)⁴⁹

Receptor modeling techniques are generally contrasted with dispersion/chemical transport models that use pollutant emissions rate estimates, meteorological transport, and chemical transformation mechanisms to estimate the contribution of each source to receptor concentrations. Receptor and dispersion/chemical transport models can be complementary in a section 179B demonstration, with each type having strengths that compensate for the weaknesses of the other.

6.4. Example Conclusion Statement in the Demonstration

A section 179B demonstration should begin with a conceptual model and follow with a demonstration that establishes a relationship between international anthropogenic emissions and the monitored exceedance(s) based on the weight of evidence. The demonstration includes

⁴⁹ The citation of specific sections of a plan or demonstration as useful examples is not intended as an endorsement of the entire document.

multiple lines of evidence and analyses such as those identified in Section 6 of this guidance and should conclude with a statement similar to the language below:

Based on the evidence, including comparisons and analyses, provided in section(s) [reference the relevant section(s)] of this demonstration with respect to ambient air concentrations measured at the [name of monitor] on [dates], [Air Agency Name] has established that the [area name] area [would attain (for section 179B(a) demonstrations) or would have attained (for section 179B(b)–(d) demonstrations) the [name of NAAQS] NAAQS by the relevant attainment date but for emissions emanating from outside the U.S.

7. Public Comment Process

In addition to providing a conceptual model and evidence of international anthropogenic emissions transport to the subject area in a demonstration, EPA encourages air agencies to conduct and document (in the demonstration) a public comment process for all section 179B demonstrations prior to submitting the demonstration to EPA. In addition to coordinating with their respective EPA Regional office throughout the development of any section 179B demonstrations, EPA also recommends that air agencies notify their respective EPA Regional office when the state public comment process begins. In the case of a section 179B(a) "prospective" demonstration, the public comment process would be documented as part of completeness requirements in the associated SIP. In the case of a section 179B(b)–(d) "retrospective" demonstration, the air agency would likely need to conduct a demonstration-specific public comment process to include in its stand-alone submission.

Documentation of a public comment process as part of a section 179B demonstration should include information about how the public comment process was publicized, such as newspaper listings, website postings, and/or places (e.g., library, agency office) where a hardcopy was available. EPA also recommends that air agencies include any comments received and the agency's responses to those public comments.

8. References

- ADEQ. (2012). 2012 State Implementation Plan. Nogales, AZ: Arizona Department of Environmental Quality. Retrieved from https://www.regulations.gov/docket?D=EPA-R09-OAR-2012-0458
- Aiken, A., Salcedo, D., Cubison, M., Huffman, J., DeCarlo, P., Ulbrich, I., . . . Laski. (2009). Mexico City aerosol analysis during MILAGRO using high resolution aerosol mass spectrometry at the urban supersite (TO) Part 1: Fine particle composition and organic source apportionment. *Atmospheric Chemistry and Physics*(9), 6633-6653.
- Baker, K. R., & Kelly, J. T. (2014, 10). Single source impacts estimated with photochemical model source sensitivity and apportionment approaches. *Atmospheric Environment*, 96, 266-274. doi:10.1016/j.atmosenv.2014.07.042
- Byrne, A. (2015). The 1979 Convention on Long-Range Transboundary Air Pollution: Assessing its Effectiveness as a Multilateral Environmental Regime after 35 Years. *Translational Environmental Law*, 37-67. doi:10.1017/S2047102514000296
- CARB. (2014). Imperial County 2013 State Implementation Plan for the 2006 24-Hour PM2.5 Moderate Nonattainment Area. California Air Resources Board.
- CARB. (2018). Imperial County Clean Air Act Section 179B(b) Retrospective Analysis for the 75 ppb 8-Hour Ozone Standard. California Air Resources Board.
- Dentener, F., Keating, T. J., & Akimoto, H. (2010). *Hemispheric transport of air pollution. Part A: Ozone and Particulate Matter*. Geneva: Economic Commission For Europe, United Nations.
- Dolwick, P., Akhtar, F., Baker, K. R., Possiel, N., Simon, H., & Tonnesen, G. (2015). Comparison of background ozone estimates over the western United States based on two separate model methodologies. *Atmospheric Environment*, 109, 282-296. doi:10.1016/j.atmosenv.2015.01.005
- Emery, C., Jung, J., Downey, N., Johnson, J., Jimenez, M., Yarwood, G., & Morris, R. (2012). Regional and global modeling estimates of policy relevant background ozone over the United States. *Atmospheric Environment*, 47, 206-217. doi:10.1016/j.atmosenv.2011.11.012
- EPA. (1987). On-site Meteorological Program Guidance for Regulatory Modeling Applications. RTP, NC: US EPA.
- EPA. (2005). Guideline on Air Quality Models, 40 CFR Part 51, Appendix W. U.S. Environmental Protection Agency, RTP, NC. Retrieved from https://www3.epa.gov/ttn/scram/guidance/guide/appw 17.pdf
- EPA. (2009, January 15). Approval and Promulgation of Air Quality Implementation Plans; Texas; Approval of the Section 110(a)(1) Maintenance Plan for the 1997 8-Hour Ozone Standard for El Paso County. *Federal Register*, 74, 2387-2392. Retrieved from https://www.federalregister.gov/d/E9-708
- EPA. (2014). *Policy Assessment for the Review of the Ozone National Ambient Air Quality Standards*. RTP, NC: US EPA. Retrieved May 2018, from https://www.epa.gov/naaqs/ozone-o3-standards-policy-assessments-current-review
- EPA. (2015). Implementation of the 2015 Primary Ozone NAAQS: Issues Associated with Background Ozone, White Paper for Discussion. RTP, NC: U.S. EPA. Retrieved from https://www.epa.gov/sites/production/files/2016-03/documents/whitepaper-bgo3-final.pdf

- EPA. (2016a). Air Quality Modeling Technical Support Document for the 2015 Ozone NAAQS

 Preliminary Interstate Transport Assessment. Office of Air Quality Planning and
 Standards. RTP, NC: US EPA. Retrieved from
 https://www.epa.gov/sites/production/files/201701/documents/aq_modeling_tsd_2015_o3_naaqs_preliminary_interstate_transport_assess
 men.pdf
- EPA. (2016b). Final Guidance on the Preparation of Exceptional Events Demonstrations for Wildfire Events that May Influence Ozone Concentrations. RTP, NC: US EPA. Retrieved 6 7, 2018, from https://www.epa.gov/air-quality-analysis/exceptional-events-rule-and-guidance
- EPA. (2016b). Guidance on the Use of Models for Assessing the Impacts of Emissions from Single Sources on the Secondarily Formed Pollutants: Ozone and PM2.5. RTP, NC: U.S. Environmental Protection Agency.
- EPA. (2016c). SO2 NAAQS Designations Modeling. RTP, NC: U.S. Environmental Protection Agency. Retrieved from https://www.epa.gov/sites/production/files/2016-06/documents/so2modelingtad.pdf
- EPA. (2017). Emissions Inventory Guidance for Implementation of Ozone and Particulate Matter National Ambient Air Quality Standards (NAAQS) and Regional Haze Regulations. RTP, NC: U.S. Environmental Protection Agency. Retrieved 6 7, 2018, from https://www.epa.gov/air-emissions-inventories/air-emissions-inventory-guidance-implementation-ozone-and-particulate
- EPA. (2018a). Guidance on the Preparation of Exceptional Events Demonstrations for Stratospheric Ozone Intrusions. RTP, NC: Environmental Protection Agency. Retrieved 4 22, 2019, from https://www.epa.gov/sites/production/files/2018-11/documents/exceptional events soi guidance 11-8-2018.pdf
- EPA. (2018b). Information on the Interstate Transport State Implementation Plan Submissions for the 2015 Ozone Nationaal Ambient Air Quality Standards under Clean Air Act Section 110(a)(2)(D)(i)(I). RTP, NC: Environmental Protection Agency. doi:https://www.epa.gov/sites/production/files/2018-03/documents/transport memo 03 27 18 1.pdf
- EPA. (2018c). *Modeling Guidance for Demonstrating Air Quality Goals for Ozone, PM2.5, and Regional Haze.* RTP, NC: Environmental Protection Agency. Retrieved 4 22, 2019, from https://www3.epa.gov/ttn/scram/guidance/guide/O3-PM-RH-Modeling_Guidance-2018.pdf
- EPA. (2018d). *User's guide for the AMS/EPA Regulatory Model (AERMOD)*. U.S. Environmental Protection Agency, RTP, NC. Retrieved from https://www3.epa.gov/ttn/scram/models/aermod/aermod_userguide.pdf
- EPA. (2019a). Technical Support Document for EPA's Updated 2028 Regional Haze Modeling. RTP, NC: United States Environmental Protection Agency. Retrieved from https://www3.epa.gov/ttn/scram/reports/Updated_2028_Regional_Haze_Modeling-TSD-2019.pdf
- EPA. (2019b). Preparation of Emissions Inventories for the Version 7.1 2016 Hemispheric Emissions Modeling Platform. Research Triangle Park, NC: U.S. Environmental Protection Agency. Retrieved from https://www.epa.gov/sites/production/files/2019-12/documents/2016fe_hemispheric_tsd.pdf

- EPA. (2020). Policy Assessment for the Review of the Ozone National Ambient Air Quality Standards. Research Triangle Park, NC: U.S. Environmental Protection Agency.
- Fleming, Z. L., Monks, P. S., & Manning, A. J. (2012). Review: Untangling the influence of airmass history in interpreting observed atmospheric composition. *Atmospheric Research*(104-105), 1-39. doi:10.1016/j.atmosres.2011.09.009
- Huang, M., Carmichael, G. R., Pierce, R. B., Jo, D. S., Park, R. J., Flemming, J., . . . Payne, V. H. (2017). Impact of intercontinental pollution transport on North American ozone air pollution: an HTAP phase 2 multi-model study. *Atmospheric Chemistry and Physics*, 17(9), 1680-7316. doi:10.5194/acp-17-5721-2017
- Im, U., Christensen, J., Geels, C., Hansen, K., Brandt, J., Solazzo, E., . . . Liu. (2018). Influence of anthropogenic emissions and boundary conditions on multi-model simulations of major air pollutants over Europe and North America in the framework of AQMEII3. *Atmospheric Chemistry and Physics*, 18, 8929–8952. doi:10.5194/acp-18-8929-2018
- Jaffe, D., Cooper, O., Fiore, A., Henderson, B., Tonneson, G., Russell, A., . . . Moore, T. (2018). Scientific assessment of background ozone over the U.S.: Implications for air quality management. *Elementa Science of the Anthropocene*(6), 56. doi:10.1525/elementa.309
- Lanz, V., Alfarra, M., Baltensperger, U., Buchmann, B., Hueglin, C., & Pervot, A. (2007). Source apportionment of submicron organic aerosols at an urban site by factor analytical modelling of aerosol mass spectra. *Atmospheric Chemistry and Physics*(7), 1503-1522.
- Larsen, R., & Baker, J. (2003). Source apportionment of polycyclic aromatic hydrocarbons in the urban atmosphere: A comparison of three models. *Environmental Science & Technology*(37), 1873-1881.
- Lin, M., Horowitz, L. W., Payton, R., Fiore, A. M., & Tonnesen, G. (2017). US surface ozone trends and extremes from 1980 to 2014: quantifying the roles of rising Asian emissions, domestic controls, wildfires, and climate. *Atmospheric Chemistry and Physics*, 17(4), 2943-2970. doi:10.5194/acp-17-2943-2017
- Nopmongcol, U., Liu, Z., Stoeckenius, T., & Yarwood, G. (2017, 08 24). Modeling intercontinental transport of ozone in North America with CAMx for the Air Quality Model Evaluation International Initiative (AQMEII) Phase 3. *Atmospheric Chemistry and Physics*, 1680-7324. doi:10.5194/acp-17-9931-2017
- NRC. (2010). Global sources of local pollution: an assessment of long-range transport of key air pollutants to and from the United States. (U. N. US National Research Council, Ed.) Washington, D.C., 978-0-309-14401-8: National Academies Press.
- Schauer, J., & Cass, G. (2000). Source apportionment of wintertime gas-phase and particle-phase air pollutants using organic compounds as tracers. *Environmental Science & Technology*, 30, 1821-1832.
- Schauer, J., Rogge, W., Hildemann, L., Maurek, M., Cass, G., & Simoneit, B. (1996). Source apportionment of airborne particulate matter using organic compounds as tracers. *Atmospheric Environment*(30), 3837-3855.
- TNRCC. (1994). REVISIONS TO THE STATE IMPLEMENTATION PLAN (SIP) FOR THE CONTROL OF OZONE AIR POLLUTION: SECTION 818 DEMONSTRATION FOR THE EL PASO NONATTAINMENT AREA. Austin, TX: Texas Natural Resource Conservation Commission. Retrieved from https://www.tceq.texas.gov/assets/public/implementation/air/sip/sipdocs/1994-09-ELP/sept94 818 el paso.pdf

- Ulbrich, I., Canagaratna, M., Zhang, Q., & Worsnop, D. J. (2009). Interpretation of organic components from Poistive Matrix Factorization of aerosol mass spectrometric data. *Atmospheric Chemistry and Physics*(9), 2891-2918.
- US Senate. (1993). *A Legislative history of the Clean Air Act Ammendments v1-6*. Washington D.C.: U.S. G.P.O. Retrieved from https://catalog.hathitrust.org/Record/006091060
- van der A, R. J., Mijling, B., Ding, J., Koukouli, M., Liu, F., Li, Q., . . . Theys, N. (2017). Cleaning up the air: effectiveness of air quality policy for SO2 and NOx emissions in China. *Atmospheric Chemistry and Physics*, 17(3), 1775-1789. doi:10.5194/acp-17-1775-2017
- Verstraeten, W. W., Neu, J. L., Williams, J. E., Bowman, K. W., Worden, J. R., & Folkert, B. K. (2015). Rapid increases in tropospheric ozone production and export from China. *Nature Geoscience*, 8(9), 690-695. doi:10.1038/ngeo2493
- Watson, J., Chow, J., & Mathai, C. (1989). Receptor models in air resources management: A summary of the APCA international specialty conference. *JAPCA*(39), 419-426.
- Watson, J., Chow, J., Lowenthal, D., Antony Chen, L., Shaw, S., Edgerton, E., & Blanchard, C. (2015). PM2.5 source apportionment with organic markers in the southeastern aerosol research and characterization (SEARCH) study. *Journal of the Air & Waste Management Association*(65), 1104-1118.
- Zaveri, R. A., Saylor, R. D., Peters, L. K., McNider, R., & Song, A. (1995, May 1). A model investigation of summertime diurnal ozone behavior in rural mountainous locations. *Atmospheric Environment*, 29(9), 1043-1065. doi:10.1016/1352-2310(94)00319-G
- Zhang, L., Jacob, D., Downey, N., Wood, D., Blewitt, D., Carouge, C., . . . Wang, Y. (2011). Improved estimate of the policy-relevant background ozone in the United States using the GEOS-Chem global model with 1/2° × 2/3° horizontal resolution over North America. *Atmospheric Environment*, 45, 6769-6776. doi:10.1016/j.atmosenv.2011.07.054

Appendix A. Statutory Text

§7509a. International border areas

(a) Implementation plans and revisions

Notwithstanding any other provision of law, an implementation plan or plan revision required under this chapter shall be approved by the Administrator if—

- (1) such plan or revision meets all the requirements applicable to it under the chapter other than a requirement that such plan or revision demonstrate attainment and maintenance of the relevant national ambient air quality standards by the attainment date specified under the applicable provision of this chapter, or in a regulation promulgated under such provision, and
- (2) the submitting State establishes to the satisfaction of the Administrator that the implementation plan of such State would be adequate to attain and maintain the relevant national ambient air quality standards by the attainment date specified under the applicable provision of this chapter, or in a regulation promulgated under such provision, but for emissions emanating from outside of the United States.

(b) Attainment of ozone levels

Notwithstanding any other provision of law, any State that establishes to the satisfaction of the Administrator that, with respect to an ozone nonattainment area in such State, such State would have attained the national ambient air quality standard for ozone by the applicable attainment date, but for emissions emanating from outside of the United States, shall not be subject to the provisions of section $7511(a)(2)^{50}$ or (5) of this title or section 7511d of this title.

(c) Attainment of carbon monoxide levels

Notwithstanding any other provision of law, any State that establishes to the satisfaction of the Administrator, with respect to a carbon monoxide nonattainment area in such State, that such State has attained the national ambient air quality standard for carbon monoxide by the applicable attainment date, but for emissions emanating from outside of the United States, shall not be subject to the provisions of section 7512(b)(2) or (9) of this title.

(d) Attainment of PM-10 levels

Notwithstanding any other provision of law, any State that establishes to the satisfaction of the Administrator that, with respect to a PM-10 nonattainment area in such State, such State would have attained the national ambient air quality standard for carbon monoxide by the applicable attainment date, but for emissions emanating from outside the United States, shall not be subject to the provisions of section 7513(b)(2) of this title.

⁵⁰ The statute contained an erroneous reference to section 7511(a)(2) instead of 7511(b)(2).

United States	Office of Air Quality Planning and Standards	Publication No. EPA-457/P-20-001F
Environmental Protection	Air Quality Policy Division	December 2020
Agency	Research Triangle Park, NC	
Agency	Research Triangle Park, NC	